

Plate 1

(Text pp. 41-44)

- | | |
|---|--|
| 1: <i>C. marmoreus</i>
Palawan, Philippines (85 mm) | 13: <i>C. bandanus</i> var.
Papua New Guinea (40 mm) |
| 2: <i>C. marmoreus</i>
Solomon Is. (68 mm) | 14: <i>C. bandanus</i> f. <i>equestris</i>
Moluccas (58 mm) |
| 3: <i>C. marmoreus</i>
Papua New Guinea (59 mm) | 15: <i>C. bandanus</i> f. <i>equestris</i>
Moluccas (60 mm) |
| 4: <i>C. marmoreus</i> ; holotype of <i>C. pseudomarmoreus</i>
Crosse
MNHN (51 mm) | 16: <i>C. bandanus</i> f. <i>vidua</i> ; syntype of <i>C. vidua</i> Reeve
Capul, Philippines; MHNG (72 mm) |
| 5: <i>C. marmoreus</i> var.
New Caledonia (65 mm) | 17: <i>C. bandanus</i> f. <i>vidua</i>
Palawan, Philippines (66 mm) |
| 6: <i>C. marmoreus</i> f. <i>suffusus</i>
New Caledonia (55 mm) | 18: <i>C. bandanus</i> f. <i>vidua</i>
Palawan, Philippines (49 mm) |
| 7: <i>C. marmoreus</i> f. <i>crosseanus</i>
New Caledonia (58 mm) | 19: <i>C. bandanus</i> f. <i>vidua</i> var.
Palawan, Philippines; coll. & photo Baer
(51 mm) |
| 8: <i>C. marmoreus</i> f. <i>crosseanus</i>
New Caledonia (51 mm) | 20: <i>C. bandanus</i> f. <i>vidua</i>
Cuyo Id., Philippines (60 mm) |
| 9: <i>C. marmoreus</i> var.
New Caledonia (49 mm) | 21: <i>C. bandanus</i> f. <i>vidua</i> ; transition to typical form
Palawan, Philippines (53 mm) |
| 10: <i>C. bandanus</i>
Zanzibar (64 mm) | 22: <i>C. bandanus</i> f. <i>vidua</i> ; transition to typical form
Philippines (46 mm) |
| 11: <i>C. bandanus</i>
Moluccas (79 mm) | 23: <i>C. bandanus</i> f. <i>vidua</i> ; transition to typical form
Palawan, Philippines (50 mm) |
| 12: <i>C. bandanus</i>
Albay, Philippines (76 mm) | |



Plate 2

(Text pp. 42-47)

- | | | | |
|-----|---|-----|---|
| 1: | <i>C. bandanus</i> f. <i>nigrescens</i>
W. Samoa (42 mm) | 11: | <i>C. araneosus araneosus</i>
S. India (98 mm) |
| 2: | <i>C. bandanus</i> f. <i>nigrescens</i>
W. Samoa (38 mm) | 12: | <i>C. araneosus araneosus</i>
S. India (72 mm) |
| 3: | <i>C. bandanus</i> f. <i>nigrescens</i>
Admiralty Is. (34 mm) | 13: | <i>C. araneosus nicobaricus</i>
Samar, Philippines (86 mm) |
| 4: | <i>C. bandanus</i> f. <i>nigrescens</i>
Solomon Is. (56 mm) | 14: | <i>C. araneosus nicobaricus</i>
Samar, Philippines (62 mm) |
| 5: | <i>C. nocturnus</i>
Moluccas (58 mm) | 15: | <i>C. araneosus nicobaricus</i>
Padang, W. Sumatra (54 mm) |
| 6: | <i>C. nocturnus</i>
Moluccas; ZMA (65 mm) | 16: | <i>C. distans</i>
Zanzibar (81 mm) |
| 7: | <i>C. nocturnus</i>
Moluccas; ZMA (70 mm) | 17: | <i>C. distans</i>
Tahiti (81 mm) |
| 8: | <i>C. nocturnus</i> [Lightfoot]; lectotype
MHNG; photo Kohn (56 mm) | 18: | <i>C. distans</i>
Seychelles (83 mm) |
| 9: | <i>C. araneosus araneosus</i>
Sri Lanka (56 mm) | 19: | <i>C. distans</i> ; juvenile
Philippines; coll. da Motta, SMNS (30 mm) |
| 10: | <i>C. araneosus araneosus</i>
S. India (76 mm) | | |



Plate 3

(Text pp. 47-51)

- | | | | |
|-----|---|-----|---|
| 1: | <i>C. imperialis</i>
Zanzibar (77 mm) | 13: | <i>C. imperialis</i> f. <i>fuscatus</i> ; lectotype of
<i>C. fuscatus</i> Born
NMW; photo Wawra (53 mm) |
| 2: | <i>C. imperialis</i>
Hawaii (67 mm) | 14: | <i>C. zonatus</i>
Maldives (47 mm) |
| 3: | <i>C. imperialis</i>
Philippines (79 mm) | 15: | <i>C. zonatus</i>
W. Thailand (56 mm) |
| 4: | <i>C. imperialis</i> f. <i>fuscatus</i>
Zanzibar (71 mm) | 16: | <i>C. zonatus</i>
Laccadives (68 mm) |
| 5: | <i>C. imperialis</i> f. <i>fuscatus</i>
Zanzibar (71 mm) | 17: | <i>C. zonatus</i>
W. Thailand; coll. da Motta, SMNS (56 mm) |
| 6: | <i>C. imperialis</i> f. <i>fuscatus</i>
Pemba Is., Tanzania (63 mm) | 18: | <i>C. biraghii</i> <i>biraghii</i> ; holotype of <i>Leptoconus biraghii</i>
G. Raybaudi Massilia; Somalia; SMNS (11 mm) |
| 7: | <i>C. imperialis</i> f. <i>fuscatus</i>
Mombasa, Kenya (63 mm) | 19: | <i>C. biraghii</i> <i>biraghii</i> ; holotype of <i>Leptoconus biraghii</i>
G. Raybaudi Massilia; apertural view |
| 8: | <i>C. imperialis</i> f. <i>fuscatus</i>
Tulear, Madagascar (52 mm) | 20: | <i>C. korni</i> G. Raybaudi Massilia; paratype
Somalia; coll. Baer (10 mm) |
| 9: | <i>C. imperialis</i> f. <i>fuscatus</i>
Zanzibar (58 mm) | 21: | <i>C. biraghii</i> <i>biraghii</i> var.
Somalia; coll. G. Raybaudi Massilia (11 mm) |
| 10: | <i>C. imperialis</i> f. <i>fuscatus</i>
Mauritiüs (63 mm) | 22: | <i>C. biraghii</i> <i>biraghii</i> var.
Somalia; coll. G. Raybaudi Massilia (11 mm) |
| 11: | <i>C. imperialis</i> f. <i>compactus</i>
Mombasa, Kenya (74 mm) | | |
| 12: | <i>C. imperialis</i> f. <i>fuscatus</i> ; figured by Kiener as
<i>C. viridulus</i>
Lamarck coll., MHNG; photo Dajoz (67 mm) | | |



Plate 4

(Text pp. 51-56)

- | | |
|--|--|
| 1: <i>C. lividus</i>
W. Thailand (43 mm) | 16: <i>C. moreleti</i>
Solomon Is. (35 mm) |
| 2: <i>C. lividus</i>
Queensland (42 mm) | 17: <i>C. moreleti</i>
Marquesas (50 mm) |
| 3: <i>C. lividus</i>
Tanzania (54 mm) | 18: <i>C. moreleti</i>
Marquesas (40 mm) |
| 4: <i>C. lividus</i>
Jeddah, Red Sea (31 mm) | 19: <i>C. caillaudii</i>
NMWC; photo Filmer (43 mm) |
| 5: <i>C. lividus</i>
S. India (62 mm) | 20: <i>C. caillaudii</i>
Mauritius; coll. da Motta, SMNS (34 mm) |
| 6: <i>C. sanguinolentus</i> Quoy & Gaimard; lectotype
New Guinea (22 mm) | 21: <i>C. caillaudii</i>
Saya de Malha, Mascarene Ridge; coll. Danila (45 mm) |
| 7: <i>C. sanguinolentus</i>
New Caledonia (39 mm) | 22: <i>C. caillaudii</i>
Réunion; MNHN (29 mm) |
| 8: <i>C. sanguinolentus</i>
Zanzibar (39 mm) | 23: <i>C. caillaudii</i>
Réunion; MNHN (29 mm) |
| 9: <i>C. sanguinolentus</i>
Jeddah (40 mm) | 24: <i>C. caillaudii</i> var.
Red Sea; coll. Sharabati, BMNH (39 mm) |
| 10: <i>C. sanguinolentus</i>
Aqaba, Red Sea (39 mm) | 25: <i>C. caillaudii</i> var.
Red Sea; coll. Sharabati, BMNH (45 mm) |
| 11: <i>C. sanguinolentus</i> var.
Marquesas (46 mm) | 26: <i>C. varius</i>
Solomon Is. (51 mm) |
| 12: <i>C. sanguinolentus</i> var.
Marquesas (61 mm) | 27: <i>C. varius</i>
Mozambique (46 mm) |
| 13: <i>C. sanguinolentus</i> var.
Tahiti (29 mm) | 28: <i>C. varius</i>
Zanzibar (43 mm) |
| 14: <i>C. moreleti</i>
Okinawa (41 mm) | |
| 15: <i>C. moreleti</i>
Hawaii (39 mm) | |




Plate 5

(Text pp. 56-59)

- | | |
|--|---|
| 1: <i>C. biliosus</i>
Rameswaram, India (56 mm) | 17: <i>C. exiguus</i>
New Caledonia (25 mm) |
| 2: <i>C. biliosus</i>
Sri Lanka (44 mm) | 18: <i>C. exiguus</i>
New Caledonia; MNHN (30 mm) |
| 3: <i>C. biliosus</i>
Madras, India (59 mm) | 19: <i>C. exiguus</i>
New Caledonia (24 mm) |
| 4: <i>C. biliosus</i>
Madras, India (53 mm) | 20: <i>C. exiguus</i>
New Caledonia (33 mm) |
| 5: <i>C. biliosus</i>
Natal, S. Africa (33 mm) | 21: <i>C. exiguus</i>
New Caledonia (25 mm) |
| 6: <i>C. biliosus</i>
Mozambique (32 mm) | 22: <i>C. exiguus</i>
New Caledonia (26 mm) |
| 7: <i>C. biliosus</i>
Karachi, Pakistan (38 mm) | 23: <i>C. exiguus</i>
New Caledonia (25 mm) |
| 8: <i>C. biliosus</i>
Madagascar (38 mm) | 24: <i>C. exiguus</i> f. <i>bougei</i>
New Caledonia; MNHN (22 mm) |
| 9: <i>C. biliosus</i>
Solomon Is. (31 mm) | 25: <i>C. exiguus</i> f. <i>bougei</i>
New Caledonia (19 mm) |
| 10: <i>C. biliosus</i>
Philippines (29 mm) | 26: <i>C. exiguus</i> Lamarck; lectotype
MHNG; photo Dajoz (18 mm) |
| 11: <i>C. biliosus</i>
Moluccas (44 mm) | 27: <i>C. exiguus</i> ; lectotype of <i>C. cabritii</i> Bernardi
New Caledonia; MNHN (20 mm) |
| 12: <i>C. biliosus</i>
Queensland (35 mm) | 28: <i>C. exiguus</i> ; lectotype of <i>C. optimus</i> Sowerby III
BMNH (23 mm) |
| 13: <i>C. biliosus</i>
Queensland (41 mm) | 29: <i>C. exiguus</i> ; lectotype of <i>C. taylorianus</i> E. A. Smith
BMNH (20 mm) |
| 14: <i>C. exiguus</i>
New Caledonia (43 mm) | 30: <i>C. exiguus</i> f. <i>bougei</i> ; lectotype of
<i>C. bougei</i> Sowerby III
New Caledonia; BMNH (21 mm) |
| 15: <i>C. exiguus</i>
New Caledonia (38 mm) | 31: <i>C. exiguus</i> ; syntype of <i>C. plumbeus</i> Reeve
New Caledonia; BMNH (25 mm) |
| 16: <i>C. exiguus</i>
New Caledonia; coll. da Motta, SMNS (47 mm) | |



Plate 6

(Text pp. 59-62)

- | | |
|--|--|
| 1: <i>C. boeticus</i>
Guam; coll. da Motta, SMNS (27 mm) | 20: <i>C. montillai</i> Röckel; paratype
Coron, Palawan, Philippines (20 mm) |
| 2: <i>C. boeticus</i>
Fiji; coll. da Motta, SMNS (24 mm) | 21: <i>C. pauperculus</i>
Japan (26 mm) |
| 3: <i>C. boeticus</i>
Fiji (22 mm) | 22: <i>C. pauperculus</i>
Japan (34 mm) |
| 4: <i>C. boeticus</i>
Flores, Indonesia; coll. Metzner (35 mm) | 23: <i>C. pauperculus</i>
Japan (24 mm) |
| 5: <i>C. boeticus</i>
Sulu Sea (21 mm) | 24: <i>C. balteatus</i>
Rodrigues, Indian Ocean (31 mm) |
| 6: <i>C. boeticus</i>
Solomon Is. (21 mm) | 25: <i>C. balteatus</i>
Réunion (29 mm) |
| 7: <i>C. boeticus</i>
Bohol, Philippines (23 mm) | 26: <i>C. balteatus</i>
Zanzibar (29 mm) |
| 8: <i>C. boeticus</i>
Halmahera, Moluccas (30 mm) | 27: <i>C. balteatus</i>
Fiji (33 mm) |
| 9: <i>C. boeticus</i>
Mozambique; coll. Ramalho (21 mm) | 28: <i>C. balteatus</i>
Mozambique (24 mm) |
| 10: <i>C. boeticus</i>
Mozambique; coll. Ramalho (29 mm) | 29: <i>C. balteatus</i> f. <i>cernicus</i>
St. Brandon, Indian Ocean (32 mm) |
| 11: <i>C. boeticus</i> f. <i>ruppellii</i>
Pangasinan, Philippines (27 mm) | 30: <i>C. balteatus</i> f. <i>cernicus</i>
St. Brandon, Indian Ocean (30 mm) |
| 12: <i>C. boeticus</i> f. <i>ruppellii</i>
Batangas Bay, Philippines (30 mm) | 31: <i>C. balteatus</i> f. <i>cernicus</i>
Réunion (27 mm) |
| 13: <i>C. boeticus</i> f. <i>ruppellii</i>
Indonesia (29 mm) | 32: <i>C. balteatus</i> f. <i>cernicus</i>
Réunion (26 mm) |
| 14: <i>C. boeticus</i> f. <i>nitidus</i>
Okinawa; coll. da Motta, SMNS (20 mm) | 33: <i>C. balteatus</i>
Saya de Malha, Mascarene Ridge; coll. Danila
(30 mm) |
| 15: <i>C. boeticus</i> Reeve; figured syntype
Philippines; BMNH (37 mm) | 34: <i>C. balteatus</i>
Seychelles (29 mm) |
| 16: <i>C. boeticus</i> ; figured syntype of <i>C. rivularis</i> Reeve
Moluccas; BMNH (33 mm) | 35: <i>C. balteatus</i> ; type of <i>C. pigmentatus</i> Adams & Reeve
BMNH (25 mm) |
| 17: <i>C. boeticus</i> ; figured type of <i>C. cerinus</i> Reeve
Philippines; BMNH (29 mm) | 36: <i>C. balteatus</i> ; holotype of <i>C. moussoni</i> Crosse
Seychelles; BMNH (28 mm) |
| 18: <i>C. boeticus</i> ; figured syntype of <i>C. lachrymosus</i>
Reeve BMNH (30 mm) | |
| 19: <i>C. montillai</i> Röckel; paratype
Coron, Palawan, Philippines (19 mm) | |



Plate 7

(Text pp. 62-68)

- | | |
|--|---|
| 1: <i>C. miliaris miliaris</i>
Queensland (27 mm) | 19: <i>C. encaustus</i>
Marquesas (30 mm) |
| 2: <i>C. miliaris miliaris</i>
Vanuatu (26 mm) | 20: <i>C. encaustus</i>
Marquesas (22 mm) |
| 3: <i>C. miliaris miliaris</i>
Queensland (26 mm) | 21: <i>C. encaustus</i>
Marquesas (26 mm) |
| 4: <i>C. miliaris miliaris</i>
Fiji (29 mm) | 22: <i>C. coronatus</i>
India (37 mm) |
| 5: <i>C. miliaris miliaris</i>
Queensland (37 mm) | 23: <i>C. coronatus</i>
India (40 mm) |
| 6: <i>C. miliaris miliaris</i>
Zanzibar (31 mm) | 24: <i>C. coronatus</i>
Seychelles (32 mm) |
| 7: <i>C. miliaris miliaris</i>
Zanzibar (31 mm) | 25: <i>C. coronatus</i>
Queensland (36 mm) |
| 8: <i>C. miliaris miliaris</i>
Zanzibar (35 mm) | 26: <i>C. coronatus</i>
Zanzibar (33 mm) |
| 9: <i>C. miliaris miliaris</i> f. <i>fulgetrum</i>
Taiwan (24 mm) | 27: <i>C. coronatus</i>
Tuamotu (28 mm) |
| 10: <i>C. miliaris miliaris</i> f. <i>fulgetrum</i>
Japan (22 mm) | 28: <i>C. coronatus</i>
Queensland (33 mm) |
| 11: <i>C. miliaris miliaris</i> f. <i>fulgetrum</i>
Philippines; coll. da Motta, SMNS (35 mm) | 29: <i>C. coronatus</i> f. <i>aristophanes</i>
Tahiti (24 mm) |
| 12: <i>C. miliaris pascuensis</i>
Easter Id. (22 mm) | 30: <i>C. coronatus</i> Gmelin; neotype
BMNH; photo Filmer (28 mm) |
| 13: <i>C. miliaris pascuensis</i>
Easter Id. (19 mm) | 31: <i>C. coronatus</i> f. <i>aristophanes</i> ; lectotype of
<i>C. aristophanes</i> Sowerby II
BMNH (35 mm) |
| 14: <i>C. miliaris pascuensis</i>
Easter Id. (30 mm) | 32: <i>C. taeniatus</i>
Muscat, Oman (33 mm) |
| 15: <i>C. abbreviatus</i>
Hawaii (36 mm) | 33: <i>C. taeniatus</i>
Muscat, Oman (33 mm) |
| 16: <i>C. abbreviatus</i>
Hawaii (26 mm) | 34: <i>C. taeniatus</i>
Red Sea (37 mm) |
| 17: <i>C. abbreviatus</i>
Hawaii (37 mm) | 35: <i>C. taeniatus</i>
Red Sea (36 mm) |
| 18: <i>C. encaustus</i>
Marquesas (33 mm) | |




Plate 8

(Text pp. 68-74)

- | | |
|--|--|
| 1: <i>C. sponsalis</i>
Palawan, Philippines (26 mm) | 19: <i>C. musicus</i> f. <i>ceylanensis</i>
W. Thailand; coll. Roussy (17 mm) |
| 2: <i>C. sponsalis</i>
Chagos Id. (25 mm) | 20: <i>C. musicus</i> f. <i>ceylanensis</i>
W. Thailand; coll. Roussy (19 mm) |
| 3: <i>C. sponsalis</i>
Samoa (20 mm) | 21: <i>C. musicus</i>
W. Thailand; coll. Roussy (19 mm) |
| 4: <i>C. sponsalis</i>
Tahiti (23 mm) | 22: <i>C. parvatus</i>
Egypt, Red Sea (23 mm) |
| 5: <i>C. sponsalis</i>
New Caledonia (20 mm) | 23: <i>C. parvatus</i>
Mozambique (20 mm) |
| 6: <i>C. sponsalis</i>
Vanuatu (20 mm) | 24: <i>C. parvatus</i>
Réunion (17 mm) |
| 7: <i>C. sponsalis</i> f. <i>nanus</i>
W. Australia (29 mm) | 25: <i>C. parvatus</i>
Madagascar (18 mm) |
| 8: <i>C. sponsalis</i> f. <i>nanus</i>
Queensland (23 mm) | 26: <i>C. parvatus</i>
Djibouti (16 mm) |
| 9: <i>C. musicus</i> Hwass in Bruguière; figure of lectotype
(18 mm) | 27: <i>C. ebraeus</i>
Samoa (38 mm) |
| 10: <i>C. musicus</i> f. <i>ceylanensis</i> ; original figure of <i>C. ceylanensis</i> Hwass in Bruguière (19 mm) | 28: <i>C. ebraeus</i>
Mauritius (41 mm) |
| 11: <i>C. musicus</i> f. <i>ceylanensis</i> ; neotype of <i>C. ceylanensis</i> Hwass in Bruguière
Sri Lanka; ZMA (10 mm) | 29: <i>C. ebraeus</i>
Zanzibar (39 mm) |
| 12: <i>C. musicus</i> f. <i>mighelsi</i> ; original figure of <i>C. mighelsi</i> Kiener
(23 mm) | 30: <i>C. ebraeus</i> ; juvenile
Solomon Is. (10 mm) |
| 13: <i>C. musicus</i>
Philippines (19 mm) | 31: <i>C. chaldaeus</i>
Cebaco Id., W. Panama (40 mm) |
| 14: <i>C. musicus</i>
New Caledonia; coll. da Motta, SMNS (19 mm) | 32: <i>C. chaldaeus</i>
Zanzibar (35 mm) |
| 15: <i>C. musicus</i>
Philippines (20 mm) | 33: <i>C. chaldaeus</i>
W. Thailand (27 mm) |
| 16: <i>C. musicus</i>
Philippines (20 mm) | 34: <i>C. dorreensis</i>
Onslow, N.W. Australia (35 mm) |
| 17: <i>C. musicus</i>
Okinawa (20 mm) | 35: <i>C. dorreensis</i>
Exmouth, N. W. Australia (29 mm) |
| 18: <i>C. musicus</i> f. <i>mighelsi</i>
W. Thailand; coll. Roussy (16 mm) | 36: <i>C. dorreensis</i>
Exmouth, N. W. Australia (29 mm) |




Plate 9

(Text pp. 74-79)

- | | |
|---|---|
| 1: <i>C. floridulus</i>
Cebu, Philippines (50 mm) | 18: <i>C. sazanka</i> ; holotype of <i>C. yoshioi</i> Azuma (with
periostracum)
Japan; coll. & photo Azuma (43 mm) |
| 2: <i>C. floridulus</i>
Cebu, Philippines (23 mm) | 19: <i>C. danilai</i> Röckel & Korn; holotype
Ras Fartak, Gulf of Aden; SMNS (31 mm) |
| 3: <i>C. floridulus</i>
Cebu, Philippines (27 mm) | 20: <i>C. danilai</i> Röckel & Korn; holotype
apertural view |
| 4: <i>C. floridulus</i>
Bohol, Philippines (44 mm) | 21: <i>C. danilai</i> Röckel & Korn; paratype
Ras Fartak, Gulf of Aden; coll. Danila (35 mm) |
| 5: <i>C. floridulus</i>
Bohol, Philippines (35 mm) | 22: <i>C. danilai</i>
Gulf of Aden; coll. Haksal (28 mm) |
| 6: <i>C. muriculatus</i> f. <i>sugillatus</i>
Marinduque, Philippines. (41 mm) | 23: <i>C. hamamotoi</i> Yoshiba & Koyama; holotype
Shiono-misaki Peninsula, Japan; WPMN; photo
Koyoma (23 mm) |
| 7: <i>C. muriculatus</i> f. <i>sugillatus</i>
Queensland (43 mm) | 24: <i>C. hamamotoi</i> Yoshiba & Koyama; holotype
apertural view |
| 8: <i>C. muriculatus</i>
Solomon Is. (14 mm) | 25: <i>C. hamamotoi</i> Yoshiba & Koyama; paratype
Shiono-misaki Peninsula, Japan; WPMN;
photo Koyama (22 mm) |
| 9: <i>C. muriculatus</i>
Solomon Is. (15 mm) | 26: <i>C. hamamotoi</i>
New Caledonia; MNHN (17 mm) |
| 10: <i>C. muriculatus</i>
Solomon Is. (15 mm) | 27: <i>C. hamamotoi</i>
New Caledonia; MNHN (24 mm) |
| 11: <i>C. muriculatus</i> f. <i>sugillatus</i>
Halmahera, Moluccas (36 mm) | 28: <i>C. axelrodi</i>
Palawan, Philippines (18 mm) |
| 12: <i>C. sazanka</i>
Japan (41 mm) | 29: <i>C. axelrodi</i>
Palawan, Philippines (21 mm) |
| 13: <i>C. sazanka</i>
Japan (32 mm) | 30: <i>C. axelrodi</i>
Palawan, Philippines (16 mm) |
| 14: <i>C. sazanka</i>
Japan (30 mm) | 31: <i>C. axelrodi</i>
Palawan, Philippines (15 mm) |
| 15: <i>C. sazanka</i>
Japan (32 mm) | |
| 16: <i>C. sazanka</i>
Bohol, Philippines (26 mm) | |
| 17: <i>C. sazanka</i>
Bohol, Philippines; coll. Korn (55 mm) | |



Plate 10

(Text pp. 79-83)

- | | |
|--|--|
| 1: <i>C. litteratus</i>
Zanzibar (84 mm) | 12: <i>C. eburneus</i>
Kantang, W. Thailand (52 mm) |
| 2: <i>C. litteratus</i>
Philippines (100 mm) | 13: <i>C. eburneus</i>
Samar, Philippines (45 mm) |
| 3: <i>C. litteratus</i>
Philippines (115 mm) | 14: <i>C. eburneus</i> f. <i>crassus</i>
Halmahera, Moluccas (39 mm) |
| 4: <i>C. litteratus</i>
Mozambique (75 mm) | 15: <i>C. eburneus</i> f. <i>crassus</i>
Papua New Guinea (33 mm) |
| 5: <i>C. leopardus</i>
Maldives (90 mm) | 16: <i>C. eburneus</i> f. <i>polyglotta</i>
Bicol Region, Philippines (42 mm) |
| 6: <i>C. leopardus</i>
Vanuatu (96 mm) | 17: <i>C. eburneus</i> f. <i>polyglotta</i>
Bicol Region, Philippines (34 mm) |
| 7: <i>C. leopardus</i> ; subadult
Mozambique (53 mm) | 18: <i>C. eburneus</i> f. <i>polyglotta</i>
Bicol Region, Philippines (33 mm) |
| 8: <i>C. eburneus</i>
E. Malaysia (48 mm) | 19: <i>C. eburneus</i> f. <i>polyglotta</i>
Bicol Region, Philippines (31 mm) |
| 9: <i>C. eburneus</i>
Philippines (47 mm) | 20: <i>C. augur</i>
Zanzibar (56 mm) |
| 10: <i>C. eburneus</i>
Rabaul, Papua New Guinea (55 mm) | 21: <i>C. augur</i>
Madagascar (59 mm) |
| 11: <i>C. eburneus</i>
Palawan, Philippines (41 mm) | 22: <i>C. augur</i>
Mozambique (61 mm) |



Plate 11

(Text pp. 83-88)

- | | |
|--|--|
| 1: <i>C. arenatus</i>
W. Thailand (64 mm) | 14: <i>C. pulicarius vaultieri</i>
Marquesas (64 mm) |
| 2: <i>C. arenatus</i> f. <i>bizona</i>
Kenya (53 mm) | 15: <i>C. stercusmuscarum</i>
Solomon Is. (45 mm) |
| 3: <i>C. arenatus</i> f. <i>aequipunctatus</i>
Massawa, Eritrea (53 mm) | 16: <i>C. stercusmuscarum</i>
Solomon Is. (55 mm) |
| 4: <i>C. arenatus</i>
Tanzania (42 mm) | 17: <i>C. stercusmuscarum</i>
Marinduque, Philippines (46 mm) |
| 5: <i>C. arenatus</i>
Egypt, Red Sea (72 mm) | 18: <i>C. stercusmuscarum</i>
Bohol, Philippines (51 mm) |
| 6: <i>C. arenatus</i>
Mozambique; coll. Amorim (40 mm) | 19: <i>C. zeylanicus</i>
Rameswaram, India (50 mm) |
| 7: <i>C. arenatus</i>
Solomon Is. (34 mm) | 20: <i>C. zeylanicus</i>
Sri Lanka (51 mm) |
| 8: <i>C. pulicarius pulicarius</i>
New Caledonia (52 mm) | 21: <i>C. zeylanicus</i>
Rameswaram, India (61 mm) |
| 9: <i>C. pulicarius pulicarius</i>
Solomon Is. (50 mm) | 22: <i>C. zeylanicus</i>
Zanzibar (64 mm) |
| 10: <i>C. pulicarius pulicarius</i>
Tahiti (43 mm) | 23: <i>C. zeylanicus</i> ; subadult
Ranong, W. Thailand (39 mm) |
| 11: <i>C. pulicarius pulicarius</i>
Philippines (45 mm) | 24: <i>C. characteristicus</i>
Mactán, Philippines (55 mm) |
| 12: <i>C. pulicarius pulicarius</i>
Taiwan (64 mm) | 25: <i>C. characteristicus</i>
Madras, India (51 mm) |
| 13: <i>C. pulicarius vaultieri</i>
Marquesas (46 mm) | 26: <i>C. characteristicus</i>
Philippines (80 mm) |



Plate 12

(Text pp. 88-92)

- | | |
|--|---|
| 1: <i>C. suturatus suturatus</i>
Langford Reef, Queensland (34 mm) | 16: <i>C. tessulatus</i>
Perlas Is., Panama (48 mm) |
| 2: <i>C. suturatus suturatus</i>
Low Id., Queensland (35 mm) | 17: <i>C. tessulatus</i>
Wakayama, Japan (48 mm) |
| 3: <i>C. suturatus suturatus</i>
Bushy Id., Queensland (41 mm) | 18: <i>C. tessulatus</i>
Muscat, Oman (50 mm) |
| 4: <i>C. suturatus suturatus</i>
Bushy Id., Queensland (45 mm) | 19: <i>C. tessulatus</i>
Palawan, Philippines (40 mm) |
| 5: <i>C. suturatus suturatus</i>
N.W. Australia; WAM (25 mm) | 20: <i>C. tessulatus</i>
E. Malaysia (34 mm) |
| 6: <i>C. suturatus suturatus</i>
Pt. Hedland, W. Australia; AMS (21 mm) | 21: <i>C. tessulatus</i>
India (67 mm) |
| 7: <i>C. suturatus sandwichensis</i>
Hawaii (44 mm) | 22: <i>C. melvilli</i> Sowerby III; holotype
NMWC (19 mm) |
| 8: <i>C. suturatus sandwichensis</i>
Hawaii (34 mm) | 23: <i>C. melvilli</i>
Oman; ZMA (23 mm) |
| 9: <i>C. kiicumulus</i>
Wakayama, Japan (32 mm) | 24: <i>C. melvilli</i>
Oman; ZMA (26 mm) |
| 10: <i>C. kiicumulus</i>
Wakayama, Japan (36 mm) | 25: <i>C. melvilli</i>
Oman (19 mm) |
| 11: <i>C. kiicumulus</i>
Kushimoto, Japan (36 mm) | 26: <i>C. melvilli</i>
Oman (20 mm) |
| 12: <i>C. kiicumulus</i> ; subadult
Amami-o-shima, Japan (20 mm) | 27: <i>C. tuticorinensis</i> Röckel & Korn; paratype
Tuticorin, India (27 mm) |
| 13: <i>C. tessulatus</i>
Batangas Bay, Philippines (44 mm) | 28: <i>C. tuticorinensis</i>
Tuticorin, India (29 mm) |
| 14: <i>C. tessulatus</i>
Aqaba, Red Sea (44 mm) | 29: <i>C. tuticorinensis</i>
Tuticorin, India (26 mm) |
| 15: <i>C. tessulatus</i>
Madras, India (46 mm) | |



Plate 13

(Text pp. 92-96)

- | | |
|--|--|
| 1: <i>C. suratensis</i>
Lingayen Gulf, Philippines (97 mm) | 14: <i>C. figulinus</i>
Brunei, Borneo (55 mm) |
| 2: <i>C. suratensis</i>
Cuddalore, India (70 mm) | 15: <i>C. figulinus</i>
Madagascar (70 mm) |
| 3: <i>C. suratensis</i> ; juvenile
Palawan, Philippines (42 mm) | 16: <i>C. figulinus</i>
Rameswaram, India (66 mm) |
| 4: <i>C. betulinus</i>
Cuddalore, India (90 mm) | 17: <i>C. lorisii</i>
Sulu Sea, Philippines (68 mm) |
| 5: <i>C. betulinus</i>
Madras, India (90 mm) | 18: <i>C. lorisii</i>
Cuddalore, India (85 mm) |
| 6: <i>C. betulinus</i>
Tuléar, Madagascar (67 mm) | 19: <i>C. lorisii</i> f. <i>insignis</i> (with periostracum)
Solomon Is. (70 mm) |
| 7: <i>C. betulinus</i>
Pamsan, India (68 mm) | 20: <i>C. lorisii</i> f. <i>insignis</i>
Solomon Is. (72 mm) |
| 8: <i>C. betulinus</i>
Cuddalore, India (67 mm) | 21: <i>C. lorisii</i> f. <i>insignis</i>
Lingayen Gulf, Philippines (80 mm) |
| 9: <i>C. betulinus</i> var. (? <i>C. betulinus</i> × <i>C. figulinus</i>)
Tuléar, Madagascar (67 mm) | 22: <i>C. lorisii</i> f. <i>insignis</i>
Cuddalore, India (64 mm) |
| 10: <i>C. betulinus</i> var. (? <i>C. betulinus</i> × <i>C. figulinus</i>)
Madras, India (55 mm) | 23: <i>C. lorisii</i> f. <i>huberorum</i> ; paratype of <i>C. huberorum</i>
da Motta
Visakhapatnam, India (45 mm) |
| 11: <i>C. betulinus</i> ; holotype of <i>C. zulu</i> Petuch
Mozambique; DMNH (63 mm) | 24: <i>C. lorisii</i> f. <i>huberorum</i>
Mahapallipuram, India; coll. Stabenow (66 mm) |
| 12: <i>C. figulinus</i>
Mozambique (83 mm) | 25: <i>C. lorisii</i> f. <i>huberorum</i>
Mahapallipuram, India; coll. Stabenow (73 mm) |
| 13: <i>C. figulinus</i>
Somalia (67 mm) | |



Plate 14

(Text pp. 96-100)

- 1: *C. glaucus*
Solomon Is. (43 mm)
- 2: *C. glaucus*
Borneo (51 mm)
- 3: *C. glaucus*
Solomon Is. (52 mm)
- 4: *C. quercinus*
Mombasa, Kenya (65 mm)
- 5: *C. quercinus*
Queensland (83 mm)
- 6: *C. quercinus*; white form
Madras, India (89 mm)
- 7: *C. quercinus*; juvenile
Oahu, Hawaii (20 mm)
- 8: *C. hyaena hyaena*
Java (47 mm)
- 9: *C. hyaena hyaena*
Bombay, India (51 mm)
- 10: *C. hyaena hyaena*
Brunei, Borneo (39 mm)
- 11: *C. hyaena hyaena*
Kilakarai, India (73 mm)
- 12: *C. hyaena hyaena*
Madras, India (73 mm)

- 13: *C. hyaena hyaena*
Java (52 mm)
- 14: *C. hyaena hyaena*
N.W. Borneo (48 mm)
- 15: *C. hyaena hyaena*
Java (50 mm)
- 16: *C. hyaena hyaena*
Bombay, India (44 mm)
- 17: *C. hyaena hyaena*
Java (43 mm)
- 18: *C. hyaena hyaena*
Bombay, India (46 mm)
- 19: *C. hyaena concolor*
Solomon Is. (50 mm)
- 20: *C. hyaena concolor*
Solomon Is. (53 mm)
- 21: *C. hyaena concolor*
Solomon Is. (55 mm)
- 22: *C. gilvus*
Solomon Is. (30 mm)
- 23: *C. gilvus*
Solomon Is. (32 mm)
- 24: *C. gilvus*
Solomon Is. (25 mm)



Plate 15

(Text pp. 100-105)

- | | |
|--|---|
| 1: <i>C. achatinus</i>
Calicut, India (66 mm) | 18: <i>C. catus</i>
Zanzibar (38 mm) |
| 2: <i>C. achatinus</i>
Bombay, India (53 mm) | 19: <i>C. catus</i>
Queensland (34 mm) |
| 3: <i>C. achatinus</i>
Roebourne, W. Australia (53 mm) | 20: <i>C. catus</i>
Samoa (38 mm) |
| 4: <i>C. achatinus</i>
Darwin, N. Australia (52 mm) | 21: <i>C. catus</i>
W. Thailand (35 mm) |
| 5: <i>C. achatinus</i>
Muscat, Oman (44 mm) | 22: <i>C. catus</i>
Seychelles (36 mm) |
| 6: <i>C. achatinus</i>
Calicut, India (50 mm) | 23: <i>C. catus</i>
Marquesas (30 mm) |
| 7: <i>C. achatinus</i>
Solomon Is. (44 mm) | 24: <i>C. catus</i> f. <i>nigropunctatus</i>
Mozambique (36 mm) |
| 8: <i>C. achatinus</i>
Oman (47 mm) | 25: <i>C. catus</i> f. <i>nigropunctatus</i>
Egypt, Red Sea (55 mm) |
| 9: <i>C. monachus</i>
Solomon Is. (47 mm) | 26: <i>C. catus</i> f. <i>nigropunctatus</i>
Muscat, Oman (32 mm) |
| 10: <i>C. monachus</i>
Solomon Is. (46 mm) | 27: <i>C. catus</i> f. <i>nigropunctatus</i>
Muscat, Oman (30 mm) |
| 11: <i>C. monachus</i>
Davao, Philippines (44 mm) | 28: <i>C. catus</i> f. <i>morrisoni</i>
Ashmore Reef, W. Australia; coll. Dan (26 mm) |
| 12: <i>C. monachus</i>
Davao, Philippines (45 mm) | 29: <i>C. fischoederi</i> Röckel & da Motta; paratype
Coron, Palawan, Philippines (41 mm) |
| 13: <i>C. monachus</i> var.; holotype of <i>C. contusus</i> Reeve
"Moluccas"; BMNH (33 mm) | 30: <i>C. fischoederi</i> Röckel & da Motta; paratype
as 29; apertural view |
| 14: <i>C. monachus</i> var.
Papua New Guinea; coll. Richards (27 mm) | 31: <i>C. fischoederi</i> Röckel & da Motta; paratype
Coron, Palawan, Philippines (37 mm) |
| 15: <i>C. striolatus</i> f. <i>decurtata</i>
Singapore (32 mm) | 32: <i>C. fischoederi</i>
Philippines; coll. da Motta, SMNS (49 mm) |
| 16: <i>C. striolatus</i> f. <i>decurtata</i>
Philippines (32 mm) | 33: <i>C. fischoederi</i>
? Albany, W. Australia; coll. da Motta, SMNS
(39 mm) |
| 17: <i>C. striolatus</i>
Papua New Guinea (35 mm) | |



Plate 16

(Text pp. 105-109)

- | | |
|---|---|
| 1: <i>C. fulmen</i>
Japan (47 mm) | 17: <i>C. bruuni</i>
New Caledonia; MNHN (47 mm) |
| 2: <i>C. fulmen</i> f. <i>kirai</i>
Japan (57 mm) | 18: <i>C. bruuni</i>
Raoul Id., Kermadec Is.; coll. Haksal (47 mm) |
| 3: <i>C. fulmen</i> f. <i>kirai</i>
Taiwan (56 mm) | 19: <i>C. bruuni</i>
Kermadec Is.; coll. & photo Baer (41 mm) |
| 4: <i>C. fulmen</i> (with periostracum)
Japan (50 mm) | 20: <i>C. bruuni</i>
New Caledonia; coll. Haksal (40 mm) |
| 5: <i>C. kinoshitai</i>
Bohol, Philippines (69 mm) | 21: <i>C. bruuni</i>
New Caledonia; MNHN (43 mm) |
| 6: <i>C. kinoshitai</i> (with periostracum)
Bohol, Philippines (46 mm) | 22: <i>C. circumcisis</i>
Moluccas (77 mm) |
| 7: <i>C. kinoshitai</i>
Bohol, Philippines (78 mm) | 23: <i>C. circumcisis</i>
Bohol, Philippines (70 mm) |
| 8: <i>C. kinoshitai</i>
Bohol, Philippines (83 mm) | 24: <i>C. circumcisis</i> f. <i>brazieri</i>
Cebu, Philippines (70 mm) |
| 9: <i>C. kinoshitai</i>
Bohol, Philippines; coll. Baer (62 mm) | 25: <i>C. circumcisis</i> ; subadult
Bohol, Philippines (49 mm) |
| 10: <i>C. kinoshitai</i>
Bohol, Philippines (54 mm) | 26: <i>C. circumcisis</i> f. <i>brazieri</i>
Solomon Is. (81 mm) |
| 11: <i>C. kinoshitai</i>
Philippines; ZMA (57 mm) | 27: ? <i>C. circumcisis</i>
Tahiti; coll. & photo Baer (56 mm) |
| 12: <i>C. kinoshitai</i> f. <i>tamikoe</i>
N.E. Taiwan; photo Lan (76 mm) | 28: <i>C. aurisiacus</i>
Moluccas (63 mm) |
| 13: <i>C. kinoshitai</i> ; holotype of <i>C. brontodes</i> Shikama
Taiwan; NSMT (48 mm) | 29: <i>C. aurisiacus</i>
Sulawesi, Indonesia; coll. Schmid (47 mm) |
| 14: <i>C. kinoshitai</i> f. <i>calliginosus</i>
Philippines (57 mm) | 30: <i>C. aurisiacus</i>
Sulawesi, Indonesia (47 mm) |
| 15: <i>C. kinoshitai</i> f. <i>tamikoe</i>
Philippines; coll. Schmid (61 mm) | 31: <i>C. aurisiacus</i>
Sulawesi, Indonesia; coll. Schmid (52 mm) |
| 16: <i>C. bruuni</i>
New Caledonia; coll. Haksal (40 mm) | 32: <i>C. aurisiacus</i>
Sulawesi, Indonesia; coll. Schmid (59 mm) |



Plate 17

(Text pp. 109-112)

- | | |
|---|--|
| 1: <i>C. timorensis</i>
Mauritius (39 mm) | 15: <i>C. neptunus</i> ; juvenile
Bohol, Philippines (22 mm) |
| 2: <i>C. timorensis</i>
St. Brandon, Indian Ocean; coll. & photo
Baer (38 mm) | 16: <i>C. neptunus</i> ; holotype of <i>C. neptunoides</i> E. A. Smith
"Australia"; BMNH (43 mm) |
| 3: <i>C. timorensis</i>
St. Brandon, Indian Ocean (37 mm) | 17: <i>C. neptunus</i> f. <i>colorvariegatus</i>
Balut Id., Philippines (62 mm) |
| 4: <i>C. nimbosus</i>
Vanuatu (43 mm) | 18: <i>C. neptunus</i> f. <i>colorvariegatus</i>
Balut Id., Philippines (63 mm) |
| 5: <i>C. nimbosus</i>
Mozambique (41 mm) | 19: <i>C. neptunus</i> f. <i>colorvariegatus</i> ; juvenile
Philippines; coll. da Motta, SMNS (23 mm) |
| 6: <i>C. nimbosus</i>
Bangka Str., Indonesia (41 mm) | 20: <i>C. lienardi</i>
New Caledonia (45 mm) |
| 7: <i>C. nimbosus</i>
Mozambique; coll. Amorim (41 mm) | 21: <i>C. lienardi</i>
New Caledonia; coll. & photo Baer (43 mm) |
| 8: <i>C. nimbosus</i>
Mozambique (40 mm) | 22: <i>C. lienardi</i>
New Caledonia (43 mm) |
| 9: <i>C. janus</i>
Mozambique; coll. Amorim (74 mm) | 23: <i>C. lienardi</i>
New Caledonia; MNHN (58 mm) |
| 10: <i>C. janus</i>
Madagascar (60 mm) | 24: <i>C. lienardi</i>
New Caledonia (36 mm) |
| 11: <i>C. janus</i>
Madagascar (64 mm) | 25: <i>C. lienardi</i>
Solomon Is.; coll. G. Raybaudi Massilia (40 mm) |
| 12: <i>C. janus</i>
Madagascar (62 mm) | 26: <i>C. lienardi</i>
Iloilo, Philippines; coll. da Motta, SMNS
(30 mm) |
| 13: <i>C. neptunus</i>
Cebu, Philippines (69 mm) | 27: <i>C. lienardi</i> ; holotype of <i>C. prevosti</i> Sowerby III
New Caledonia; BMNH (40 mm) |
| 14: <i>C. neptunus</i>
Balut Id., Philippines (69 mm) | |



Plate 18

(Text pp. 112-117)

- | | |
|--|---|
| 1: <i>C. radiatus</i>
Cebu, Philippines (68 mm) | 18: <i>C. ochroleucus tmetus</i>
Vanuatu (54 mm) |
| 2: <i>C. radiatus</i>
Taiwan (77 mm) | 19: <i>C. ochroleucus tmetus</i>
Solomon Is. (51 mm) |
| 3: <i>C. radiatus</i>
Fiji (49 mm) | 20: <i>C. ochroleucus tmetus</i> ; juvenile; holotype of
<i>C. tmetus</i> Tomlin
New Ireland; BMNH (22 mm) |
| 4: ? <i>C. parius</i>
Solomon Is. (40 mm) | 21: <i>C. ochroleucus tmetus</i>
Papua New Guinea (69 mm) |
| 5: <i>C. parius</i>
Vanuatu (40 mm) | 22: <i>C. flavus</i> Röckel; paratype
Solomon Is. (48 mm) |
| 6: <i>C. parius</i>
Vanuatu (37 mm) | 23: <i>C. flavus</i>
Batangas Bay, Philippines (40 mm) |
| 7: <i>C. parius</i>
Philippines (41 mm) | 24: <i>C. flavus</i>
Papua New Guinea (61 mm) |
| 8: <i>C. cinereus cinereus</i>
Marinduque, Philippines (52 mm) | 25: <i>C. flavus</i>
Batangas Bay, Philippines (55 mm) |
| 9: <i>C. cinereus cinereus</i> f. <i>bernardii</i>
Sulu Sea, Philippines (50 mm) | 26: <i>C. flavus</i> var.
Palawan, Philippines; coll. Fischöder (77 mm) |
| 10: <i>C. cinereus cinereus</i>
Karimata Str., Indonesia (47 mm) | 27: <i>C. flavus</i> var.
Palawan, Philippines; coll. Fischöder (63 mm) |
| 11: <i>C. cinereus cinereus</i>
Java (48 mm) | 28: <i>C. oishii</i>
Taiwan (35 mm) |
| 12: <i>C. cinereus cinereus</i> f. <i>bernardii</i>
Sulu Sea, Philippines (46 mm) | 29: <i>C. oishii</i>
Taiwan (33 mm) |
| 13: <i>C. cinereus cinereus</i>
Sorsogon, Philippines (38 mm) | 30: <i>C. oishii</i>
Taiwan (40 mm) |
| 14: <i>C. cinereus gubba</i>
coll. Haksal (57 mm) | 31: <i>C. oishii</i>
N.E. Sumba, Indonesia; RMNH (44 mm) |
| 15: <i>C. cinereus gubba</i>
Manus Id., Papua New Guinea (43 mm) | 32: <i>C. oishii</i>
N.E. Sumba, Indonesia; RMNH (38 mm) |
| 16: <i>C. ochroleucus ochroleucus</i>
Batangas Bay, Philippines (65 mm) | |
| 17: <i>C. ochroleucus ochroleucus</i>
Batangas Bay, Philippines (62 mm) | |



Plate 19

(Text pp. 117-120)

- | | |
|--|--|
| 1: <i>C. consors</i>
Coral Sea (70 mm) | 17: <i>C. furvus</i>
Palawan, Philippines (42 mm) |
| 2: <i>C. consors</i> f. <i>daullei</i>
Mozambique (72 mm) | 18: <i>C. furvus</i>
Balabac, Palawan, Philippines (38 mm) |
| 3: <i>C. consors</i>
Mactan, Philippines (59 mm) | 19: <i>C. furvus</i> f. <i>granifer</i>
Sulu Sea (35 mm) |
| 4: <i>C. consors</i> f. <i>turschi</i>
Kantang, W. Thailand (79 mm) | 20: <i>C. furvus</i> f. <i>granifer</i>
Turtle Id., Philippines (35 mm) |
| 5: <i>C. consors</i>
Bohol, Philippines (79 mm) | 21: <i>C. furvus</i>
Sorsogon, Philippines (57 mm) |
| 6: <i>C. consors</i>
Vanuatu (51 mm) | 22: <i>C. furvus</i>
Culion, Philippines (42 mm) |
| 7: <i>C. consors</i>
Mozambique; coll. Korn (73 mm) | 23: <i>C. furvus</i> f. <i>aegrotus</i>
Sulu Sea (52 mm) |
| 8: <i>C. consors</i>
Philippines; NMC (85 mm) | 24: <i>C. furvus</i> f. <i>polygrammus</i> ; holotype of
<i>C. polygrammus</i> Tomlin
NMWC; (39 mm) |
| 9: <i>C. consors</i>
Solomon Is. (99 mm) | 25: <i>C. furvus</i> f. <i>albicans</i>
Balabac, Palawan, Philippines (35 mm) |
| 10: <i>C. consors</i> f. <i>poehlianus</i>
Solomon Is. (67 mm) | 26: <i>C. furvus</i> f. <i>turritinus</i>
E. coast of Malay. Peninsula (50 mm) |
| 11: <i>C. consors</i> f. <i>poehlianus</i>
Papua New Guinea (55 mm) | 27: <i>C. furvus</i> f. <i>nivalis</i>
Sorsogon, Philippines (59 mm) |
| 12: <i>C. consors</i> ; white form
Guam; coll. da Motta, SMNS (55 mm) | 28: <i>C. furvus</i> f. <i>lignarius</i> ; syntype of <i>C. lignarius</i> Reeve
Leyte, Philippines; BMNH (49 mm) |
| 13: <i>C. furvus</i>
Sulu Sea (59 mm) | 29: <i>C. furvus</i> ; holotype of <i>C. albus</i> Sowerby III
BMNH; photo Filmer (49 mm) |
| 14: <i>C. furvus</i>
Palawan, Philippines (46 mm) | 30: ? <i>C. furvus</i> var.
Cebu, Philippines (35 mm) |
| 15: <i>C. furvus</i> f. <i>neobuxeus</i>
Amami-o-shima, Japan (47 mm) | 31: ? <i>C. furvus</i> var.
as 30; apertural view |
| 16: <i>C. furvus</i>
Sulu Sea (45 mm) | |




Plate 20

(Text pp. 120-123)

- | | |
|--|---|
| 1: <i>C. magus</i>
New Caledonia (53 mm) | 17: <i>C. magus</i> f. <i>metcalfii</i> ; syntype of <i>C. metcalfii</i> Reeve
BMNH (28 mm) |
| 2: <i>C. magus</i>
Mauritius; NMWC (ca. 40 mm) | 18: <i>C. magus</i> ; holotype of <i>C. cernohorskyi</i> da Motta
MHNG (47 mm) |
| 3: <i>C. magus</i>
Solomon Is. (56 mm) | 19: <i>C. magus</i> f. <i>metcalfii</i>
Palawan, Philippines (56 mm) |
| 4: <i>C. magus</i>
Zamboanga, Philippines (66 mm) | 20: <i>C. magus</i> f. <i>epistomium</i> ; lectotype of
<i>C. epistomium</i> Reeve
"Mauritius"; BMNH (48 mm) |
| 5: <i>C. magus</i>
Queensland (67 mm) | 21: <i>C. magus</i> f. <i>raphanus</i>
Cebu, Philippines (65 mm) |
| 6: <i>C. magus</i>
Sorsogon, Philippines (54 mm) | 22: <i>C. magus</i> f. <i>raphanus</i>
Banda Id., Moluccas; coll. Salminger (60 mm) |
| 7: <i>C. magus</i>
Pt. Moresby, Papua New Guinea (54 mm) | 23: <i>C. magus</i>
Siasi, Sulu Sea; coll. da Motta, SMNS (55 mm) |
| 8: <i>C. magus</i>
New Britain, Papua New Guinea; coll. Richards
(49 mm) | 24: <i>C. magus</i>
Sulu Sea; coll. Baer (53 mm) |
| 9: <i>C. magus</i>
Nimoo Id., Papua New Guinea; coll. Danila
(49 mm) | 25: <i>C. magus</i> f. <i>signifer</i> ; holotype of <i>C. signifer</i> Crosse
Fiji; BMNH (40 mm) |
| 10: <i>C. magus</i> f. <i>assimilis</i> ; lectotype of <i>C. assimilis</i> A.
Adams
"Australia"; BMNH (53 mm) | 26: <i>C. magus</i> f. <i>ustulatus</i> ; syntype of <i>C. ustulatus</i> Reeve
"New Holland"; BMNH (42 mm) |
| 11: <i>C. magus</i> ; holotype of <i>C. fulvobullatus</i> da Motta
E. coast of Malaysia; MHNG (53 mm) | 27: <i>C. magus</i> f. <i>ustulatus</i>
Cebu, Philippines; coll. Baer (54 mm) |
| 12: <i>C. magus</i> f. <i>assimilis</i>
Taiwan (76 mm) | 28: <i>C. magus</i> ; holotype of <i>C. rollandi</i> Bernardi
MNHN (33 mm) |
| 13: <i>C. magus</i> f. <i>consul</i> ; original figure of
<i>C. consul</i> Boivin (43 mm) | 29: <i>C. magus</i> ; juvenile; holotype of <i>C. epistomioides</i>
Weinkauff
LMD (33 mm) |
| 14: <i>C. magus</i> f. <i>consul</i>
Sorsogon, Philippines (70 mm) | 30: <i>C. magus</i> ; juvenile; holotype of <i>C. ambaroides</i>
Shikama
"Philippines"; KPM (33 mm) |
| 15: <i>C. magus</i> f. <i>carinatus</i>
Sulu Sea (63 mm) | 31: <i>C. magus</i> ; juvenile; syntype of <i>C. fucatus</i> Reeve
Philippines; BMNH (19 mm) |
| 16: <i>C. magus</i> f. <i>borneensis</i>
N. Borneo; coll. da Motta, SMNS (60 mm) | 32: <i>C. magus</i> Linné; neotype
ZMUU (43 mm) |

