

Plate 21

(Text pp. 123-126)

- | | |
|---|--|
| 1: <i>C. planorbis</i> ; lectotype of <i>C. vulpinus</i> Hwass in Bruguière
MHNG; photo Dajoz (51 mm) | 17: <i>C. planorbis</i> f. <i>vitulinus</i>
Philippines (53 mm) |
| 2: <i>C. ferrugineus</i>
Philippines (50 mm) | 18: <i>C. planorbis</i> f. <i>vitulinus</i>
Sulu Sea (55 mm) |
| 3: <i>C. ferrugineus</i>
Outer Swain Reefs, Queensland (45 mm) | 19: <i>C. planorbis</i> f. <i>vitulinus</i>
Solomon Is. (50 mm) |
| 4: <i>C. ferrugineus</i>
Philippines (50 mm) | 20: ? <i>C. planorbis</i>
Kwajalein (41 mm) |
| 5: <i>C. ferrugineus</i>
Sulu Sea (61 mm) | 21: <i>C. planorbis</i>
Samoa (33 mm) |
| 6: <i>C. ferrugineus</i>
Philippines (57 mm) | 22: <i>C. striatellus</i>
Philippines (59 mm) |
| 7: <i>C. ferrugineus</i>
Solomon Is. (47 mm) | 23: <i>C. striatellus</i>
Solomon Is. (40 mm) |
| 8: <i>C. ferrugineus</i> f. <i>chenui</i> ; lectotype of
<i>C. chenui</i> Crosse
New Caledonia; MNHN (49 mm) | 24: <i>C. striatellus</i>
Kenya (51 mm) |
| 9: <i>C. ferrugineus</i> Hwass in Bruguière; lectotype
MHNG; photo Dajoz (52 mm) | 25: <i>C. striatellus</i>
Aden (60 mm) |
| 10: <i>C. ferrugineus</i> ; holotype of <i>C. sophiae</i> Brazier
AMS; photo Filmer (39 mm) | 26: <i>C. striatellus</i>
Mozambique (51 mm) |
| 11: <i>C. planorbis</i> Born; lectotype
NMW; photo Wawra (41 mm) | 27: <i>C. striatellus</i>
Madras, India (64 mm) |
| 12: <i>C. planorbis</i>
Marinduque, Philippines (46 mm) | 28: <i>C. striatellus</i>
Maldives (43 mm) |
| 13: <i>C. planorbis</i>
Solomon Is. (48 mm) | 29: <i>C. striatellus</i>
Zanzibar (52 mm) |
| 14: <i>C. planorbis</i>
Kwajalein, Marshall Is. (49 mm) | 30: <i>C. striatellus</i> ; original figure of <i>C. lictor</i> Crosse
(40 mm) |
| 15: <i>C. planorbis</i>
Philippines (50 mm) | 31: <i>C. striatellus</i>
Solomon Is. (44 mm) |
| 16: <i>C. planorbis</i> f. <i>vitulinus</i>
Marinduque, Philippines (49 mm) | |



Plate 22

(Text pp. 126-131)

- | | |
|--|--|
| 1: <i>C. circumactus</i>
Siasi, Sulu Sea (43 mm) | 17: <i>C. litoglyphus</i> f. <i>seychellensis</i>
Phuket, W. Thailand (46 mm) |
| 2: <i>C. circumactus</i>
Hawaii (38 mm) | 18: <i>C. litoglyphus</i>
Zanzibar (43 mm) |
| 3: <i>C. circumactus</i>
Siasi, Sulu Sea (49 mm) | 19: <i>C. litoglyphus</i>
Bohol, Philippines (42 mm) |
| 4: <i>C. circumactus</i>
Bohol, Philippines (57 mm) | 20: <i>C. litoglyphus</i>
Zanzibar (50 mm) |
| 5: <i>C. circumactus</i>
Mozambique (42 mm) | 21: <i>C. litoglyphus</i>
W. Thailand; coll. da Motta, SMNS (35 mm) |
| 6: <i>C. circumactus</i>
Mozambique (48 mm) | 22: <i>C. litoglyphus</i> f. <i>carpenteri</i> ; holotype of
<i>C. carpenteri</i> Crosse
BMNH (47 mm) |
| 7: <i>C. circumactus</i> ; juvenile; holotype of <i>C. hammatus</i>
Bartsch & Rehder
Hawaii (23 mm) | 23: <i>C. fumigatus</i>
Massawa, Ethiopia (51 mm) |
| 8: <i>C. swainsoni</i> Estival & von Cosel; holotype
off Noumea, New Caledonia; MNHN (56 mm) | 24: <i>C. fumigatus</i>
Djibouti (43 mm) |
| 9: <i>C. swainsoni</i> Estival & von Cosel; holotype
apertural view | 25: <i>C. fumigatus</i>
Syrte, Libya, Mediterranean Sea (30 mm) |
| 10: <i>C. swainsoni</i> ; juvenile
New Caledonia (26 mm) | 26: <i>C. fumigatus</i> f. <i>blainvillei</i> ; original figure of
<i>C. blainvillei</i> Kiener
(37 mm) |
| 11: <i>C. swainsoni</i>
New Caledonia; coll. Baer (54 mm) | 27: <i>C. fumigatus</i> f. <i>blainvillei</i>
Dahlak Is., Red Sea (43 mm) |
| 12: <i>C. swainsoni</i>
New Caledonia; coll. Baer (38 mm) | 28: <i>C. fumigatus</i> f. <i>blainvillei</i>
Red Sea (46 mm) |
| 13: <i>C. rawaiensis</i>
Raya Id., W. Thailand (42 mm) | 29: <i>C. fumigatus</i> f. <i>excavatus</i> ; holotype of <i>C. excavatus</i>
Sowerby II
BMNH (41 mm) |
| 14: <i>C. rawaiensis</i>
Raya Id., W. Thailand (35 mm) | 30: <i>C. fumigatus</i> f. <i>henoquei</i> ; holotype of <i>C. henoquei</i>
Bernardi
MNHN (38 mm) |
| 15: <i>C. rawaiensis</i>
Raya Id., W. Thailand (33 mm) | |
| 16: <i>C. litoglyphus</i>
Samar, Philippines (48 mm) | |



Plate 23

(Text pp. 131-137)

- | | |
|--|--|
| 1: <i>C. vexillum vexillum</i>
Marquesas (80 mm) | 13: <i>C. capitaneus</i>
N.W. Australia (67 mm) |
| 2: <i>C. vexillum vexillum</i>
Palawan, Philippines (66 mm) | 14: <i>C. capitaneus</i> f. <i>ceciliae</i>
Phuket, W. Thailand (30 mm) |
| 3: <i>C. vexillum sumatrensis</i>
Jeddah, Saudi Arabia (82 mm) | 15: <i>C. capitaneus</i>
Philippines (50 mm) |
| 4: <i>C. vexillum sumatrensis</i>
Berbera, Somalia (67 mm) | 16: <i>C. namocanus</i>
Zanzibar (51 mm) |
| 5: <i>C. vexillum vexillum</i> ; juvenile
Philippines (27 mm) | 17: <i>C. namocanus</i>
Zanzibar (70 mm) |
| 6: <i>C. vexillum vexillum</i> ; juvenile
Cebu, Philippines (31 mm) | 18: <i>C. namocanus</i>
N. Madagascar (50 mm) |
| 7: <i>C. mustelinus</i>
Solomon Is. (55 mm) | 19: <i>C. namocanus</i> f. <i>badius</i>
Red Sea; coll. Danila (62 mm) |
| 8: <i>C. mustelinus</i>
Marinduque, Philippines (70 mm) | 20: <i>C. namocanus</i> var.
Masirah Id., Oman; ZMA (63 mm) |
| 9: <i>C. mustelinus</i>
Singapore (50 mm) | 21: <i>C. miles</i>
Philippines (67 mm) |
| 10: <i>C. mustelinus</i>
Cebu, Philippines (58 mm) | 22: <i>C. miles</i>
Zanzibar (52 mm) |
| 11: <i>C. capitaneus</i>
Tanga, Tanzania (62 mm) | 23: <i>C. miles</i>
Zanzibar (58 mm) |
| 12: <i>C. capitaneus</i>
Queensland (55 mm) | 24: <i>C. miles</i>
Solomon Is. (68 mm) |



Plate 24

(Text pp. 137-141)

- | | |
|--|--|
| 1: <i>C. trigonus</i>
W. Australia (48 mm) | 14: <i>C. lischkeanus</i>
Muscat, Oman (46 mm) |
| 2: <i>C. trigonus</i>
W. Australia (52 mm) | 15: <i>C. lischkeanus</i>
Mozambique (36 mm) |
| 3: <i>C. trigonus</i>
N.W. Australia (46 mm) | 16: <i>C. lischkeanus</i>
Somalia; coll. G. Raybaudi Massilia (28 mm) |
| 4: <i>C. trigonus</i>
N. Australia (39 mm) | 17: <i>C. lischkeanus</i>
Somalia; coll. G. Raybaudi Massilia (38 mm) |
| 5: <i>C. trigonus</i> f. <i>adami</i>
Gulf of Carpentaria, N. Australia (54 mm) | 18: <i>C. rattus</i>
Kenya (41 mm) |
| 6: <i>C. lischkeanus</i>
Japan (48 mm) | 19: <i>C. rattus</i>
Solomon Is. (37 mm) |
| 7: <i>C. lischkeanus</i>
Japan (46 mm) | 20: <i>C. rattus</i>
Mauritius (46 mm) |
| 8: <i>C. lischkeanus</i>
Cape Moreton, Queensland (39 mm) | 21: <i>C. rattus</i> f. <i>semivelatus</i>
Eilat, Red Sea (37 mm) |
| 9: <i>C. lischkeanus</i>
Exmouth Gulf, W. Australia (44 mm) | 22: <i>C. rattus</i> f. <i>taitensis</i>
Marquesas (41 mm) |
| 10: <i>C. lischkeanus</i>
Pt. Hedland, W. Australia (37 mm) | 23: <i>C. rattus</i> f. <i>taitensis</i>
Tahiti (54 mm) |
| 11: <i>C. lischkeanus</i>
N. Zululand, S. Africa (34 mm) | 24: <i>C. capitanellus</i>
Japan (36 mm) |
| 12: <i>C. lischkeanus</i>
Durban, S. Africa; coll. Connell (48 mm) | 25: <i>C. capitanellus</i>
Taiwan (36 mm) |
| 13: <i>C. lischkeanus</i>
Tanzania; coll. Lorenz (45 mm) | 26: <i>C. capitanellus</i>
Taiwan (35 mm) |




Plate 25

(Text pp. 141-145)

- | | |
|--|--|
| 1: <i>C. eximius</i>
Sri Lanka (40 mm) | 15: <i>C. tribblei queenslandis</i> var.
Lady Musgrave Id., Queensland; coll. Ford
(83 mm) |
| 2: <i>C. eximius</i>
Madras, India (40 mm) | 16: <i>C. lenavati</i> da Motta & Röckel; holotype
Cebu, Philippines; MHNG (48 mm) |
| 3: <i>C. eximius</i>
Cuddalore, India (50 mm) | 17: <i>C. lenavati</i>
Cebu, Philippines (58 mm) |
| 4: <i>C. eximius</i>
W. Thailand (40 mm) | 18: <i>C. lenavati</i>
Cebu, Philippines (65 mm) |
| 5: <i>C. eximius</i>
S. Vietnam; coll. Bondarev (28 mm) | 19: <i>C. lenavati</i>
S. Vietnam; coll. Bondarev (38 mm) |
| 6: <i>C. malacanus</i>
Cuddalore, India (57 mm) | 20: ? <i>C. lenavati</i>
Cebu, Philippines (75 mm) |
| 7: <i>C. malacanus</i>
Cuddalore, India (45 mm) | 21: <i>C. sugimotonis</i> (with periostracum)
Tosa Bay, Japan (88 mm) |
| 8: <i>C. malacanus</i>
Kilakarai, India (65 mm) | 22: <i>C. sugimotonis</i>
Queensland (100 mm) |
| 9: <i>C. malacanus</i>
Cuddalore, India (71 mm) | 23: <i>C. sugimotonis</i>
Bohol, Philippines (61 mm) |
| 10: <i>C. tribblei tribblei</i>
Taiwan (77 mm) | 24: <i>C. sugimotonis</i>
Bohol, Philippines (77 mm) |
| 11: <i>C. tribblei tribblei</i>
Taiwan (84 mm) | 25: <i>C. ione</i>
Cebu, Philippines (57 mm) |
| 12: <i>C. tribblei tribblei</i>
Cebu, Philippines (82 mm) | 26: <i>C. ione</i>
Mozambique; coll. Amorim (58 mm) |
| 13: <i>C. tribblei queenslandis</i>
Queensland (59 mm) | 27: <i>C. ione</i>
Tosa Bay, Japan; coll. da Motta, SMNS (66 mm) |
| 14: <i>C. tribblei queenslandis</i>
Queensland (82 mm) | 28: <i>C. ione</i>
Bohol, Philippines (54 mm) |



Plate 26

(Text pp. 145-150)

- | | |
|---|---|
| 1: <i>C. teramachii</i> (with periostracum)
Japan (91 mm) | 16: <i>C. pergrandis</i>
Bohol, Philippines; coll. da Motta, SMNS
(127 mm) |
| 2: <i>C. teramachii</i> (with periostracum)
Majunga, Madagascar; coll. Schmidt (62 mm) | 17: <i>C. pergrandis</i>
Bohol, Philippines (129 mm) |
| 3: <i>C. teramachii</i>
Broome, N.W. Australia (65 mm) | 18: <i>C. pergrandis</i>
Philippines (110 mm) |
| 4: <i>C. teramachii</i>
Somalia (73 mm) | 19: <i>C. pergrandis</i> ; subadult
Philippines; coll. Korn (80 mm) |
| 5: <i>C. teramachii</i>
as 4; apertural view | 20: <i>C. tisii</i>
N.E. Taiwan (123 mm) |
| 6: <i>C. bozzettii</i>
Somalia (57 mm) | 21: <i>C. tisii</i>
Balut Id., Philippines; coll. Schmid (98mm) |
| 7: <i>C. bozzettii</i>
Yemen; coll. Haksal (53 mm) | 22: <i>C. tisii</i> Lan; paratype
N.E. Taiwan; coll. & photo Lan (154 mm) |
| 8: <i>C. bozzettii</i>
Somalia; coll. Haksal (58 mm) | 23: <i>C. tisii</i> ; subadult
Balut Id., Philippines; coll. G. Raybaudi Massi-
lia (70 mm) |
| 9: <i>C. bozzettii</i>
Somalia (46 mm) | 24: <i>C. excelsus</i>
Philippines; coll. Pagobo (76 mm) |
| 10: <i>C. sieboldii</i>
Taiwan (77 mm) | 25: <i>C. excelsus</i>
Solomon Is. (74 mm) |
| 11: <i>C. sieboldii</i>
Taiwan (79 mm) | 26: <i>C. excelsus</i>
Philippines; coll. Pagobo (65 mm) |
| 12: <i>C. sieboldii</i>
Taiwan (76 mm) | 27: <i>C. excelsus</i>
Solomon Is. (65 mm) |
| 13: <i>C. luciae</i> Moolenbeek; holotype
Chesterfield Is.; MNHN (56 mm) | 28: <i>C. excelsus</i>
Balut Id., Philippines (52 mm) |
| 14: <i>C. luciae</i>
Coral Sea; MNHN (56 mm) | |
| 15: <i>C. luciae</i>
New Caledonia (37 mm) | |



Plate 27

(Text pp. 150-155)

- | | |
|---|--|
| 1: <i>C. gratacapii</i> Pilsbry; syntype
Kikai, Japan; ANSP (31 mm) | 15: <i>C. profundorum</i>
Tosa, Japan; MNHN (74 mm) |
| 2: <i>C. gratacapii</i> Pilsbry; syntype
Kikai, Japan; ANSP (30 mm) | 16: <i>C. profundorum</i>
New Caledonia; MNHN (72 mm) |
| 3: <i>C. gratacapii</i>
Taiwan; coll. Lyn (29 mm) | 17: <i>C. profundorum</i> ; holotype of <i>C. soyomaruae</i>
Okutani
Hachijo Id., Japan; GIUT (76 mm) |
| 4: <i>C. smirna</i> Bartsch & Rehder; holotype
(with periostracum)
Kauai, Hawaii; USNM (61 mm) | 18: <i>C. ikedai</i> Ninomiya; paratype
Sagami Bay, Japan; AMNH (30 mm) |
| 5: <i>C. smirna</i> Bartsch & Rehder; holotype
apertural view | 19: <i>C. ikedai</i> Ninomiya; paratype
as 18; apertural view |
| 6: <i>C. smirna</i> var.
Kermadec Ridge; NMNZ; photo Filmer | 20: <i>C. scopulicola</i> Okutani; holotype
Hyotanse, Japan; NSMT (22 mm) |
| 7: <i>C. smirna</i> var.
New Caledonia; MNHN (64 mm) | 21: <i>C. scopulicola</i> Okutani; holotype
apertural view |
| 8: <i>C. smirna</i> var.
Oahu, Hawaii; MNHN (61 mm) | 22: <i>C. darkini</i> Röckel, Korn & Richard; holotype
Loyalty Is.; MNHN (56 mm) |
| 9: <i>C. smirna</i> var.
as 8; apertural view | 23: <i>C. darkini</i>
Loyalty Is.; MNHN (50 mm) |
| 10: <i>C. lani</i> Crandall; holotype
N.E. Taiwan; coll. & photo Lan (53 mm) | 24: <i>C. darkini</i>
Kita-Koho Bank, Philippine Sea; coll. Darkin
(70 mm) |
| 11: <i>C. lani</i>
Bohol, Philippines (50 mm) | 25: <i>C. darkini</i>
Balut Id., Philippines; coll. Schmid (84 mm) |
| 12: <i>C. lani</i>
Loyalty Is.; MNHN (39 mm) | 26: <i>C. jeanmartini</i> G. Raybaudi Massilia; holotype
Réunion; MNHN (47 mm) |
| 13: <i>C. profundorum</i>
Balut Id., Philippines; coll. Schmid (111 mm) | 27: <i>C. jeanmartini</i> G. Raybaudi Massilia; holotype
apertural view |
| 14: <i>C. profundorum</i>
Tosa, Japan (81 mm) | 28: <i>C. jeanmartini</i> G. Raybaudi Massilia; paratype
Réunion; coll. G. Raybaudi Massilia (43 mm) |



Plate 28

(Text pp. 155-157)

- | | |
|---|---|
| 1: <i>C. stupa</i>
Solomon Is. (89 mm) | 14: <i>C. recluzianus</i> f. <i>gloriakiensis</i>
Japan (61 mm) |
| 2: <i>C. stupa</i>
Taiwan (83 mm) | 15: <i>C. recluzianus</i>
Taiwan (66 mm) |
| 3: <i>C. stupa</i>
as 2; apertural view | 16: <i>C. recluzianus</i>
Madras, India (58 mm) |
| 4: <i>C. stupella</i>
Taiwan (68 mm) | 17: <i>C. recluzianus</i>
W. Thailand (63 mm) |
| 5: <i>C. stupella</i>
Japan (55 mm) | 18: <i>C. recluzianus</i>
Cebu, Philippines (56 mm) |
| 6: <i>C. stupella</i>
Taiwan (76 mm) | 19: <i>C. recluzianus</i>
Cape Moreton, Queensland; coll. Ford (70 mm) |
| 7: <i>C. hirasei</i>
Taiwan (54 mm) | 20: <i>C. recluzianus</i>
Bathurst Id., N.W. Australia; coll. Limpus (39 mm) |
| 8: <i>C. hirasei</i>
Philippines; coll. Korn (63 mm) | 21: <i>C. recluzianus</i> f. <i>roseorapum</i>
Bohol, Philippines (57 mm) |
| 9: <i>C. hirasei</i>
Bohol, Philippines; coll. Pagobo (63 mm) | 22: <i>C. recluzianus</i> f. <i>roseorapum</i>
Cebu, Philippines (63 mm) |
| 10: <i>C. recluzianus</i> Bernardi; holotype
"mers de Chine"; LMD (58 mm) | 23: <i>C. recluzianus</i> f. <i>roseorapum</i>
Bohol, Philippines (49 mm) |
| 11: <i>C. recluzianus</i>
Japan (59 mm) | 24: <i>C. recluzianus</i> f. <i>roseorapum</i> ?
South China Sea (31 mm) |
| 12: <i>C. recluzianus</i>
Japan (52 mm) | 25: <i>C. recluzianus</i> f. <i>roseorapum</i> (fragment)
Solomon Is. |
| 13: <i>C. recluzianus</i> f. <i>urashimanus</i>
Japan (66 mm) | |



Plate 29

(Text pp. 157-164)

- | | |
|--|---|
| 1: <i>C. sukhadwalai</i> Röckel & da Motta; paratype
Madras, India (45 mm) | 18: <i>C. lentiginosus</i>
Bombay, India (35 mm) |
| 2: <i>C. sukhadwalai</i> Röckel & da Motta; paratype
Madras, India (43 mm) | 19: <i>C. dictator</i>
Rameswaram, India (42 mm) |
| 3: <i>C. sukhadwalai</i> Röckel & da Motta; paratype
Madras, India (50 mm) | 20: <i>C. dictator</i>
Rameswaram, India (36 mm) |
| 4: <i>C. shikamai</i>
Bohol, Philippines (49 mm) | 21: <i>C. dictator</i>
Karachi, Pakistan (42 mm) |
| 5: <i>C. shikamai</i>
Bohol, Philippines (57 mm) | 22: <i>C. dictator</i>
Sri Lanka; coll. Huber (27 mm) |
| 6: <i>C. shikamai</i>
Cebu, Philippines (62 mm) | 23: <i>C. milesi</i>
Oman; BMNH (24 mm) |
| 7: <i>C. shikamai</i>
South China Sea; coll. Bondarev (25 mm) | 24: <i>C. milesi</i>
Oman; BMNH (26 mm) |
| 8: <i>C. bayani</i>
Madras, India (48 mm) | 25: <i>C. milesi</i> E. A. Smith; holotype
Oman; BMNH (21 mm) |
| 9: <i>C. bayani</i>
Madras, India (53 mm) | 26: <i>C. traversianus</i>
Aden (39 mm) |
| 10: <i>C. bayani</i>
Cuddalore, India (53 mm) | 27: <i>C. traversianus</i>
Aden (36 mm) |
| 11: <i>C. bayani</i>
Madras, India (54 mm) | 28: <i>C. traversianus</i>
Somalia; coll. G. Raybaudi Massilia (27 mm) |
| 12: <i>C. bayani</i>
Red Sea; coll. Bondarev (54 mm) | 29: <i>C. stocki</i>
Muscat, Oman (36 mm) |
| 13: <i>C. capreolus</i> Röckel; paratype
Visakhapatnam, India (44 mm) | 30: <i>C. stocki</i>
Muscat, Oman (31 mm) |
| 14: <i>C. capreolus</i> Röckel; paratype
Visakhapatnam, India (42 mm) | 31: <i>C. stocki</i>
Muscat, Oman; coll. da Motta, SMNS (41 mm) |
| 15: <i>C. capreolus</i>
Kantang, W. Thailand (65 mm) | 32: <i>C. lizarum</i>
Somalia; coll. G. Raybaudi Massilia (29 mm) |
| 16: <i>C. lentiginosus</i>
Bombay, India (40 mm) | 33: <i>C. lizarum</i>
as 32; apertural view |
| 17: <i>C. lentiginosus</i>
Bombay, India (33 mm) | 34: <i>C. lizarum</i>
Somalia; coll. G. Raybaudi Massilia (24 mm) |



Plate 30

(Text pp. 164-166)

- | | |
|---|--|
| 1: <i>C. voluminalis</i> Reeve; original figure
(38 mm) | 17: <i>C. voluminalis</i>
Palawan, Philippines; coll. Korn (52 mm) |
| 2: <i>C. voluminalis</i>
Kantang, W. Thailand (43 mm) | 18: <i>C. voluminalis</i>
Sorigao, Mindanao, Philippines; coll. Pagobo
(54 mm) |
| 3: <i>C. voluminalis</i> ; holotype of <i>C. macarae</i> Bernardi
MNHN (39 mm) | 19: <i>C. papuensis</i>
Hansa Bay, Papua New Guinea (27 mm) |
| 4: <i>C. voluminalis</i> ; holotype of <i>C. filicinctus</i> Schepman
ZMA (27 mm) | 20: <i>C. papuensis</i>
Hansa Bay, Papua New Guinea; SMNS (24 mm) |
| 5: <i>C. voluminalis</i>
Gulf of Thailand (51 mm) | 21: <i>C. papuensis</i>
Hansa Bay, Papua New Guinea; SMNH (26 m) |
| 6: <i>C. voluminalis</i>
W. Thailand (49 mm) | 22: ? <i>C. papuensis</i>
Solomon Is. (25 mm) |
| 7: <i>C. voluminalis</i>
Sulu Sea, Philippines; coll. Schmid (43 mm) | 23: <i>C. locumtenens</i>
Aden (46 mm) |
| 8: <i>C. voluminalis</i>
Solomon Is. (42 mm) | 24: <i>C. locumtenens</i>
Massawa, Red Sea (59 mm) |
| 9: <i>C. voluminalis</i>
W. Thailand (73 mm) | 25: <i>C. locumtenens</i>
Massawa, Red Sea (49 mm) |
| 10: <i>C. voluminalis</i>
Philippines; coll. Schmid (46 mm) | 26: <i>C. locumtenens</i>
Red Sea (50 mm) |
| 11: <i>C. voluminalis</i>
Maldives; coll. Frantzen (32 mm) | 27: <i>C. locumtenens</i> ; juvenile
Saudi Arabia; coll. Sharabati, BMNH (23 mm) |
| 12: <i>C. voluminalis</i>
Bangka Str., Indonesia (44 mm) | 28: <i>C. locumtenens</i> f. <i>multicatenatus</i>
MNHN (32 mm) |
| 13: <i>C. voluminalis</i>
Pt. Hedland, W. Australia (25 mm) | 29: <i>C. locumtenens</i> f. <i>multicatenatus</i> ; holotype
of <i>C. multicatenatus</i> Sowerby II
BMNH (32 mm) |
| 14: <i>C. voluminalis</i>
Turtle Id., Sulu Sea (41 mm) | 30: <i>C. locumtenens</i> f. <i>schech</i> ; syntype of
<i>C. schech</i> Weinkauff
Massawa; ZMB (31 mm) |
| 15: <i>C. voluminalis</i>
Moluccas; ZMA (39 mm) | |
| 16: <i>C. voluminalis</i>
Philippines; coll. Schmid (43 mm) | |



Plate 31

(Text pp. 166-172)

- | | |
|---|--|
| 1: <i>C. virgo</i>
Moluccas (67 mm) | 14: <i>C. frigidus</i>
Queensland (42 mm) |
| 2: <i>C. virgo</i>
Zanzibar (81 mm) | 15: <i>C. frigidus</i>
Solomon Is. (49 mm) |
| 3: <i>C. virgo</i>
W. Thailand (75 mm) | 16: <i>C. terebra terebra</i>
Marquesas (79 mm) |
| 4: <i>C. emaciatus</i>
Vanuatu (48 mm) | 17: <i>C. terebra terebra</i>
New Caledonia (86 mm) |
| 5: <i>C. emaciatus</i>
E. Malaysia (46 mm) | 18: <i>C. terebra terebra</i>
Pamsam, India (66 mm) |
| 6: <i>C. emaciatus</i>
Philippines (44 mm) | 19: <i>C. terebra terebra</i>
Solomon Is. (85 mm) |
| 7: <i>C. flavidus</i>
Oman (46 mm) | 20: <i>C. terebra terebra</i>
W. Thailand; coll. Roussy (46 mm) |
| 8: <i>C. flavidus</i>
Philippines (50 mm) | 21: <i>C. terebra thomasi</i>
Aden (63 mm) |
| 9: <i>C. flavidus</i>
Tanzania (48 mm) | 22: <i>C. terebra thomasi</i>
Jeddah, Saudi Arabia (72 mm) |
| 10: <i>C. flavidus</i>
Taiwan (52 mm) | 23: <i>C. bondarevi</i> Röckel & G. Raybaudi Massilia;
holotype
Somalia; SMNS (41 mm) |
| 11: <i>C. frigidus</i>
Solomon Is. (40 mm) | 24: <i>C. bondarevi</i> Röckel & G. Raybaudi Massilia;
paratype
Somalia (35 mm) |
| 12: <i>C. frigidus</i>
W. Thailand (34 mm) | 25: <i>C. bondarevi</i> Röckel & G. Raybaudi Massilia;
paratype
Somalia; coll. G. Raybaudi Massilia (41 mm) |
| 13: <i>C. frigidus</i>
Fiji (38 mm) | |




Plate 32

(Text pp. 173-175)

- | | |
|--|---|
| 1: <i>C. martensi</i> E. A. Smith; holotype
Providence Reef, Mascarenes; BMNH (24 mm) | 14: <i>C. coelinae coelinae</i>
New Caledonia (78 mm) |
| 2: <i>C. martensi</i> ; holotype of <i>C. alconnelli</i> da Motta
Richard Bay, Natal; MHNG (62 mm) | 15: <i>C. coelinae coelinae</i>
Coral Sea (82 mm) |
| 3: <i>C. martensi</i>
Majunga, Madagascar (78 mm) | 16: <i>C. coelinae coelinae</i>
Queensland (77 mm) |
| 4: <i>C. martensi</i>
Ras Fartak, Gulf of Aden; coll. Danila (43 mm) | 17: <i>C. coelinae spiceri</i>
Hawaii (110 mm) |
| 5: <i>C. martensi</i>
Park Rynie, Natal; NM (50 mm) | 18: <i>C. coelinae spiceri</i>
as 17; apertural view |
| 6: <i>C. martensi</i>
Gulf of Aden; coll. Bondarev (36 mm) | 19: <i>C. coelinae spiceri</i>
Hawaii; coll. da Motta, SMNS (104 mm) |
| 7: <i>C. berdulinus</i> Veillard; holotype
Réunion; MNHN (68 mm) | 20: <i>C. coelinae</i> Crosse; holotype
New Caledonia; IRSN; photo Delsaerd
(115 mm) |
| 8: <i>C. berdulinus</i>
Madras, India (59 mm) | 21: <i>C. kintoki</i>
Cebu, Philippines (56 mm) |
| 9: <i>C. berdulinus</i>
Nicobar Is.; coll. Peled (73 mm) | 22: <i>C. kintoki</i>
Bohol, Philippines (59 mm) |
| 10: <i>C. berdulinus</i>
Park Rynie, Natal; NM (45 mm) | 23: <i>C. kintoki</i>
Bohol, Philippines (71 mm) |
| 11: <i>C. berdulinus</i>
Kuria Muria Is., Oman; coll. Bondarev (67 mm) | 24: <i>C. kintoki</i>
Bohol, Philippines (69 mm) |
| 12: <i>C. berdulinus</i>
Zululand; NM (57 mm) | |
| 13: <i>C. berdulinus</i>
Somalia; coll. G. Raybaudi Massilia (77 mm) | |



Plate 33

(Text pp. 175-179)

- | | |
|---|--|
| 1: <i>C. generalis</i>
Palawan, Philippines (70 mm) | 16: <i>C. maldivus</i>
Mozambique (60 mm) |
| 2: <i>C. generalis</i>
Samar, Philippines (91 mm) | 17: <i>C. maldivus</i>
Kenya (74 mm) |
| 3: <i>C. generalis</i>
Solomon Is. (59 mm) | 18: <i>C. maldivus</i>
Zanzibar (73 mm) |
| 4: <i>C. generalis</i>
Java, Indonesia (71 mm) | 19: <i>C. maldivus</i>
Mozambique (55 mm) |
| 5: <i>C. generalis</i>
Solomon Is. (60 mm) | 20: <i>C. maldivus</i> ; juvenile; holotype of <i>C. spirogloxus</i>
Deshayes
Réunion; MNHN (22 mm) |
| 6: <i>C. generalis</i>
Balut Id., Philippines (60 mm) | 21: <i>C. monile</i>
Madras, India (91 mm) |
| 7: <i>C. generalis</i>
Philippines (57 mm) | 22: <i>C. monile</i>
Cuddalore, India (79 mm) |
| 8: <i>C. generalis</i>
Taiwan (73 mm) | 23: <i>C. monile</i>
Cuddalore, India (61 mm) |
| 9: <i>C. generalis</i> f. <i>krabiensis</i>
W. Thailand | 24: <i>C. monile</i>
W. Thailand (66 mm) |
| 10: <i>C. generalis</i> f. <i>krabiensis</i>
W. Thailand | 25: <i>C. rufimaculosus</i>
Queensland (50 mm) |
| 11: <i>C. generalis</i> ; juvenile
Solomon Is. (24 mm) | 26: <i>C. rufimaculosus</i>
Cape Moreton, Queensland (45 mm) |
| 12: <i>C. generalis</i> ; juvenile; syntype of <i>C. spiculum</i>
Reeve
BMNH (23 mm) | 27: <i>C. sculletti</i>
Cape Moreton, Queensland (42 mm) |
| 13: <i>C. maldivus</i>
Zanzibar (24 mm) | 28: <i>C. sculletti</i>
Cape Moreton, Queensland (45 mm) |
| 14: <i>C. maldivus</i>
Eilat, Red Sea (62 mm) | 29: <i>C. sculletti</i>
Cape Moreton, Queensland (43 mm) |
| 15: <i>C. maldivus</i>
Zanzibar (62 mm) | |



Plate 34

(Text pp. 179-183)

- | | |
|---|--|
| 1: <i>C. ammiralis ammiralis</i>
E. Thailand (73 mm) | 14: <i>C. thomae</i>
Davao, Philippines; SMNS (73 mm) |
| 2: <i>C. ammiralis ammiralis</i>
Philippines (73 mm) | 15: <i>C. thomae</i>
Philippines; ZMA (83 mm) |
| 3: <i>C. ammiralis ammiralis</i> f. <i>temnes</i>
Queensland (85 mm) | 16: <i>C. splendidulus</i>
Aden (57 mm) |
| 4: <i>C. ammiralis ammiralis</i>
Bohol, Philippines (74 mm) | 17: <i>C. splendidulus</i>
Aden (57 mm) |
| 5: <i>C. ammiralis ammiralis</i>
Moluccas (61 mm) | 18: <i>C. splendidulus</i>
Aden (62 mm) |
| 6: <i>C. ammiralis ammiralis</i> f. <i>architalassus</i>
Solomon Is. (46 mm) | 19: <i>C. splendidulus</i>
Berbera, Somalia (65 mm) |
| 7: <i>C. ammiralis ammiralis</i>
Solomon Is. (56 mm) | 20: <i>C. splendidulus</i>
Berbera, Somalia (50 mm) |
| 8: <i>C. ammiralis pseudocedonulli</i>
Seychelles (56 mm) | 21: <i>C. fumigatus</i> ; syntype of <i>C. luctificus</i> Reeve
BMNH (38 mm) |
| 9: <i>C. ammiralis pseudocedonulli</i>
Kenya (52 mm) | 22: <i>C. ciderryi</i> da Motta; holotype
Strait of Taiwan; MHNG (31 mm) |
| 10: <i>C. ammiralis pseudocedonulli</i>
Mozambique; coll. Amorim (53 mm) | 23: <i>C. ciderryi</i>
Strait of Taiwan; coll. da Motta, SMNS (42 mm) |
| 11: <i>C. ammiralis pseudocedonulli</i>
W. Thailand (42 mm) | 24: <i>C. ciderryi</i>
Vietnam; coll. D. Dan (40 mm) |
| 12: <i>C. thomae</i>
S. Mindanao, Philippines (83 mm) | 25: <i>C. ciderryi</i>
Taiwan (33 mm) |
| 13: <i>C. thomae</i>
Davao, Philippines (77 mm) | |



Plate 35

(Text pp. 183-186)

- | | |
|---|--|
| 1: <i>C. amadis</i>
Rameswaram, India (81 mm) | 14: <i>C. thalassiarachus</i> ; holotype of <i>C. mariei</i> Jousseaume
MNHN (47 mm) |
| 2: <i>C. amadis</i>
Madras, India (80 mm) | 15: <i>C. thalassiarachus</i>
Palawan, Philippines (67 mm) |
| 3: <i>C. amadis</i>
Cuddalore, India (78 mm) | 16: <i>C. thalassiarachus</i>
Sulu Sea, Philippines (72 mm) |
| 4: <i>C. amadis</i>
Cuddalore, India (78 mm) | 17: <i>C. thalassiarachus</i>
Cebu, Philippines (80 mm) |
| 5: <i>C. amadis</i>
Cuddalore, India (76 mm) | 18: <i>C. thalassiarachus</i>
Coron, Palawan, Philippines (73 mm) |
| 6: <i>C. amadis</i> f. <i>castaneofasciata</i>
Phuket, W. Thailand (71 mm) | 19: <i>C. thalassiarachus</i>
Coron, Palawan, Philippines (63 mm) |
| 7: <i>C. amadis</i> f. <i>castaneofasciata</i>
Phuket, W. Thailand (64 mm) | 20: <i>C. thalassiarachus</i>
Manobel Id., Philippines (58 mm) |
| 8: <i>C. amadis</i> f. <i>arbornatalis</i>
Ranong, W. Thailand (50 mm) | 21: <i>C. thalassiarachus</i>
S. Mindanao, Philippines (77 mm) |
| 9: <i>C. amadis</i> f. <i>castaneofasciata</i>
W. Thailand; coll. Roussy (63 mm) | 22: <i>C. iodostoma</i>
Conducia Bay, Mozambique (38 mm) |
| 10: <i>C. amadis</i> f. <i>castaneofasciata</i>
Ranong, W. Thailand (41 mm) | 23: <i>C. iodostoma</i>
Nacala, Mozambique (42 mm) |
| 11: <i>C. amadis</i> f. <i>castaneofasciata</i>
W. Thailand; coll. Peled (39 mm) | 24: <i>C. iodostoma</i>
N. Mozambique (38 mm) |
| 12: <i>C. amadis</i> ; holotype of <i>C. lozeti</i> Richard
"S.E. Madagascar"; MNHN (56 mm) | 25: <i>C. iodostoma</i> ; holotype of <i>C. hedgesi</i> Sowerby II
BMNH (33 mm) |
| 13: <i>C. amadis</i> ; holotype of <i>C. lozeti</i> Richard
apertural view | |



Plate 36

(Text pp. 186-190)

- | | |
|---|--|
| 1: <i>C. nobilis nobilis</i> ; lectotype of <i>C. nobilis</i> Linné
LSL; photo Kohn (41 mm) | 15: <i>C. cordigera</i>
Philippines (43 mm) |
| 2: <i>C. nobilis nobilis</i>
Java, Indonesia; coll. da Motta, SMNS (58 mm) | 16: <i>C. cordigera</i>
Borneo; ZMA (39 mm) |
| 3: <i>C. nobilis nobilis</i>
N.W. Sumatra; coll. da Motta, SMNS (56 mm) | 17: <i>C. cordigera</i> f. <i>bitleri</i>
Tawi Tawi Id., Philippines (35 mm) |
| 4: <i>C. nobilis skinneri</i>
Bali (57 mm) | 18: <i>C. marchionatus</i>
Marquesas (50 mm) |
| 5: <i>C. nobilis skinneri</i>
Bali (56 mm) | 19: <i>C. marchionatus</i>
Marquesas (33 mm) |
| 6: <i>C. nobilis skinneri</i>
"Moluccas" (48 mm) | 20: <i>C. marchionatus</i> (with periostracum)
Marquesas (29 mm) |
| 7: <i>C. nobilis victor</i>
Flores (41 mm) | 21: <i>C. jickelii</i> Weinkauff; lectotype
Eritrea; ZMB (51 mm) |
| 8: <i>C. nobilis victor</i> ; holotype of <i>C. victor</i> Broderip
BMNH (37 mm) | 22: <i>C. jickelii</i>
Djibouti; coll. Nicolay (45 mm) |
| 9: <i>C. nobilis renatae</i>
Andaman Is., IRSN (37 mm) | 23: <i>C. jickelii</i>
Djibouti (43 mm) |
| 10: <i>C. nobilis friedae</i>
Sri Lanka; coll. Huber (46 mm) | 24: <i>C. jickelii</i>
Massawa; coll. da Motta, SMNS (38 mm) |
| 11: <i>C. nobilis friedae</i> ; holotype of <i>Eugeniconus</i>
<i>friedae</i> da Motta
Sri Lanka; MHNG (51 mm) | 25: <i>C. jickelii</i>
Djibouti; coll. Ramalho (40 mm) |
| 12: <i>C. cordigera</i> Sowerby II; lectotype
Cuyo, Philippines; BMNH, photo Filmer
(62 mm) | 26: <i>C. jickelii</i>
Djibouti (44 mm) |
| 13: <i>C. cordigera</i>
Palawan, Philippines (48 mm) | 27: <i>C. jickelii</i> ; lectotype of <i>C. quadratus</i> Röding
Red Sea; ZMUC (34 mm) |
| 14: <i>C. cordigera</i>
Philippines (44 mm) | 28: <i>C. jickelii</i> ; lectotype of <i>C. quadratus</i> Röding
apertural view |



Plate 37

(Text pp. 190-193)

- | | |
|---|---|
| 1: <i>C. angioiorum</i> Röckel & Moolenbeek; holotype Djibouti; SMNS (36 mm) | 17: <i>C. erythraeensis dillwynii</i> ; paralectotype of <i>C. dillwynii</i> Reeve BMNH (31 mm) |
| 2: <i>C. angioiorum</i> Djibouti (37 mm) | 18: ? <i>C. erythraeensis dillwynii</i> S. of Jeddah; coll. da Motta, SMNS (29 mm) |
| 3: <i>C. angioiorum</i> Röckel & Moolenbeek; paratype Djibouti (34 mm) | 19: ? <i>C. erythraeensis dillwynii</i> Jeddah; coll. Filmer (24 mm) |
| 4: <i>C. angioiorum</i> Röckel & Moolenbeek; paratype Djibouti; coll. Mailly (34 mm) | 20: <i>C. nigromaculatus</i> Röckel & Moolenbeek; holotype Ras Andade, Eritrea; SMNS (32 mm) |
| 5: <i>C. erythraeensis erythraeensis</i> ; lectotype of <i>C. erythraeensis</i> Reeve ZMUC (24 mm) | 21: <i>C. nigromaculatus</i> Röckel & Moolenbeek; paratype Ras Andade; coll. Peled (35 mm) |
| 6: <i>C. erythraeensis erythraeensis</i> ; holotype of <i>C. quadratomaculatus</i> Sowerby II BMNH (21 mm) | 22: <i>C. nigromaculatus</i> Röckel & Moolenbeek; paratype Dahlak Archipelago; coll. Nicolay (35 mm) |
| 7: <i>C. erythraeensis erythraeensis</i> ; holotype of <i>C. adustus</i> Sowerby II BMNH (23 mm) | 23: <i>C. nigromaculatus</i> Röckel & Moolenbeek; paratype Eritrea; coll. Ramalho (31 mm) |
| 8: <i>C. erythraeensis erythraeensis</i> Eritrea (28 mm) | 24: <i>C. blanfordianus</i> Philippines; ZMA (40 mm) |
| 9: <i>C. erythraeensis erythraeensis</i> Eritrea (26 mm) | 25: <i>C. blanfordianus</i> Philippines; coll. Korn (50 mm) |
| 10: <i>C. erythraeensis erythraeensis</i> Aden (27 mm) | 26: <i>C. blanfordianus</i> Philippines (31 mm) |
| 11: <i>C. erythraeensis erythraeensis</i> ; Massawa form Massawa, Eritrea (17 mm) | 27: <i>C. blanfordianus</i> Philippines (31 mm) |
| 12: <i>C. erythraeensis erythraeensis</i> ; Massawa form Massawa, Eritrea (19 mm) | 28: <i>C. blanfordianus</i> Cebu, Philippines (29 mm) |
| 13: <i>C. erythraeensis induratus</i> ; original figure of <i>C. induratus</i> Reeve (19 mm) | 29: <i>C. blanfordianus</i> Crosse; original figure (36 mm) |
| 14: <i>C. erythraeensis induratus</i> Pt. Sudan; coll. Vink (23 mm) | 30: <i>C. wittigi</i> Lesser Sunda Is. (37 mm) |
| 15: ? <i>C. erythraeensis</i> Bank Dankali, Djibouti; coll. Reichert (27 mm) | 31: <i>C. wittigi</i> as 30; apertural view |
| 16: <i>C. erythraeensis dillwynii</i> ; lectotype of <i>C. dillwynii</i> Reeve BMNH (28 mm) | 32: <i>C. wittigi</i> f. <i>kongaensis</i> Konga Id., S.E. Flores (43 mm) |
| | 33: <i>C. wittigi</i> f. <i>kongaensis</i> as 32; apertural view |



Plate 38

(Text pp. 194-196)

- | | |
|--|---|
| 1: <i>C. inscriptus inscriptus</i> ; lectotype of <i>C. inscriptus</i> Reeve
BMNH (34 mm) | 16: <i>C. inscriptus inscriptus</i>
Kilakarai, India (49 mm) |
| 2: <i>C. inscriptus inscriptus</i>
Ras Andade, Eritrea (44 mm) | 17: <i>C. inscriptus inscriptus</i>
Madras, India (32 mm) |
| 3: <i>C. inscriptus inscriptus</i>
Aden (39 mm) | 18: <i>C. inscriptus inscriptus</i> ; syntype of <i>C. cuneiformis</i> E. A. Smith
BMNH (29 mm) |
| 4: <i>C. inscriptus inscriptus</i> f. <i>bangladeshianus</i>
Tuticorin, India (51 mm) | 19: <i>C. inscriptus inscriptus</i> var.
Diego Suarez; coll. Monteiro (33 mm) |
| 5: <i>C. inscriptus inscriptus</i>
Madras, India | 20: <i>C. inscriptus inscriptus</i> var.
Réunion; MNHN (36 mm) |
| 6: <i>C. inscriptus inscriptus</i> ; holotype of <i>C. keatiformis</i> Shikama
KPM; photo Filmer (62 mm) | 21: <i>C. inscriptus inscriptus</i> var.
Mauritius (44 mm) |
| 7: <i>C. inscriptus inscriptus</i>
S. India (62 mm) | 22: <i>C. lynceus</i>
Bohol, Philippines (73 mm) |
| 8: <i>C. inscriptus inscriptus</i>
S. India (41 mm) | 23: <i>C. lynceus</i>
Manila Bay, Philippines (62 mm) |
| 9: <i>C. inscriptus inscriptus</i> f. <i>maculospira</i> ; holotype of <i>C. planiliratus</i> Sowerby III
BMNH (41 mm) | 24: <i>C. lynceus</i>
Queensland (65 mm) |
| 10: <i>C. inscriptus inscriptus</i> f. <i>maculospira</i>
W. Thailand (50 mm) | 25: <i>C. lynceus</i>
Philippines (67 mm) |
| 11: <i>C. inscriptus adenensis</i>
Mozambique (58 mm) | 26: <i>C. pretiosus</i>
Tuticorin; India (74 mm) |
| 12: <i>C. inscriptus adenensis</i>
Somalia (62 mm) | 27: <i>C. pretiosus</i>
as 26; apertural view |
| 13: <i>C. inscriptus adenensis</i>
Mozambique (55 mm) | 28: <i>C. pretiosus</i>
Bay of Bengal; coll. Peled (84 mm) |
| 14: <i>C. inscriptus adenensis</i> var.; syntype of <i>C. keatii</i> Sowerby II
Seychelles; BMNH (48 mm) | 29: <i>C. pretiosus</i>
Kantang, W. Thailand; coll. Schmid (76 mm) |
| 15: <i>C. inscriptus adenensis</i> var.; syntype of <i>C. keatii</i> Sowerby II
Seychelles; BMNH (46 mm) | |



Plate 39

(Text pp. 196-199)

- | | |
|--|---|
| 1: <i>C. collisus</i> Reeve; paralectotype
BMNH (43 mm) | 17: <i>C. andamanensis</i>
Andaman Sea; MNHN (29 mm) |
| 2: <i>C. collisus</i>
W. Thailand (49 mm) | 18: <i>C. andamanensis</i>
Andaman Sea; MNHN (28 mm) |
| 3: <i>C. collisus</i>
W. Thailand (46 mm) | 19: <i>C. andamanensis</i>
Andaman Sea; MNHN (29 mm) |
| 4: <i>C. collisus</i>
S. India (44 mm) | 20: <i>C. andamanensis</i>
Andaman Is.; KBIN (21 mm) |
| 5: <i>C. collisus</i>
W. Thailand (46 mm) | 21: <i>C. subulatus</i> Kiener; original figure
(35 mm) |
| 6: <i>C. collisus</i>
Malaysia (49 mm) | 22: <i>C. subulatus</i>
Iloilo, Philippines (33 mm) |
| 7: <i>C. collisus</i>
W. Thailand (41 mm) | 23: <i>C. subulatus</i>
Solomon Is.; coll. da Motta, SMNS (34 mm) |
| 8: <i>C. collisus</i>
Timor (34 mm) | 24: <i>C. subulatus</i>
Solomon Is.; coll. da Motta, SMNS (34 mm) |
| 9: <i>C. collisus</i> ; syntype of <i>C. stigmaticus</i> A. Adams
BMNH (45 mm) | 25: <i>C. subulatus</i>
Solomon Is.; coll. da Motta, SMNS (33 mm) |
| 10: <i>C. collisus</i> f. <i>kieneri</i> ; original figure of
<i>C. kieneri</i> Reeve
(29 mm) | 26: <i>C. broderipii</i> Reeve; holotype
Moluccas; BMNH (28 mm) |
| 11: <i>C. collisus</i> f. <i>kieneri</i>
BMNH (32 mm) | 27: <i>C. broderipii</i>
"Andaman", MNHN (30 mm) |
| 12: <i>C. collisus</i> f. <i>kieneri</i>
Madagascar; MNHN (32 mm) | 28: <i>C. broderipii</i>
Moluccas, ZMA (32 mm) |
| 13: <i>C. collisus</i> f. <i>straturatus</i> ; syntype of <i>C. straturatus</i>
Sowerby II
BMNH (32 mm) | 29: ? <i>C. broderipii</i>
Solomon Is. (39 mm) |
| 14: <i>C. collisus</i> f. <i>straturatus</i>
MNHN (24 mm) | 30: ? <i>C. broderipii</i>
Philippines (38 mm) |
| 15: <i>C. andamanensis</i> E. A. Smith; holotype
Andaman Is.; BMNH (22 mm) | 31: ? <i>C. broderipii</i>
Philippines; coll. Korn (41 mm) |
| 16: <i>C. andamanensis</i> E. A. Smith; original figure
Andaman Is. (31 mm) | 32: ? <i>C. broderipii</i>
Philippines (39 mm) |



Plate 40

(Text pp. 199-204)

- | | |
|--|--|
| 1: <i>C. scalptus</i> Reeve; syntype
(BMNH 27 mm) | 18: <i>C. stramineus mulderi</i>
Philippines (47 mm) |
| 2: <i>C. scalptus</i>
W. Zambelis, Philippines; coll. da Motta, SMNS
(28 mm) | 19: <i>C. stramineus mulderi</i>
Philippines (47 mm) |
| 3: <i>C. scalptus</i>
MNHN (28 mm) | 20: <i>C. stramineus mulderi</i>
Papua New Guinea (35 mm) |
| 4: <i>C. sertacinctus</i> Röckel; paratype
Solomon Is. (32 mm) | 21: <i>C. stramineus mulderi</i> ; holotype of <i>C. mulderi</i> Fulton
Mindoro, Philippines; BMNH (29 mm) |
| 5: <i>C. sertacinctus</i> Röckel; paratype
Solomon Is. (29 mm) | 22: <i>C. zapatosensis</i> Röckel; paratype
Zapatos Is., Philippines (44 mm) |
| 6: <i>C. sertacinctus</i> f. <i>solomonensis</i>
Solomon Is. (30 mm) | 23: <i>C. zapatosensis</i> Röckel; paratype
Zapatos Is., Philippines (36 mm) |
| 7: <i>C. sertacinctus</i> f. <i>solomonensis</i>
Philippines (40 mm) | 24: <i>C. zapatosensis</i> Röckel; paratype
Burias Strait, Philippines (38 mm) |
| 8: ? <i>C. sertacinctus</i> dark form
Solomon Is. (37 mm) | 25: <i>C. zapatosensis</i> Röckel; paratype
Zapatos Is., Philippines; ZSM (49 mm) |
| 9: ? <i>C. sertacinctus</i> dark form
Solomon Is. (32 mm) | 26: <i>C. zebra</i> Lamarck; holotype
MNHN (27 mm) |
| 10: ? <i>C. sertacinctus</i>
Flores; coll. G. Raybaudi Massilia (32 mm) | 27: <i>C. zebra</i>
Solomon Is.; coll. da Motta ex coll. Bitler,
SMNS (29 mm) |
| 11: ? <i>C. sertacinctus</i>
Flores; coll. G. Raybaudi Massilia (32 mm) | 28: <i>C. zebra</i>
New Britain, Papua New Guinea (30 mm) |
| 12: ? <i>C. sertacinctus</i>
Batangas Bay, Philippines (32 mm) | 29: <i>C. zebra</i>
New Britain, Papua New Guinea (32 mm) |
| 13: ? <i>C. sertacinctus</i>
Makira Id., Solomon Is. (37 mm) | 30: <i>C. zebra</i>
Moluccas (33 mm) |
| 14: <i>C. stramineus stramineus</i> ; neotype of
<i>C. stramineus</i> Lamarck
MNHN (34 mm) | 31: <i>C. zebra</i> f. <i>nahoniaraensis</i>
Solomon Is. (27 mm) |
| 15: <i>C. stramineus stramineus</i>
Moluccas (43 mm) | 32: <i>C. zebra</i> f. <i>nahoniaraensis</i>
Solomon Is. (33 mm) |
| 16: <i>C. stramineus stramineus</i>
Moluccas (43 mm) | 33: <i>C. zebra</i> f. <i>nahoniaraensis</i>
Solomon Is. (28 mm) |
| 17: <i>C. stramineus stramineus</i> ; holotype of
<i>C. fuscomaculatus</i> E. A. Smith
BMNH (39 mm) | |

