




Figures 2-6. *C. bulbus*. 2-3: lectotype, 24.0 mm (BMNH); 4: Baia Binga, 24.6 mm (CER); 5-6: Caota Bay, 20.0 and 17.5 mm (SMNS). **Figures 7-11.** *C. aemulus*. 7-10: Mussulo Bay, 36.6, 36.8, 34.1 and 39.3 mm (SMNS); 11: Luanda, 40.1 (CER). **Figures 12-17.** *C. africanus*. 12-13: holotype in Kiener, 30 mm; 14-15: S. Nicolau, 19.9 mm (CER); 16: Ponta das Vacas, 24.6 mm (SMNS); 17: Bentiaba, 17.5 mm (CER). **Figures 18-21.** *C. neoguttatus*. 18: Meva, 39.5 mm (CER); 19: Limagens, 33.9 mm (SMNS); 20-21: Santa Maria, 32.9 and 24.9 mm (SMNS).




Figures 22-26. *C. variegatus*. 22: lectotype, in Kiener, 19 mm; 23: Baia Azul, 24.0 mm (SMNS); 24: Lucira, 26.5 mm (SMNS); 25: Lucira, 25.0 mm (CER); 26: Limagens, 23.3 mm (CER). **Figures 27-33.** *C. zebroides*. 27: Bissonga, 51.2 mm (CER); 28-29: Caota, 35.8 and 47.7 mm (SMNS); 30: shell with periostracum, Caota, 28.5 mm (CER); 31: Limagens, 30.0 mm (SMNS); 32: S. Nicolau, 31.0 mm (SMNS); 33: Chapeu Armado, 29.0 mm (CER). **Figures 34-36.** *C. carnalis*. 34: Santa Maria, 39.6 mm (SMNS); 35: Canoco, 39.9 mm (SMNS); 36: Lucira, 41.1 mm (SMNS). **Figures 37-41.** *C. chytreus*. 37: Salinas Bay, 23.4 mm (SMNS); 38: Limagens, 27.4 mm (SMNS); 39: Lucira, 27.0 mm (CER); 40: Lucira, 25.2 mm (CER); 41: Limagens, 27.0 mm (SMNS).



Figures 42-46. *C. fuscolineatus*. 42-43: holotype, 21.8 mm (BMNH); 44-46: Ponta de Noronha, 26.3, 22.7 and 20.7 mm (SMNS). **Figures 47-49.** *C. cepasi*. 47-49: Chapeu Armado, 39.2, 35.9 and 26.4 mm (SMNS). **Figures 50-51.** *C. nobrei*. 50-51: Lucira, 18.5 and 14.7 mm (CER). **Figures 52-53.** *C. musivus*: 52: Benguela, 22.7 mm (SMNS); 53: Lucira, 25.5 mm (CER). **Figures 54-56.** *C. naranjus*. 54-55: Santa Maria, 18.5 and 18.5 mm (SMNS); 56: Lucira, 15.3 mm (CER). **Figures 57-59.** *C. albuquerquei*. 57-58: Santa Maria, 17.0 and 15.2 mm (SMNS); 59: Santa Maria, 13.2 mm (CER). **Figures 60-61.** *C. bocagei*. 60-61: Lobito, 22.8, 26.0 mm (SMNS).



Figures 62-66. *C. xicoi*. 62-63: *C. hyrcoglyphicus* in Kiener. 64-66: paratypes, Praia Santiago, 23.7, 30.2 and 23.4 mm (SMNS). **Figures 67-71.** *C. gabrielae* n. sp. 67-68: holotype, Chapeu Armado, 24.7 mm (MNCN); 69-71: paratypes, Chapeu Armado, 29.4, 26.2 and 22.7 mm (CER). **Figures 72-76.** *C. micropunctatus* n. sp. 72-73: holotype, Lucira, 32.5 mm (MNCN); 74-75: Limagens, 30.1 mm (SMNS); 76: Limagens, 28.0 (CER). **Figures 77-80.** *C. filmeri* n. sp. 77-78: holotype, Saco Mar, 33.2 (SMNS); 79-80: paratypes, Saco Mar, 22.0 and 30.2 mm (CER). **Figure 81.** *C. dealbatus*, holotype (BMNH).



Figures 82-86. *C. franciscoi* n. sp. 82-83: holotype, Chapeu Armado, 28.4 mm (MNCN); 84-85: paratypes, Chapeu Armado, 34.8 and 36.4 (CER); 86: paratype, Chapeu Armado, 31.3 mm (MNHN), with periostracum.

Figures 87-91. *C. trovaoi* n. sp. 87-88: holotype, Limagens, 38.5 (MNCN); 89-91: paratypes, Limagens, 37.3, 42.4 and 38.0 (SMNS). **Figures 92-96.** *C. flavusalbus* n. sp. 92-93: holotype, Baia das Pipas, 23.7 mm (MNCN); 94: paratype, Baia das Pipas, 25.9 mm (SMNS); 95-96: paratype, Baia das Pipas, 27.4 (SMNS).