

Section C cabritii to cylindraceus

Copyright © 2011.R Filmer Conditions of use. The content of this website is provided for personal and scientific use and may be downloaded for this purpose. It may not be used in whole or part for any commercial activity and publication of any of the content on the internet is limited to the Cone Collector website ($\underline{\text{www.TheConeCollector.com}}$). Authors wishing to publish any of the pictures may do so on a limited basis but should inform M Filmer so that the original owner of the rights to the picture can be acknowledged.

Version 1.0 October 2011.

(mike@mfilmer.fsnet.co.uk)

cabritii Bernardi, 1858.

Published in J. Conch. Paris, 7: p. 377, pl. 13, figs 2 & 2a, (1861, Monographie de Genre Conus: p. 5, pl. 1, fig. 8). Holotype whereabouts unknown, (fig. 29 x 15 mm), (a specimen in MNHN type collection is not the holotype, dimensions $19.8 \times 11.9 \times 11$

Type locality New Caledonia, (Rolland).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. exiquus Lamarck, 1810.



Figure in Monographie.

cacao Ferrario, 1983.

Published in La Piu Grande Enciclopedia delle Conchiglie di Tutto il Mondo, (Italy), vol. 1: p. 146, & figs.

Lectotype (Lauer) in MNHN, (36.6 x 20.9 mm).

Type locality La Somone River Estuary, Senegal, (West Africa), (between rocks).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. mercator Linnaeus, 1758.





caelatus A. Adams, 1854 ("1853").

Published in Proc. zool. Soc. Lond. pt. 21, no. 255: p. 117, no. 5, not figured.

Lectotype (C, M & W) in NHMUK, $(11.9 \times 5.9 \text{ mm})$.

Type locality Chinese seas, [erroneous].

Nomenclatural status, an available name.

Taxonomic status, a synonym (juvenile) of *C. marchionatus* Hinds, 1843.





caerulans Küster, 1837.

Published in Syst. Conch.-Cab. 4 (2): p. 85, no. 69, (1838, pl. 14, figs. 3 & 4.

Lectotype representative (Kohn) figure in Küster, pl. 14, figs 3 & 4, (fig. 41 x 24 mm, orig. desc. 44 x 24 mm).

Type locality St. Thomas Island, Caribbean, (Leeward Islands).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ermineus Born, 1778.



caerulescens "Chemnitz" Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 423, no. 130, not figured.

Lectotype representative (Kohn) figure in Chemnitz, 1795, pl. 183, figs 1776 & 1777. (figs 48 x 23 mm).

Type locality Islands of Moluccas, Indonesia.

Nomenclatural status, an available name, an invalid name, a homonym of *C. coerulescens* Schröter, 1803.

Taxonomic status, a synonym of C. cinereus Hwass, 1792.



<u>caerulescens</u> (variety mediterraneus Hwass) Bucquoy, Dautzenberg & Dollfus, 1882.

Published in Mollusques Marins Roussillon, (Paris), 1 (2): p. 83, pl. 13, fig. 15.

Holotype was in collection Guilliou, present whereabouts unknown, (fig. $22 \times 10 \text{ mm}$).

Type locality Barbary coast, (North Africa), (among sponges). Nomenclatural status, an available name, an invalid name, a homonym of *C. coerulescens* Schröter, 1803 and of Dillwyn, 1817 and of *C. caerulescens* Lamarck, 1810.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.



Figure in Bucquoy et al.

caesius (Röding), 1798.

Published in Mus. Bolten 2: p. 48, no. 616/109, not figured. Lectotype representative (Kohn) figure in Martini, 1773, pl. 52, fig. 580, (fig. 48 x 21 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. magus Linnaeus, 1758.



caffer (Röding), 1798.

Published in Mus. Bolten 2: p. 48, no. 606/100, not figured. Holotype in NBC-ZMA, (27.3 x 16.1 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of *C. coffeae* Gmelin, 1791, same type.





caffer Krauss, 1848.

Published in Die südafrikanischen Mollusken, (Stuttgart): p. 131, pl. 6, fig. 24.

Holotype was in SMNS, destroyed in World War 2, (37 x 16.5 mm), Lectotype (Janus, invalid as this specimen was not part of the type material), in SMNS, (37.5 x 20 mm).

Type locality Cape Province and Natal, South Africa.

Nomenclatural status, an available name, an invalid name, a homonym of *C. caffer* (Röding), 1798.

Taxonomic status, a synonym of C. mozambicus Hwass, 1792.



Figure in Krauss.

caillaudi Kiener, 1845 (1846).

Published in Coq. Viv. 2: pl. 55, fig. 5, (1850, Coq. Viv. 2: p. 285, no. 251, spelt *caillaudii*).

Holotype in MHNN, (51.5 x 24.8 mm, orig. desc. 50 x ? mm). Type locality not mentioned, designated (C, M & W) Cargados Carajos Islands, Indian Ocean.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





Holotype in MNHN Figure in Coq.Viv.

Note the holotype is glued tightly to a card and no picture of the dorsal side is possible.

cailliaudii Jay, 1846.

Published in Ann. Lyceum nat. Hist. 4: p. 169, pl. 10, figs 8a & b.

Holotype was in collection J.C. Jay, transferred to AMNH but no longer identifiable, $(41 \times 17 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. caillaudi* Kiener, 1845 (1846).

Taxonomic status, a synonym (form) of *C. ventricosus* Gmelin, 1791.



Figure 8a in Jay

cakobaui Moolenbeek, Röckel & Bouchet, 2008.

Published in Vita Malacologica 6: p. 37, pl. 1, figs 5 - 7 & pl. 4, fig. 39.

Holotype in MNHN, $(18.9 \times 8.4 \text{ mm})$.

Type locality Fiji, South of Vanua Levu, Somo-somo Strait, (426 - 487 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





caledonicus Hwass in Bruguière, 1792.


Published in Enc. Méth. 1: p. 634, no. 33, (1798, Tab. Enc. pl. 321, fig. 10).

Holotype in MHNG, $(55.5 \times 31 \text{ mm})$.

Type locality the Pacific coasts of New Caledonia, (erroneous), corrected (C, M & W) St. Vincent, Lesser Antilles, (Windward Islands).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. cedonulli* Linnaeus, 1767.





<u>calhetae</u> (subspecies *navarroi* Rolán) Rolán, 1990.

Published in Iberus, Sup. 2: p. 41, pl. 1, fig. 7, pl. 2, fig. 7 & pl. 5, fig. 0.

Holotype in MNCM, $(26.8 \times 15 \text{ mm})$.

Type locality Calheta, eastern Maio Island, Cape Verde Islands, (stones, 1 mtr).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. navarroi Rolán, 1986.





californicus "Hinds" Reeve, 1844.

Published in Conch. Icon. 1 (Conus): pl. 42, sp. 224, (published Jan).

Holotype in NHMUK, $(23.5 \times 12.9 \text{ mm})$.

Type locality California.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





californicus Hinds, 1844.

Published in The Zoology of the Voyage of "H.M.S. Sulphur" 2, Mollusca: p. 7, pl. 1, figs 3 - 5, (published Jul).

Two syntypes were in collection Sir Edward Belcher, present whereabouts unknown, (figs $26 \times 15 \text{ mm}$).

Type locality Bay of Magdalena, California, (sandy bottom, 7 fths).

Nomenclatural status, an available name, an invalid name, a homonym of C. californicus Reeve, 1844, (published six months earlier).

Taxonomic status, a synonym of C. californicus Reeve, 1844.








Figures in Hinds.

(Chelyconus) <u>calliginosus</u> (form kinoshitai Kuroda) Shikama, 1979. Published in Sci. Rep. Yokosuka Cy. Mus. (26): p. 5, pl. 1, figs 15 & 16.

Specimen described was in collection R. Kawamura, to be deposited in NSMT, (51.3 \times 21.3 mm), [missing in May 1990 & April 1995]. Locality Taiwan.

Nomenclatural status, an unavailable name (nomen nudum), described as a form after 1960.

Taxonomic status, none, (a specimen of *C. kinoshitai* Kuroda, 1956).



Figures in Shikama.

camelinus (Röding), 1798.

Published in Mus. Bolten 2: p. 48, no. 618/110, not figured. Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

canaliculatus (Röding), 1798.

Published in Mus. Bolten 2: p. 40, no. 501/26, not figured. Five specimens named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

canaliculatus Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 360, no. 13, not figured.

Lectotype representative (Walls) figure in Tab. Enc. 1798, pl. 325, fig. 9, (fig. 53 x 31 mm, orig. desc. app. 51 x 28 mm). Type locality the coasts of Ceylon (Sri Lanka) and the Nicobar Islands, (Chemnitz), Straits of Malacca, (Bruguière). Nomenclatural status, an available name, an invalid name, a homonym of *C. canaliculatus* Brocchi, 1814 (fossil).

Taxonomic status, an objective junior synonym of *C. malacanus* Hwass, 1792, same type.



"<u>canariensis</u>" (forma *pulcher* Lightfoot) van Rossum, 1997.

Published in World Shells (20): p. 40, figs 8 & 9.

Specimens described whereabouts not indicated, (figs 63 x 36 & 58 x 34 mm, orig. desc. up to 100 mm).

Locality from the Canary Islands to the Gulf of Cameroon, (West Africa), (deep water).

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name, stated to be infrasubspecific after 1960.

Taxonomic status, none, (a specimen (form) of *C. pulcher* Lightfoot, 1786).







cancellatus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 712, no. 107, (1798, Tab. Enc. pl. 338, fig. 1).

Holotype in MHNG, $(39 \times 19 \text{ mm})$.

Type locality Hawaii, (erroneous), corrected (Vink) Off Dry Tortugas, Florida.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





candidus Born, 1778.

Published in Ind. Mus. Caes. Vindob: p. 130, (1780, Test. Mus. Caes. p. 150, pl. 7, fig. 9).

Holotype whereabouts unknown, (fig. 46 x 27.5 mm, orig. desc. app. 46 x 22 mm).

Type locality not known.

Nomenclatural status, an available name, a totally suppressed name, on Official Index of Rejected and Invalid Specific Names in Zoology, ICZN, opinion 757, (1966), see Section 1 B (2). Taxonomic status, none.



Figure in Born.

candidus (Röding), 1798.

Published in Mus. Bolten 2: p. 50, no. 643/129, not figured. Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

candidus Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 97, fig. 1, (1848) (1849), Coq. Viv. 2: p. 214, no. 183).

Holotype was in collection A.C. Bernardi, acquired by M. Gubba and then in MHNH, destroyed in World War 2, $(31 \times ? mm)$, lectotype representative (Lauer) figure in C. V. pl. 97, fig. 1,

Type locality not known.

 $(fig. 31 \times 13 mm).$

Nomenclatural status, an available name, a homonym of *C. candidus* Born, 1778.

Taxonomic status, a synonym (form) of *C. anabathrum* Crosse, 1865.



Figures in Coq. Viv.

canonicus Hwass in Bruguière, 1792.


Published in Enc. Méth. 1: p. 749, no. 143, (1798, Tab. Enc. pl. 345, figs 1 & 5).

Neotype (Kohn) in MHNG, $(53 \times 28.5 \text{ mm})$.

Type locality East Indies, restricted (C, M & W) to the Maldives, (Indian Ocean), neotype locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





canonicus (Röding), 1798.

Published in Mus. Bolten 2: p. 43, no. 535/54, not figured.

Lectotype representative (Kohn) figure in Martini, 1773, pl. 57, fig. 629, (fig. 63 \times 35 mm).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. canonicus* Hwass, 1792.

Taxonomic status, a synonym of C. vexillum Gmelin, 1791.



capitanellus Kuroda, 1935.

Published in Venus, Kyoto, 5 (1): p. 54, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none, (a specimen of *C. capitanellus* Fulton, 1938).

capitanellus Fulton, 1938.

Published in Proc. malac. Soc. Lond. 23 (1): p. 55, pl. 3, figs 1 & 1a.

Holotype in NHMUK, $(36 \times 20 \text{ mm. orig. desc. } 36 \times 19 \text{ mm})$. Type locality Kii, Japan.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





capitaneus Linnaeus, 1758.

Published in Syst. Nat. 10th ed. (1): p. 713, no. 254, not figured.

Lectotype (Kohn) in LSL, $(51 \times 32 \text{ mm})$.

Type locality Asia, restricted (C, M & W) to Larantuka, Flores, Indonesia.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





capitatus (Röding), 1798.

Published in Mus. Bolten 2: p. 51, no. 650/136, not figured.

Specimen named whereabouts unknown, (? $\ensuremath{\mathsf{mm}}\xspace)$.

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

capreolus Röckel, 1985.

Published in Arch. Molluskenk, 115 (4/6): p. 267, pl. 1, figs 1 - 3.

Holotype in NMSF, $(40.7 \times 18.7 \text{ mm})$.

Type locality 50 kms off East coast of Visakhapatnam State, India, (50 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





capricorni van Mol, Tursch & Kempf, 1967.

Published in Annls. Inst. Oceanogr. Univ. Recife, 45 (2) 16: p. 238, fig. 3, pl. 10, fig. 1a - 1b.

Holotype in MNHN, $(49.2 \times 25 \text{ mm})$.

Type locality Puerto Alegre, off Rio Grande do Sul, South Brazil, (135 - 141 mtrs).

Nomenclatural status, an available name.

Taxonomic status, uncertain (incertae cedis), possibly a synonym (fossil form) of *C. cancellatus* Hwass, 1792.





<u>caracanus</u> (subspecies <u>cedonulli</u> Linnaeus) Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 603, no. 1I, (1798, Tab. Enc. pl. 316, fig. 6).

Holotype in MHNG, $(47 \times 25.5 \text{ mm})$.

Type locality coast of Caracas, North of Guyana, South America. Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. trinitarius* Hwass, 1792, (p. 603, no.1D), (a subspecies of *C. mappa* Lightfoot, 1786).





caracteristicus G. Fischer von Waldheim, 1807

Published in Mus. Démidoff 3: p. 139, nos 113 - 116, not figured. Type series was in collection P. de Démidoff, in ZMMU, destroyed by fire in 1812, lectotype representative (Kohn) figure in Chemnitz, 1795, pl. 182, fig. 1761. (fig. 43 x 29 mm), second variety is in ZMMU, $(45 \times 29 \text{ mm})$.

Type locality not mentioned, designated (C, M & W) Java Sea, (Indonesia).

Nomenclatural status, an available name.

Taxonomic status, a valid species, (second variety is a specimen of *C. eburneus* Hwass, 1792).



carcellesi Martins, 1945.

Published in Notas Mus. La Plata, 10 (88): p. 260, figs A - G2. Holotype in MNRJ, (29 x 14 mm).

Type locality Mar del Plata, Buenos Aires Province, Argentina. Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. lemniscatus* Reeve, 1849.





cardinalis Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 632, no. 30, (1798, Tab. Enc. pl. 322, fig. 6).

Probable holotype in MHNG, (27 x 16 mm), type figure designated (Clench) figure in Martini, 1773, pl. 61, fig. 680, (fig. 25 x 15.5 mm).

Type locality the Indian Ocean (erroneous), and the coasts of Santo Domingo and Martinique, corrected (Clench) Tortola Ils., Virgin Islands, restricted (Kohn) to Santo Domingo, (Windward Islands).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cardinalis (Röding), 1798.

Published in Mus. Bolten 2: p. 42, no. 531/50, not figured. Lectotype representative (Walls) figure in Chemnitz, 1788, pl. 144A, fig. L, (fig. 42 x 24 mm).

Type locality not mentioned, (West Indies, Chemnitz).

Nomenclatural status, an available name, an invalid name, a homonym of *C. cardinalis* Hwass, 1792.

Taxonomic status, an objective junior synonym of *C. daucus* Hwass, 1792, same type.



cardonensis (subspecies puncticulatus Hwass) Vink, 1990.

Published in Conchiglia 22 (253 - 255): p. 16, fig. 39C.

Holotype in MHNG, $(20.9 \times 11.9 \text{ mm})$.

Type locality Punta Cardon, Paraguana Peninsula, Venezuela, (mud, 0 - 10 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. puncticulatus* Hwass, 1792.





cargilei Coltro, 2004.

Published in Strombus no. 011: p. 5, pl. 5, fig. A, pl. 13 figs H1

& P1 - 11, pl. 16, fig. H.

Holotype in MZUSP, $(20.8 \times 10.6 \text{ mm})$.

Type locality 75 kms NNE Abrolhos Archipelago off Alcobaça, Bahia State, Brazil, (rubble and coral sand bottom at 20-35 mtrs on offshore reefs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. archetyphus* Crosse, 1865.





<u>caribaensis</u> (subspecies <u>centurio</u> Born) Nowell-Usticke, 1968. Published in Caribbean Cones from St. Croix and the Lesser Antilles: p. 13, pl. 2, fig. 998.

Lectotype (Nowell-Usticke 1971) in AMNH, (31.5 x 18 mm).

Type locality Krause's Bay, St. Croix, (Virgin Islands) and South of Barbados, (Leeward Islands).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. centurio Born, 1778.





caribbaeus Clench, 1942.

Published in Johnsonia 1 (6): p. 23, pl. 11, figs 4 & 5.

Holotype in MCZ, $(31 \times 15 \text{ mm})$.

Type locality off Palm Beach County, Florida.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. flavescens* Sowerby I & II, 1834.





carinata (variety mediterraneus Hwass) Bucquoy, Dautzenberg &
Dollfus, 1882.

Published in Mollusques Marins Roussillon, (Paris), 1 (2): p. 82, pl. 13, figs 16 & 17.

Type series was in collection Guilliou, present whereabouts unknown, lectotype representative (C, M & W) larger figure in Bucquoy, Dautzenberg & Dollfus, pl. 13, fig. 17, (fig. 22 x 12 mm).

Type locality Barbary coast, (North Africa), (on sponges), (Guilliou).

Nomenclatural status, an available name, an invalid name, a homonym of *C. carinatus* Swainson, 1822.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.




carinatus Swainson, 1822.

Published in Zoological Illustrations (1) 2: pl. 112. Holotype was in collection C. Dubois, present whereabouts unknown, holotype representative (Kohn) figure in Swainson, pl. 112, (fig. $69 \times 33.5 \text{ mm}$).

Type locality Asiatic Ocean, restricted (C, M & W) to the Philippines.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. magus Linnaeus, 1758.



carioca Petuch, 1986.

Published in Proc. biol. Soc. Wash. 99 (1): p. 9, figs 1 & 2. Holotype in MOFU, (52 x 24 mm).

Type locality Cape Frio, Rio de Janeiro, Brazil, (trawled, 100 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. sanderi* Wils & Moolenbeek, 1975.





carmeli Tenison-Woods, 1877 ("1876").

Published in Pap. Pro. R. Soc. Tasm: p. 134, no. 7, not figured. Holotype in TMH, (22 x 9 mm).

Type locality North coast of Tasmania, (Australia), (Petterd). Nomenclatural status, an available name.

Taxonomic status, a synonym (juvenile) of C. anemone Lamarck, 1810.





carnalis Sowerby III, 1879 ("1878").

Published in Proc. zool. Soc. Lond. unnumbered (46), pt. 4: p. 796, pl. 48, fig. 2,

Holotype in NMWC, $(48.1 \times 25.1 \text{ mm})$.

Type locality not known, designated (C, M & W) Santa Maria Bay, Angola.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





carota (Röding), 1798.

Published in Mus. Bolten 2: p. 47, no. 597/93, not figured. Lectotype representative (C, M & W) figure in Martini, 1773, pl. 52, fig. 581, (fig. 52×25 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. spectrum* Linnaeus, 1758



carpenteri Crosse, 1865.

Published in J. Conch. Paris, 13: p. 302, no. 3, pl. 9, fig. 1. Holotype in NHMUK, (46 x 25 mm).

Type locality New Guinea, Oceania.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. litoglyphus* Hwass, 1792.





cassis "Meuschen" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 355, no. 4, not figured.

Specimen named whereabouts unknown, cited species in Meuschen, 1766, p. 360, no. 1041, not figured.

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. arachnoideus* Gmelin, 1791.

Taxonomic status, none, (a specimen of *C. araneosus* Lightfoot, 1786).

cassis "Crosse" O. & J.B. Lozet, 1978.

Published in Le Peigne de Venus, Coquillages du Monde Entier: p. 96, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none, (stated to be a specimen of C.

"broderipi" Kiener but Kiener did not introduce this name, presumably an error for *C. broderipii* Reeve, 1843).

castanea
1933.
(variety mediterraneus Hwass) "Danilo & Sandri" Coen,

Published in Memorie R. Comit. talassogr. ital. (192): p. 70, not figured.

Specimens named were in collection Danilo Sandri in MCG.

destroyed in World War 2, (? mm).

Locality East coast of the Adriatic.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none, (a specimen of *C. ventricosus* Gmelin, 1791).

castaneofasciata
1937.
"Sowerby" (variety amadis Gmelin) Dautzenberg,

Published in Mem. Mus. r. Hist. nat. Belg. 2 (18): p. 15, not figured. Lectotype representative (C, M & W) figure in Kiener (1847) pl. 84, fig. 2, (80 x 40.5 mm), (this specimen was in the collections of Delessert and Sollier in MHNG). Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a subspecies of C. amadis Gmelin, 1791.



Figure in Coq. Viv. 2.

<u>castaneus</u> Kiener, 1845 (1848).

Published in Coq. Viv. 2: pl. 104, fig. 3, (1848) (1849), Coq. Viv. 2: p. 209, no. 178).

Holotype in NHMUK, $(45 \times 23.5 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. cingulatus Lamarck, 1810.





castrensis Gould, 1842.

Published in Boston J. nat. Hist. 4 (1), back cover: (1843, Proc. Boston Soc. nat. Hist. 5 (1): p. 138).

Holotype whereabouts unknown, (75 x 38 mm), lectotype (Johnson, invalid as it is not part of type material) owned by NYSM, on permanent loan to MCZ, (42.5 x 23.4 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of $C.\ thalassiarchus$ Sowerby I & II, 1834.





Specimen in MCZ.

<u>castus</u> Reeve, 1844 ("1843").

Published in Conch. Icon. 1 (*Conus*): pl. 47, sp. 267, (published Feb), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 180, not figured, published Jun).

Holotype in NHMUK, $(19 \times 11 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (subadult) of C. daucus Hwass, 1792.





catenatus
(variety protheus Lightfoot) Lightfoot, 1786.
Publish in Cat. Portland Mus: p. 158, no. 3452, not figured.
Specimen(s) named sold at auction, present whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

catenatus Sowerby III, 1879 ("1878").


Published in Proc. zool. Soc. Lond. unnumbered (46), pt. 4: p. 796, pl. 48, fig. 3. (In Thes. Conch. 5, 1887, pl.507 (29) and in the index p.277, Sowerby miss-spelled it *C. concatenatus*) Holotype in NMWC, (31.4 x 15 mm).

Type locality Panama, (East coast).

Nomenclatural status, an available name, an invalid name, a homonym of *C. catenatus* Sowerby, 1849 (fossil), renamed *C. desmotus* Tomlin, 1937,

Taxonomic status, a synonym of *C. granarius* Kiener, 1845 (1847)





catus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 707, no. 99, (1798, Tab. Enc. pl. 332, figs 3, 4 & 7).

Lectotype (Kohn) in MHNG, $(40 \times 23 \text{ mm})$.

Type locality Santo Domingo Isle, Martinique, (erroneous) and Ile de France, (Mauritius), restricted (Kohn) to Mauritius.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cavailloni Fenaux, 1942

Published in Bull. Inst. oceanogr. Monaco, (814): p. 4, fig. 12. Type series was in collection Valero, present whereabouts unknown, (fig. $47.5 \times 18.5 \text{ mm}$).

Type locality Bermuda, [erroneous].

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. inscriptus Reeve, 1843.



Figure in Bull. Inst.

caysalensis L. Raybaudi & Prati, 1994.

Published in World Shells (8): p. 8 & figs.

Holotype in IMT, $(18.8 \times 10.5 \text{ mm})$, not present (apparently never deposited and now probably lost Raybaudi G. pers. corr.).

Type locality Cay Sal Bank, South western Bahamas, (coral reefs, 5 - 10 fths).

Nomenclatural status, an available name.

Taxonomic status, a valid species.



Figure of holotype.

cebuensis (subspecies proximus Sowerby) Wils, 1990.

Published in Gloria Maris 29 (2): p. 25. figs 4 - 6.

Holotype in IRSN, $(37.1 \times 19.2 \text{ mm})$.

Type locality Punta Engano, Mactan Isl., Cebu, Philippines, (2 - 40 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cebuganus da Motta & R. Martin, 1982.

Published in Carfel 4 (3): p. 1, fig. 1.

Holotype in MHNG, $(36 \times 14.5 \text{ mm})$.

Type locality Visayan Sea, between Malapascua Isl. and Bantayan Isl. and northward, Cebu, Philippines, (dredged, sandy - mud bottom, 20 - 40 fths).

Nomenclatural status, an available name.

Taxonomic status, a synonym (semi adult form) of *C. australis* Holten, 1802.





cecilei Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 98, fig. 4, & var. pl. 107, fig. 3, (1850, Coq. Viv. 2: p. 286, no. 252).

Holotype was in collection Largilliert, present whereabouts unknown, (42 x ? mm), (fig. 44 x 20 mm).

Type locality China Sea.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. furvus Reeve, 1843.



Figures in Coq. Viv.

ceciliae "Chenu" Crosse, 1858.

Described in J. Conch. Paris, 7: p. 381, pl. 14, fig. 5. Holotype in MHNG, (33 x 19.5 mm, orig. desc. 36 x 21 mm). Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (ridged form) of *C. capitaneus* Linnaeus, 1758.





cedonulli (variety ammiralis Linnaeus) Linnaeus, 1767.
Published in Syst. Nat. 12th ed. (1): p. 1167, no. 298E, not figured.

Holotype was in collection J.B. de la Faille, sent to the King of Denmark and later bought by P. Lyonnet and sold at auction, present whereabouts unknown, (? mm), lectotype representative (Kohn) and holotype representative (C, M & W), figure in Seba, 1758, pl. 48, fig. 8, (fig. $56 \times 30 \text{ mm}$).

Type locality South American Ocean, restricted (Vink & von Cosel) to St. Vincent, (Windward Islands).

Nomenclatural status, an available name.

Taxonomic status, a valid species.



cedonulli Perry, 1811.

Published in Conchology: pl. 24, fig. 1.

Type series whereabouts unknown, $(67 \times 36 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. cedonulli* Linnaeus, 1767.

Taxonomic status, a synonym of C. cedonulli Linnaeus, 1767.




Figure in Perry.

centurio Born, 1778.

Published in Ind. Mus. Caes. Vindob: p. 133, (1780, Test. Mus.

Caes. p. 153, pl. 7, fig. 10).

Holotype in NHMW, $(35.5 \times 21.2 \text{ mm})$.

Type locality not known, designated (Clench) Puerto Plata, Santo Domingo, (Dominican Republic).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cepasi Trovão, 1975.

Published in Bolm. Cent. port. Activ. Subaqua. 4 (1): p. 3, pl. 1, figs 1a - 1f.

Holotype in CPAS, $(46.6 \times 26.4 \text{ mm})$

Type locality (near São Nicolau) Angola, (rocks).

Nomenclatural status, an available name.

Taxonomic status, a valid species.

cereolus (Röding), 1798.

Published in Mus. Bolten 2: p. 44, no. 557/67, not figured.

Lectotype representative (Kohn) figure in Chemnitz, 1788, pl. 140, fig. 1301, (fig. 56 x 27 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. monile Hwass, 1792.



cerinus Reeve, 1848.

Published in Conch. Icon. 1 (Conus): Sup. pl. 3, sp. 283a & b. Holotype in NHMUK, (29 x 15 mm).

Type locality Island of Mindanao, Philippines.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. boeticus Reeve, 1844.





(Coronaxis) cernicus "Barclay MS" H. Adams, 1869.

Published in Proc. zool. Soc. Lond. unnumbered (37), pt. 2: p. 272, pl. 19, fig. 1.

Holotype was in collection D.W. Barclay, sold in 1891 and again in 1898, present whereabouts unknown, (fig. $25 \times 12 \text{ mm}$).

Type locality Barkly Island, Mauritius, (Barclay).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of $C.\ balteatus$ Sowerby I & II, 1833.



Figure in Adams

cernohorskyi da Motta, 1983.

Published in Publcoes. Occ. Soc. port. malac. (2): p. 2, figs 9 - 13.


Holotype in MHNG, $(46.4 \times 26.8 \text{ mm})$.

Type locality Borogon, East Samar, Philippines, (diving, shallow water).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. magus Linnaeus, 1758.





ceruttii Cargile, 1997.

Published in Conchiglia 29 (282): p. 48 & figs.

Holotype in SBMNH, $(31.6 \times 15.6 \text{ mm})$.

Type locality Isla Grande de Maiz, Nicaragua, (East coast), (SCUBA, 10 - 25 mtrs, sand bottom).

Nomenclatural status, an available name.

Taxonomic status, a valid species.



cervulus "Meuschen" Dillwyn, 1817

Published in Desc. Cat. Recent Shells 1: p. 380, no. 45, not figured.

Specimen named whereabouts unknown, cited species in Meuschen, 1766, p. 362, no. 1056, not figured.

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. nobilis* Linnaeus, 1758.

Taxonomic status, none, (a specimen of C. nobilis Linnaeus, 1758).

cervus Lamarck, 1822.

Published in Hist. Nat. Anim. sans Vert. 7: p. 510, no. 151, not figured.

Holotype in MHNG, $(94 \times 44 \text{ mm})$.

Type locality not known, designated (C, M & W) the Moluccas, Indonesia.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cervus Sowerby I & II, 1838.

Published in Conch. Ill. (Conus): pl. 147, fig. 94.

Lectotype (C, M & W) in IRSN, (51.2 x 27.4 mm).

Type locality Swan River, Australia, (Reeve, 1843), (erroneous), corrected (C, M & W) Djibouti, (Somalia).

Nomenclatural status, an available name, an invalid name, a homonym of *C. cervus* Lamarck, 1822, renamed and redescribed as *C. deshayesii* Reeve, 1843, (Sowerby misidentified this shell). Taxonomic status a synonym of *C. cuvieri* Crosse, 1858,





ceylanensis Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 636, no number, (1798, Tab. Enc. pl. 322, fig. 10).

Lectotype representative (Kohn) figure in Tab. Enc. pl. 322, fig. 10, (fig. 18 x 10 mm). neotype (C, M & W) in NBC-ZMA, (10 x 6.2 mm).

Type locality coasts of Ceylon Island, (Sri Lanka), neotype locality, Hikkaduwa, Sri Lanka.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. musicus* Hwass, 1792, (p. 629, no. 25).



ceylonicus "Chemnitz" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 401, no. 85, not figured.

Specimen named whereabouts unknown, cited figure in Chemnitz, 1788, pl. 142, fig. 1318, (fig. $42 \times 23 \text{ mm}$).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. obesus* Hwass, 1792.

Taxonomic status, none, (figure cited is lectotype representative of C. meningeus (Röding), 1798), (a synonym (form) of C. zeylanicus Gmelin, 1791).



Cited figure in Chemnitz

<u>ceylonicus</u> "Chemnitz" Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Emend. p. 1, not figured. Lectotype of C. obesus Hwass, 1792 in MHNG, (65 x 45.5 mm). Type locality of C. obesus East Indies.

Nomenclatural status, an available name, a new replacement name (nomen novum), for $C.\ obesus$ Hwass, 1792, an invalid name, unjustified as Hwass's name is available.

Taxonomic status, a synonym of C. zeylanicus Gmelin, 1791.





Lectotype of *C. obesus*.

ceylonicus "Chemnitz" Sowerby II, 1857.

Published in Thes. Conch. 3 (*Conus*), pt. 17: p. 8, no. 56, pl. 2 (188), figs 22 & 16 (var).

Type series whereabouts unknown, (fig. 22, 33 x 22 mm).

Type locality Ceylon, (Sri Lanka).

Nomenclatural status, an available name, an invalid name, a homonym of *C. ceylonicus* Reeve, 1849.

Taxonomic status, a synonym of C. zeylanicus Gmelin, 1791.





Figures 22 & 16 in Thes. Conch. 3.

chaldaeus (Röding), 1798.

Published in Mus. Bolten 2: p. 42, no. 525/47, not figured. Lectotype representative (Kohn) figure in Knorr, 1768, pl. 4, fig. 2, (fig. $36 \times 22 \text{ mm}$).

Type locality not mentioned, designated (C, M & W) the Moluccas, Indonesia.

Nomenclatural status, an available name.

Taxonomic status, a valid species.



characteristicus Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 367, no. 26, not figured.

Holotype representative (Kohn) figure in Chemnitz, 1795, pl. 182, figs 1760 - 1761, (figs. 43 x 29 mm, orig. desc. app. 45 x 30 mm).

Type locality West Indian seas, on the coast of the Island of St. Bartholomew, (Chemnitz), (erroneous), corrected (C, M & W) Moluccas, (Indonesia).

Nomenclatural status, an available name, an invalid name, a homonym of *C. caracteristicus* Fischer, 1807.

Taxonomic status, an objective junior synonym of *C. caracteristicus* Fischer, 1807, same type.



chemnitzii Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 363, no. 18, not figured.

Holotype representative (Kohn) figure in Chemnitz, 1795, pl. 182, figs 1764 - 1765, (figs 45×28 mm, orig. desc. 45×25 mm). Type locality the coasts of Ceylon, (Sri Lanka), (Chemnitz). Nomenclatural status, an available name.

Taxonomic status, a synonym of C. rattus Hwass, 1792.



chenui Crosse, 1857 ("1856").

Published in J. Conch. Paris, 6: p. 381, pl. 11, figs 3 & 4, (1858, Revue Mag. Zool. 2 (10): p. 120).

A syntype in MNHN, (49.1 x 26.9 mm. figured), orig. desc. 50 x 28 mm), (three probable syntypes in NHMUK)

Type locality New Caledonia.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. ferrugineus* Hwass, 1792.





Figured syntype in MNHN.

<u>chersoideus</u> "Ch" (variety mediterraneus Hwass) Nardo, 1847. Published in Sinon. Moderna, pt. 2: p. 41/42, sp. 10, fig. 519. Type series was in collection S. Chiereghin, present whereabouts unknown, (fig. ? mm).

Type locality Cherso Island, Adriatic Sea, (on the beach). Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. ventricosus* Gmelin, 1791.




Figure in Chiereghin MSS.

(Taranteconus) chiangi Azuma, 1972.

Published in Venus, Kyoto, 31 (2): p. 56, figs 5 & 6, (published Jul).

Holotype in BRIT, $(17.8 \times 10 \text{ mm})$.

Type locality South China Sea, (on coral bed, about 200 fths).

Nomenclatural status, an available name.





chiapponorum Lorenz, 2004.

Published in Visaya vol. 1, no. 2: p. 20, pl. 2. (published November)

Holotype in MNHN, $(41.4 \times 22 \text{ mm})$.

Type locality From Fort Dauphin to Lavonono [Lavanono ?], southern Madagascar, (beach).

Nomenclatural status, an available name.

Taxonomic status, a valid species, possibly a synonym (colour form) of *C. solangeae* Bozzetti, 2004 (May).





chinensis (Röding), 1798.

Published in Mus. Bolten 2: p. 47, no. 595/92, not figured. Lectotype representative (Walls) figures in Chemnitz, 1788, pl. 144A, figs g & h, (fig. g, 38 x 21 & fig. h, 39 x 23 mm). Type locality not mentioned, (Chemnitz indicates this specimen was from Sumatra)

Nomenclatural status, an available name. Taxonomic status, doubtful (nomen dubium).



(Rhizoconus ?) chinoi Shikama, 1970.

Published in Venus, Kyoto, 29 (4): p. 115, figs 1 - 4.

Holotype in NSMT, $(32 \times 17.6 \text{ mm})$.

Type locality Ogokuda, Shiono-misaki, Wakayama Prefecture, Japan. Nomenclatural status, an available name.

Taxonomic status, a synonym (subadult) of C. distans Hwass, 1792.





cholmondeleyi Melvill, 1900.

Published in J. Conch. Lond. 9 (10): p. 308, & fig.

Holotype in MMMU, $(43 \times 18 \text{ mm, orig. desc. } 45 \times 12 \text{ mm an error})$. Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. textile Linnaeus, 1758.





chrysalis (Röding), 1798.

Published in Mus. Bolten 2: p. 50, no. 645/131, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

chrysocestus Berry, 1968.

Published in Leafl. Malac. 1 (25): p. 157, not figured.

Holotype in SBMNH, $(45.3 \times 24.3 \text{ mm})$.

Type locality Morro Colorada, Sonora, Mexico, (West coast), (trawled, 30 - 45 fths).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. xanthicus Dall, 1910.





chusaki da Motta, 1978.

Published in Centre Thai Natural Study, 6 (1), (Bangkok): p. 8 & figs.

Holotype in MHNG, $(65 \times 30 \text{ mm})$.

Type locality off Raya Island, Phuket, Thailand, (sand & rocks, 80 - 120 ft).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. floridus Sowerby II, 1858, recognized by da Motta in 1995, (a synonym (form) of C. striatus Linnaeus, 1758).





chytreus
("1883").
(variety figulinus Linnaeus) Melvill in Tryon, 1884

Published in Manual Conchology, (1) 6, (Philadelphia,

Pennsylvania, U.S.A.): p. 17, pl. 27, fig. 1.

Holotype in NMWC, $(16.6 \times 10 \text{ mm})$.

Type locality not mentioned, designated (C, M & W) Lucira Bay, Angola.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cibieli Kiener, 1845 (1849).

Published in Coq. Viv. 2: pl. 107, fig. 2, (1849 (1850), Coq. Viv. 2: p. 242, no. 210, spelt *cibielii*).

Holotype wherebouts unknown, was in collection Cibiel then probably in MHNH destroyed in World War 2, (orig. desc. 28 x 17 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of $C.\ gladiator$ Broderip, 1833.



Figure in Coq. Viv.

cidaris Kiener, 1845 (1846).

Published in Coq. Viv. 2: pl. 63, figs 1 & 1a, (1846) (1847), Coq. Viv. 2: p. 57, no. 47).

Lectotype (Richard) in MNHN, (28.4 x 14.1 mm. orig. desc. 25 x ? mm).

Type locality Indian Ocean, [erroneous].

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. magellanicus Hwass, 1792.





ciderryi da Motta, 1985.

Published in Publcoes. Occ. Soc. port. malac. (4): p. 4, pl. 1, figs 3a & b.

Holotype in MHNG, $(31 \times 14.4 \text{ mm})$.

Type locality Straits of Taiwan, (trawled).

Nomenclatural status, an available name.





cinamomeus (Röding), 1798.

Published in Mus. Bolten 2: p. 43, no. 534/53, not figured. Lectotype representative (Kohn) figure in Martini, 1773, pl. 57, fig. 631, (fig. 52×25 mm), (the figures cited of 680 & 681 are erroneous and should be 630 & 631).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. litoglyphus* Hwass, 1792.



cinctus "S" Lightfoot, 1786.

Published in Cat. Portland Mus: p. 149, no. 3280, not figured. Specimen(s) named sold at auction, present whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

cinctus Bosc, 1801.

Published in Hist. Nat. Coq. 5: p. 140, pl. 40, fig. 2.

Holotype was in collection Famin in MHNH, destroyed in World War 2, holotype representative (Kohn) figure in Bosc, pl. 40, fig. 2. (fig. $26 \times 13 \text{ mm}$).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.



cinctus Link, 1807.

Published in Beschr. Nat.-Samml. Univ. 3: p. 102, not figured.

Type series whereabouts unknown, (? mm).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. cinctus* Bosc, 1801.

Taxonomic status, doubtful (nomen dubium), lost type, no locality or figure and an inadequate description.

cinctus Swainson, 1822.


Published in Zoological Illustrations (1) 2: pl. 110.

Holotype was in collection C. Dubois, (ex Mrs. Angus), present whereabouts unknown, lectotype representative (Kohn) figure in Swainson, pl. 110, (fig. $65 \times 35 \text{ mm}$).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. cinctus* Bosc. 1801, of Link, 1807 and of Borson, 1820 (fossil), renamed *C. circumactus* Iredale, 1929.

Taxonomic status, a synonym of C. circumactus Iredale, 1929.



cinctus Valenciennes, 1832.

Published in Humboldt & Bonpland, Voyage dans l'Intérieur de l'Amérique dans les années 1799 - 1804 Recueil d'Observations de Zoologie et Anatomie Comparée, pt. 2, book 14: p. 337, not figured.

Holotype whereabouts unknown, (34 x ? mm).

Type locality Acapulco, (West Mexico).

Nomenclatural status, an available name, an invalid name, a homonym of *C. cinctus* Bosc, 1801, of Link, 1807, of Borson, 1820 (fossil) and of Swainson, 1822.

Taxonomic status, a synonym of C. emarginatus Reeve, 1844.

cinerarius (Röding), 1798.

Published in Mus. Bolten 2: p. 46, no. 590/89, not figured. Lectotype representative (Kohn) figure in Knorr, 1768, pl. 16, fig. 3. (fig. 43 x 21 mm)

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. monachus Linnaeus, 1758.



cinereus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 673, no. 67, (1798, Tab. Enc. pl. 331, figs 3, 4 & 7).

Lectotype (Kohn) in MHNG, $(48 \times 24 \text{ mm})$.

Type locality Asiatic Ocean, restricted (C, M & W) to Moluccas, Indonesia.

Nomenclatural status, an available name.





cinereus (Röding), 1798.

Published in Mus. Bolten 2: p. 46, no. 589/88, not figured. Lectotype representative (C, M & W) figure in Martini, 1773, pl. 52, fig. 578, (fig. 53 x 26 mm).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of $C.\ cinereus$ Hwass, 1792.

Taxonomic status, a synonym of C. cinereus Hwass, 1792.



cinereus Schröter, 1803.

Published in Arch. Zool. Zoot. 3 (2): p. 72, not figured. Holotype whereabouts unknown, (app. 15 x ? mm).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of C. cinereus Hwass, 1792 and of (Röding), 1798. Taxonomic status, doubtful (nomen dubium), lost holotype, no figure or locality and an inadequate description.

cingulatus Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 274, no. 84, not figured.

Neotype (Walls) in MHNG, (44 x 21 mm).

Type locality Asiatic Ocean, neotype, Indian Ocean, (Kiener), (both erroneous), corrected (C, M & W) Santa Marta, Colombia, (East coast).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cingulatus Sowerby I, 1825.

Published in Cat. Shells Tankerville: p. 34 (no entry), App. no. 2467, not figured, (1834, C. I. pl. 152, fig. 108). Holotype was in collection Earl of Tankerville, present

whereabouts unknown, (51 x 28 mm), holotype representative (Kohn) figure in Conch. Ill. (*Conus*), 1834, pl. 152, fig. 108. (fig. 50.5 x 28 mm).

Type locality not known.

Nomenclatural status, an available name, an invalid name, a homonym of $C.\ cingulatus$ Lamarck, 1810 and of Schlotheim, 1820 (fossil).

Taxonomic status, a synonym of C. adamsonii Broderip, 1833.



cingulum Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3378, no. 72, not figured.

Holotype was in collection Mrs. Fordyce, present whereabouts unknown, holotype representative (Kohn) figure in Martyn, 1789, pl. 39, (fig. $85 \times 52 \text{ mm}$).

Type locality Tonga Islands, (Martyn).

Nomenclatural status, an available name.

Taxonomic status, a synonym (teratological specimen) of *C. quercinus* Lightfoot, 1786.



circae "Chemnitz" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 422, no. 136, not figured.

Specimen named in ZMUC, $(45.9 \times 20.3 \text{ mm})$.

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. magus* Linnaeus, 1758.

Taxonomic status, none, (specimens of C. magus Linnaeus, 1758).



Specimen named in ZMUC.

circae "Chemnitz" Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Emend. p. 2, not figured. Specimen named in ZMUC, ($45.9 \times 20.3 \text{ mm}$).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. magus* Linnaeus, 1758.

Taxonomic status, none, (a specimen of C. magus Linnaeus, 1758).



Specimen named in ZMUC.

circae "Chemnitz" Sowerby II, 1858.

Published in Thes. Conch. 3, (Conus) pt. 18: p. 39, no. 338, pl. 21 (207), figs 513 & 514, variety pl. 22 (208), fig. 525. Syntypes were in collection H. Cuming, present whereabouts unknown, (fig. 513, 52 x 26 & fig. 514, 51 x 28 mm). Type locality Philippines, (Cuming).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. magus Linnaeus, 1758.



Figures in Thes. Conch. 3.

circumactus Iredale, 1929.


Published in Mem. Qd. Mus. 9 (1): p. 281.

Lectotype representative of $C.\ cinctus$ Swainson, 1822 figure in Swainson pl. 110, (fig. 65 x 35 mm).

Type locality of *C. cinctus* not mentioned.

Nomenclatural status, an available name, a new replacement name (nomen novum) for *C. cinctus* Swainson, 1822.

Taxonomic status, a valid species.



Lectotype representative of C. cinctus.

circumcisus Born, 1778.

Published in Ind. Mus. Caes. Vindob: p. 147, (1880, Test. Mus. Caes. p. 163, not figured).

Lectotype (Kohn) in MHNG, $(71 \times 27 \text{ mm})$.

Type locality not known, designated (C, M & W) Moluccas, Indonesia.

Nomenclatural status, an available name.





circumclausus Fenaux, 1942

Published in Bull. Inst. oceanogr. Monaco, (814): p. 3, fig. 8. Type series whereabouts unknown, (fig. 18 x 10 mm).

Type locality Mauritius.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. cernicus* Adams, 1869, (a synonym of *C. balteatus* Sowerby I & II, 1833).

circumductus (Röding), 1798.

Published in Mus. Bolten 2: p. 48, no. 611/105, not figured. Two specimens named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

<u>circumpunctatus</u> (variety <u>caribbaeus</u> Clench) Nowell-Usticke, 1968. Published in Caribbean Cones from St. Croix and the Lesser Antilles: p. 15, pl. 2, fig. 1002. (raised to species status by Nowell-Usticke 1971, p. 18, pl. 3, fig. 1002). Lectotype (Nowell-Usticke 1971) in AMNH, (33.5 x 19 mm). Type locality Anguilla, Haiti, St. Croix and Antigua, restricted (Nowell-Usticke 1971) to Anguilla, (Leeward Islands). Nomenclatural status, an available name. Taxonomic status, a synonym (colour form) of *C. daucus* Hwass, 1792.





circumsignatus Crosse, 1865.

Published in \overline{J} . Conch. Paris, 13: p. 311, no. 10, pl. 10, fig. 4. Holotype in NHMUK, 30 x 15 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of $C.\ floccatus$ Sowerby I & II, 1841.





circumvolutus (Röding), 1798.

Published in Mus. Bolten 2: p. 48, no. 613/106, not figured. Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

citrinus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3389, no. 37, not figured.

Holotype was in collection B. Feldmann, present whereabouts unknown, type figure designated (Clench) figure in Martini, 1773, pl. 61, fig. 681, (fig. $26 \times 15 \text{ mm}$).

Type locality Curacas, (Curacao), Netherlands Antilles.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. regius* Gmelin, 1791, Clench and Abbott separately redescribed and established *C. regius* Gmelin, 1791 as the nominate species.



citrinus Kiener, 1845 (1846).

Published in Coq. Viv. 2: pl. 59, fig. 6, (1849) (1850), Coq. Viv. 2: p. 248, no. 216).

Holotype was in collection Declerc, present whereabouts unknown, (40 x ? mm), (fig. 43.5 x 23 mm).

Type locality the coasts of Ceylon, (Sri Lanka), Indian Ocean, [erroneous].

Nomenclatural status, an available name, an invalid name, a homonym of *C. citrinus* Gmelin, 1791, renamed *C. kieneri* Crosse, 1858.

Taxonomic status, a synonym of C. tinianus nHwass, 1792.



Figure in Coq. Viv.

(Rhizoconus) clandestinatous Shikama, 1979.

Published in Sci. Rep. Yokosuka Cy. Mus. (26): p. 2, pl. 1, figs 3 & 4.

Holotype in NSMT, $(43.5 \times 22.5 \text{ mm})$.

Type locality South China Sea.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. voluminalis Reeve, 1843.





clandestinus "Chemnitz" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 422, no. 136, not figured.

Specimen named whereabouts unknown, cited figure in Chemnitz, 1788, pl. 140, fig. 1296, (fig. 52 x 24 mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. magus* Linnaeus, 1758.

Taxonomic status, none, (a specimen of C. magus Linnaeus, 1758).



Cited figure in Chemnitz

clandestinus "Chemnitz" Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Emend. p. 2, not figured. Specimen named whereabouts unknown, cited figure in Chemnitz, 1788, pl. 140, fig. 1296, (fig. 52 x 24 mm).

Typer locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of $C.\ magus$ Linnaeus, 1758.

Taxonomic status, none, (a specimen of C. magus Linnaeus, 1758).



Cited figure in Chemnitz

(Rhizoconus) clandestinus "Chemnitz" Shikama, 1979.

Published in Sci. Rep. Yokosuka Cy. Mus. (26): p. 3, pl. 1, figs 9 & 10.

Specimen described holotype of C. purpuratus Shikama, 1979 in NSMT, (45.7 x 21.6 mm).

Type locality of *C. purpuratus* off Taiwan, South China Sea. Nomenclatural status, an unavailable name (nomen nudum), Shikama used this unavailable name in conjunction with a new subspecies, *C. purpuratus*, later raised to species status and renamed *C. shikamai* Coomans, Moolenbeek & Wils, 1985.

Taxonomic status, a synonym of *C.shikamai* Coomans, Moolenbeek & Wils 1985.





Holotype of C. purpuratus.

clarki Rehder & Abbott, 1951.

Published in J. Wash. Acad. Sci. 41 (1): p. 22, figs 1-6. Holotype in USNM, (34 x 16 mm. orig. desc. 36 x 16 mm. specimen is now broken at the base).

Type locality 50 miles S, S-W of Marsh Island, Iberia County, Louisiana, (dredged, 29 fths).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. armiger Crosse, 1858.





clarus E.A. Smith, 1881.

Published in Ann. Mag. nat. Hist. 5 (8): p. 442, not figured. Holotype in NHMUK, $(26.7 \times 14 \text{ mm})$.

Type locality western Australia, (Sowerby).

Nomenclatural status, an available name.





classiarius Hwass in Bruguière, 1792

Published in Enc. Méth. 1: p. 705, no. 96, (1798, Tab. Enc. pl. 335, fig. 7).

Holotype whereabouts unknown, $(27 \times 17 \text{ mm})$.

Type locality Indian Ocean.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost type, a discrepancy between the figure and the description and a question mark added to the figure.



Figure in Enc. Méth.

claudiae Tenorio & Afonso, 2004.

Published in Visaya vol. 1, no. 2: p. 27, pl. 3.

Holotype in MNCM, $(17.2 \times 9.6 \text{ mm})$.

Type locality Praia Real, North coast of Maio Island, Cape Verde Islands, (under rocks or coral slabs on sandy bottom, 1-2 mtrs).

Nomenclatural status, an available name.





clavatus (Röding), 1798.

Published in Mus. Bolten 2: p. 46, no. 588/87, not figured. Lectotype representative (C, M & W) figure in Martini, 1773, pl. 52, fig. 578, (fig. 53 x 26 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of C. cinereus (Röding), 1798, (p. 46, no. 589/88), same type, (a synonym of C. cinereus Hwass, 1792).



clavus Linnaeus, 1758.

Published in Syst. Nat. 10th ed. (1): p. 716, no. 272, not figured.

Type series whereabouts unknown, (? mm).

Type locality not known.

Nomenclatural status, an available name, a totally suppressed name, on Official Index of Rejected and Invalid Specific Names in Zoology, ICZN opinion 753, (1965), see Section 1 B (1). Taxonomic status, none.

clenchi Martins, 1943.

Published in Bolm. Mus. nac. Rio de J. (12): p. 2 & fig. Holotype in MNRJ, $(36 \times 19.5 \text{ mm. orig. desc. } 38 \times 20 \text{ mm})$.

Type locality Barra do Furado, Municipio de Campos, State of Rio de Janeiro, Brazil, (at the shore line).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. lemniscatus Reeve, 1849.





clerii Reeve, 1844.

Published in Conch. Icon. 1 (Conus): pl. 43, sp. 229, (published Jan), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 175, not figured, published Jun).

Holotype in NHMUK, (31.9 x 15.7 mm). Type locality Cape St. Thomas, (State of Rio de Janeiro), Brazil, (in sandy mud, 35 fths).

Nomenclatural status, an available name.





<u>clodianus</u> "Ch" (variety mediterraneus Hwass) Nardo, 1847. Published in Sinon. Moderna, pt. 2: p. 41/42, sp. 12, fig. 521. Type series was in collection S. Chiereghin, present whereabouts unknown, (fig. ? mm).

Type locality beach of Chioggia, Gulf of Venice, (in beach sediment).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.




Figure in Chiereghin MSS.

cloveri Walls, 1978.

Published in Pariah (2): p. 2, fig. on p. 5, (published 30.01.78), (1979, Cone Shells: p. 338, figs on p. 224). Holotype in DMNH, (25.7 x 13 mm).

Type locality near Anse Bernard, Dakar Harbour, Senegal. Nomenclatural status, an available name. Taxonomic status, a valid species.





(Cylinder) clytospira Melvill & Standen, 1899.

Published in Ann. Mag. nat. Hist. 7 (4): p. 461, not figured, (1901, Proc. zool. Soc. Lond. unnumbered (2), pt. 1: p. 430, pl. 21, fig. 12).

Lectotype (C, M & W) in NHMUK, (108 x 33 mm).

Type locality Arabian Sea, about 125 miles W. S-W. of Bombay, (45 fths).

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. milneedwardsii* Jousseaume, 1894.





cocceus Reeve, 1844 ("1843").

Published in Conch. Icon. 1 (*Conus*): pl. 42, sp. 228, (published Jan), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 174, not figured, published Jun).

Lectotype (C, M & W) in NHMUK, $(31.2 \times 16.2 \text{ mm})$.

Type locality New Holland, (Australia), restricted (C, M & W) to Geographe Bay, West Australia.

Nomenclatural status, an available name.






coccineus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3390, no. 46, not figured.


Lectotype representative (Kohn) figure in Knorr, 1771, pl. 24, fig. 2, (fig. 31 x 15 mm). As this figure is ambiguous and could represent a specimen of C. granulatus Linnaeus, 1758 a neotype [possibly invalid ?] was designated (C, M & W) in NBC-ZMA, (49.4 x 24.3 mm),

Type locality not known, neotype from Samar Island, Philippines. Nomenclatural status, an available name. Taxonomic status, a valid species.











coelebs Hinds, 1843.

Published in Ann. Mag. nat. Hist. 11 (70): p. 256, not figured. Holotype was in collection Sir E. Belcher, present whereabouts unknown, (25.5 x ? mm).

Type locality Ambow, Fiji Islands, (on coral reefs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (juvenile) of C. terebra Born, 1778.

coelinae Crosse, 1858, (spelt caelinae p. 126).

Published in Revue Mag. Zoo. 2 (10): p. 81, p. 117 & p. 126, pl. 2, fig. 1.

Holotype in IRSN, (116 x 57 mm).

Type locality New Caledonia, corrected, (Delsaerdt) Hawaii Islands, (erroneous).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





coerulescens Schröter, 1803.

Published in Arch. Zool. Zoot. 3 (2): p. 67, not figured.

Holotype in MNG, (52 x 29 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. ermineus* Born, 1778.





coerulescens "Chemnitz" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 368, no. 27, not figured.

Lectotype representative (Walls) figures in Chemnitz, 1795, pl. 182, figs 1762 & 1763, (figs 41 x 25 mm, orig. desc. app. 44 x 25 mm).

Type locality St. Thomas Island, (Leeward Islands), Caribbean. Nomenclatural status, an available name, an invalid name, a homonym of *C. coerulescens* Schröter, 1803 and of *C. caerulescens* Lamarck, 1810.

Taxonomic status, a synonym of C. ermineus Born, 1778.



coffea Weinkauff, 1874.

Published in Jb. dt. malakozool. Ges. 1: p. 260.

Holotype of C. coffeae Gmelin, 1791 in NBC-ZMA, (27.3 x 16.1 mm). Type locality of C. coffeae not known.

Nomenclatural status, an available name, an invalid name, an unjustified emendation for *C. coffeae* Gmelin, 1791 and a homonym of *C. coffeae* Gmelin, 1791.

Taxonomic status, a synonym of C. coffeae Gmelin, 1791.





coffeae Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3388, no. 31, not figured.

Holotype in NBC-ZMA, $(27.3 \times 16.1 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





collisus Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 8, sp. 273 (323). Lectotype representative (Walls) figure in C. I. pl. 8, sp. 273, (fig. 41 x 20 mm), (three paralectotypes [dubious] in NHMUK, 43.5 x 20.5, 40.0 x 19.5 & 38.5 x 18.5 mm).

Type locality not known, designated (C, M & W) Phuket, Thailand. Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium).







Lectotype representative.

Paralectotype 1 in NHMUK. ©NHMUK.

colmani Röckel & Korn, 1990.

Published in Acta Conch. (2): p. 16, pl. 8, figs 1 - 4, pl. 9, figs 1 - 7, pl. 10, row 5, figs 3 - 5.

Holotype in AMS, $(42 \times 23.6 \text{ mm})$.

Type locality S.E. of Swains Reef, Great Barrier Reef,

Queensland, (Australia), (185 - 225 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species, possibly a semi fossil.





©AMS.

Dauciconus colombi Monnier & Limpalaër, 2012.

Published in Visaya vol. 3, no. 5: p. 15 & plates.

Holotype in MNHN, $(19.31 \times 10.67 \text{ mm})$.

Type locality La Vauclin Bay, East coast of Martinique Island, (in shallow water).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





colorovariegatus Kosuge, 1981.

Published in Bull. Inst. malac. Tokyo, 1 (6): p. 94, pl. 32, figs 1 - 5.

Holotype in IMT, $(63.3 \times 27.4 \text{ mm})$.

Type locality Bohol Island, Philippines.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. neptunus* Reeve, 1843.





colubrinus Lamarck, 1810.


Published in Annls. Mus. Hist. nat. Paris, 15: p. 433, no. 163, not figured.

Lectotype (Kohn) in MNHN, $(53.5 \times 24.5 \text{ mm})$.

Type locality "Mers des Grandes Indes", (Indian Ocean), restricted (Lauer) to northern coasts of Mozambique.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. praelatus* Hwass, 1792, (a synonym of *C. pennaceus* Born, 1778).





columba Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 709, no. 101, (1798, Tab. Enc. pl. 334, fig. 3).

Type figure designated, (Clench) figure in Gualtieri, 1742, pl. 25, fig. G, (fig. $29 \times 19 \text{ mm}$).

Type locality Indian Ocean, (erroneous), corrected (Vink) Fort de France, Martinique, (Windward Islands).

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. puncticulatus* Hwass, 1792, (p. 702, no. 92).



comatosa Pilsbry, 1904.

Published in Proc. Acad. nat. Sci. Philad. 56 (2): p. 550. Lectotype of *C. dormitor* Pilsbry, 1904 in ANSP, (44 x 16 mm). Type locality of *C. dormitor* Kikai, Osumi, South Japan, (in a deposit, probably Pliocene).

Nomenclatural status, an available name, a new replacement name (nomen novum) for C. dormitor Pilsbry, 1904 (fossil). Taxonomic status, a valid species, although the holotype of C. dormitor Pilsbry, 1904, (43.5 x 16.5 mm), is a fossil, living specimens exist.





Lectotype of C. dormitor.

(Leptoconus) commodus A. Adams in H. & A. Adams, 1853. Published in The Genera of Recent Mollusca 1: p. 251, not figured, (published Dec), (1854, Proc. zool. Soc. Lond. pt. 21, no. 255: p. 117, no. 9, (as Conus), not figured, published Nov). Holotype was in collection Grüner, bought by Landauer, then in collection Von Maltzan and sold again to a dealer, present whereabouts unknown, (figure in Weinkauff, 1875, p. 330, pl. 61, figs 4 & 6, is the same shell), (fig. 31 x 14 mm). Type locality not known.

Nomenclatural status, an available name. Taxonomic status, doubtful (nomen dubium), possibly a synonym (subadult) of *C. eximius* Reeve, 1849.



Figure in Weinkauff, 1875.

(Textilia) communis Swainson, 1840.

Published in Treatise on Malacology pt. 2: p. 312, not figured. Lectotype representative (C, M & W) figure in Tab. Enc. 1798, pl. 346, fig. 1. (fig. 68 x 36 mm), (a syntype fig. 2 in Tab. Enc. is in MHNG, $68 \times 35 \text{ mm}$).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of C. archiepiscopus Hwass, 1792, same type, (a synonym (form) of C. textile Linnaeus, 1758), the syntype in MHNG is a specimen of C. textile Linnaeus, 1758).





Figure in Tab. Enc.

Syntype in MHNG.

compactus
(subspecies imperialis Linnaeus) Wils, 1970.
Published in Family Conidae 8: p. 12, no. 15, pl. 2, fig. 7.
Lectotype (C, M & W) in NBC-ZMA, (71 x 43.1 mm).
Type locality Nosy Be, Madagascar.
Nomenclatural status, an available name.
Taxonomic status, a synonym (form) of C. imperialis Linnaeus, 1758.





complanatus Sowerby II, 1866.

Published in Thes. Conch. 3 (Conus), pt. 24 - 25: App. p. 330, no. 441, pl. 28 (289), figs 650 (variety) & 651, (1857, pl. 23, fig. 576 variety as C. victoriae Reeve, 1843).

Two probable syntypes in NHMUK, (fig. 576, 47.9 x 26.9 mm. & fig. 650, variety, $52.3 \times 29.6 \text{ mm}$).


Type locality Australia.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of C. victoriae Reeve, 1843, as Sowerby II first suspected in 1857.









Probable syntype no. 1 in NHMUK.

Probable syntype no. 2 in NHMUK.

compressus Sowerby II, 1866.

Published in Thes. Conch. 3 (Conus), pt. 24 - 25: App. p. 325, no. 404, pl. 25 (286), figs 602 & 603.

Holotype in NHMUK, $(24.8 \times 10.6 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





comptus Gould, 1853.

Published in Boston J. nat. Hist. 6 (3): p. 387, pl. 14, fig. 23. Holotype was in collection E. Jewett, present whereabouts unknown, $(31 \times 19 \text{ mm})$.

Type locality Santa Barbara, California, [dubious].

Nomenclatural status, an available name.

Taxonomic status, a synonym (juvenile) of *C. purpurascens* Sowerby I & II, 1833, recognised by Gould in 1862.



Figure in Gould.

comptus A. Adams, 1854 ("1853").

Published in Proc. zool. Soc. Lond. pt. 21, no. 255: p. 119, no. 19, not figured.

Holotype in NHMUK, $(20.9 \times 12 \text{ mm})$.

Type locality Natal, (South Africa), (Cuming), [erroneous]. Nomenclatural status, an available name, an invalid name, a homonym of *C. comptus* Gould, 1853.

Taxonomic status, a synonym of *C. papilliferus* Sowerby I & II, 1834.






"<u>conakryensis</u>" (forma *pulcher* Lightfoot), van Rossum, 1997. Published in World Shells (20): p. 41, figs 10 & 11.

Specimen described whereabouts unknown, (figs 62 x 32 mm. orig. desc. up to 51 mm).

Locality off the coast of Guinea (West Africa), (depth to 60 mtrs).

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name, stated to be infrasubspecific after 1960.

Taxonomic status, none, (a specimen of *C. pulcher* Lightfoot, 1786).



Original illustration.

concatenatus Kiener, 1845 (1850).

Published in Coq. Viv. 2: pl. 110, fig. 1, (1850, Coq. Viv. 2: Sup. p. 362, no. 324).

Holotype was in collection Lorois, present whereabouts unknown, $(35 \times ? \text{mm})$, lectotype representative (Lauer) figure in C. V. pl. 110, fig. 1, (fig. 34 x 18 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. textile Linnaeus, 1758.



concinnulus Crosse, 1858.

Published in Revue Mag. Zool. 2 (10): p. 200, no. 2.

Lectotype of C. concinnus Broderip, 1833 in NHMUK, (20.3 x 10.7 mm).

Type locality of C. concinnus Gulf of California, (sand). Nomenclatural status, an available name, a new replacement name,

(nomen novum) for *C. concinnus* Broderip, 1833. Taxonomic status, an invalid name, not in Family Conidae, in Family Columbellidae.





Lectotype of C. concinnus in NHMUK.

concinnus Broderip, 1833.

Published in Proc. zool. Soc. Lond. pt. 1, no. 4: p. 53, not figured, (figure in C. I. pl. 32. no. 32).

Lectotype (Kohn) in NHMUK, (20.3 x 10.7 mm).

Type locality Gulf of California, (sand).

Nomenclatural status, an available name, an invalid name, a homonym of *C. concinnus* J. de C. Sowerby, 1821 (fossil), renamed *C. concinnulus* Crosse, 1858.

Taxonomic status, an invalid name, not in Family Conidae, in Family Columbellidae.





concinnus Sowerby II, 1866.

Published in Thes. Conch. 3 (*Conus*), pt. 24 - 25: App. p. 329, no. 438, pl. 28 (289), fig. 646.

Holotype was in collection T.L. Taylor, present whereabouts unknown, (fig. $25 \times 14 \text{ mm}$).

Type locality not known.

Nomenclatural status, an available name, an invalid name, a homonym of *C. concinnus* J. de C. Sowerby, 1821 (fossil) and of Broderip, 1833, renamed *C. sapphirostoma* Weinkauff, 1874. Taxonomic status, a synonym of *C. biliosus* (Röding), 1798



Figure in Thes. Conch. 3.

concolor Sowerby I & II, 1841.

Published in Conch. Ill. (Conus): Index, p. 3, no. 59. Lectotype of C. unicolor Sowerby I & II, 1834 in NHMUK, (45.5 x 23.0 mm.)

Type locality of $C.\ unicolor$ not known, (designated Röckel & Korn) Solomon Islands.

Nomenclatural status, an available name, a new replacement name (nomen novum) for *C. unicolor* Sowerby I & II, 1834.

Taxonomic status, a subspecies of C. hyaena Hwass, 1792.





Lectotype of C. unicolor in NHMUK.

concolor
1933.
(subspecies pigmentatus Adams & Reeve) Barros E Cunha,

Published in Mems Estud. Mus. zool. Univ. Coimbra, 1 (71): p. 52, no. 43, not figured.

Holotype in MZUC, $(22.9 \times 15.1 \text{ mm})$.

Type locality eastern seas, exact locality unknown.

Nomenclatural status, an available name, an invalid name, a homonym of *C. concolor* Sowerby I & II, 1841.

Taxonomic status, a synonym of *C. pigmentatus* Adams & Reeve, 1848, (a synonym of *C. balteatus* Sowerby I & II, 1833).





condensus Sowerby II, 1866.

Published in Thes. Conch. 3 (*Conus*), pt. 24 - 25: App. p. 326, no. 417, pl. 26 (287), fig. 622.

Holotype whereabouts unknown, lectotype representative (Lauer) figure in T. C. pl. 26, fig. 622, (fig. 46 x 21 mm).

Type locality Sandwich Islands, (Hawaii), [dubious].

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. canonicus Hwass, 1792.



<u>condoriana</u> (variety *minimus* Linnaeus) Crosse & P. Fischer, 1864. Published in J. Conch. Paris, 12: p. 334, not figured.

Holotype was probably in collection A. Michau, present whereabouts unknown, $(21 \times 12 \text{ mm})$.

Type locality Condor Island, off Saigon, South China Sea, (in holes formed by polyps), (Michau).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. coronatus Gmelin, 1791.

Darioconus pennaceus f "confusa" Bozzetti, 2010.

Published in Malacologia no.69, p.10.

Specimen described whereabouts unknown, (dimensions not given). Locality Southwest Madagascar,

Nomenclatural status, an unavailable name, cited as a form after 1961.

Taxonomic status, none, (Specimens are C. pennaceus Born, 1778).



Figure in Bozzetti.

Leptoconus (Thoraconus) <u>congruens</u> (subspecies biraghii Raybaudi) Korn & G. Raybaudi (Massilia), 1993.

Published in Conchiglia 25 (268): p. 33 & figs 1 - 1c, 3, 6, & pl. 1, fig. 3.

Holotype in SNMS, $(10.9 \times 5.8 \text{ mm})$.

Type locality off northern Somalia, Gulf of Aden, (dredged). Nomenclatural status, an available name.

Taxonomic status, a subspecies of C. biraghii Raybaudi, 1992.





conicus Galindo, 1977.

Published in Index & Register of Seashells: p. 116, not figured. Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none.

<u>connectens</u> A. Adams, 1855 ("1854").

Published in Proc. zool. Soc. Lond. pt. 22, no. 268: p. 136, pl. 28.

Holotype in NHMUK, $(54 \times 29 \text{ mm})$.

Type locality China, (Cuming), (erroneous).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. daucus* Hwass, 1792.





consanguineus E.A. Smith, 1880.

Published in Proc. zool. Soc. Lond. unnumbered (48), pt. 3: p. 478, pl. 48, fig. 1.

Holotype in NHMUK, $(87.3 \times 48.8 \text{ mm. orig. desc. } 88 \times 49 \text{ mm})$. Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. fergusoni Sowerby II, 1873.





consiliarius "S" Lightfoot, 1786.

Published in Cat. Portland Mus: p. 153, no. 3352, not figured. Specimen(s) named sold at auction, present whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

consors Sowerby I & II, 1833.

Published in Conch. Ill. (Conus): pl. 36, fig. 42.

Lectotype representative (Kohn) figure in C. I. pl. 36, fig. 42, $(fig. 63 \times 34 \text{ mm})$.

Type locality not mentioned, designated (C, M & W) Singapore. Nomenclatural status, an available name. Taxonomic status, a valid species.



conspersus Reeve, 1844 ("1843").

Published in Conch. Icon. 1 (*Conus*): pl. 47, sp. 262, (published Feb), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 180, not figured, published Jun).

Holotype was in collection Grüner, probably in MGB (defunct), present whereabouts unknown, (fig. $30 \times 16 \text{ mm}$).

Type locality not known, designated (C, M & W) the Moluccas, Indonesia.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. spectrum* Linnaeus, 1758.



Figure in Conch. Icon.

conspicuus "S" Lightfoot, 1786.

Published in Cat. Portland Mus: p. 167, no. 3623, not figured. Specimen(s) named sold at auction, present whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

consul Boivin, 1864.

Published in J. Conch. Paris, 12: p. 33, pl. 1, figs 5 & 6. Holotype was in collection A. Boivin, in MHNH, destroyed in World War 2, (43 x 19 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. magus Linnaeus, 1758.



Figures in Boivin.


Dendroconus betulinus rufoluteus f. "continua" Bozzetti, 2010. Published in Malacologia no.68, p.12 & figs.

Specimens described whereabouts unknown, (no dimensions given). Locality Tulear Madagascar.

Nomenclatural status, an unavailable name, described as a form after 1961.

Taxonomical status, none, (specimens are C. betulinus Linnaeus, 1758).





Specimens illustrated in Malacologia.

contusus Reeve, 1848.

Published in Conch. Icon. 1 (*Conus*): Sup. pl. 2, sp. 276. Holotype in NHMUK, (32.6 x 15.2 mm).

Type locality Moluccas, Indonesia.

Nomenclatural status, an available name

Taxonomic status, a synonym of C. cinerarius (Röding), 1798, (a synonym (form) of C. monachus Linnaeus, 1758).





convolutus Sowerby II, 1858.

Published in Thes. Conch. 3 (*Conus*), pt. 18: p. 44, no. 380, pl. 23 (209), fig. 564.

Lectotype (C, M & W) in NHMUK, $(59.3 \times 24 \text{ mm})$.

Type locality not mentioned, designated (Lauer) Madagascar.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. omaria Hwass, 1792.





cooki Brazier, 1870.

Published in Proc. zool. Soc. Lond. unnumbered (38), pt. 1: p. 109, not figured, (published Feb), (J. Conch. Paris, 18: p. 300, not figured, in French, published Jul).

Lectotype (C, M & W) in AMS, (17.2 x 11.1 mm, orig. desc. 21.1 x 11.6 mm).

Type locality Cooks Landing, Botany Bay, N.S.W., Australia, (rocks).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. aplustre Reeve, 1843.



"coralinus" Habe & Kosuge, 1970.

Published in Pacific Shell News (1): p. 1, fig. frontispiece. (published Jan)

Specimen named in NSMT, $(38.4 \times 21.5 \text{ mm})$.

Locality Zamboanga, Mindanao, Philippines (erroneous).

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name.

Taxonomic status, none, (a specimen of C. coralinus Habe & Kosuge, 1970.





Specimen named in NSMT.

(Rhizoconus) coralinus Habe & Kosuge, 1970.

Published in Venus, Kyoto, 29 (3): p. 81, pl.5, figs 3 & 4, (published Aug).

Holotype in NSMT, $(38.4 \times 21.5 \text{ mm})$.

Type locality Zamboanga, Mindanao, Philippines, (erroneous). Nomenclatural status, an available name, an invalid name, a homonym of *C. corallinus* Kiener, 1845 (1847).

Taxonomic status, a synonym of C. klemae Cotton, 1953.





corallinus "S" Lightfoot, 1786.

Published in Cat. Portland Mus: p. 149, no. 3280, not figured. Specimen(s) named sold at auction, present whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

corallinus "Solander's MSS" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 404, no. 94, not figured.

Specimen named whereabouts unknown, cited species in Mus.

Callone. p. 11, no. 173, not figured.

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. coccineus* Gmelin, 1791.

Taxonomic status, none, (a specimen of *C. coccineus* Gmelin, 1791).

corallinus Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 73, fig. 2, (1849) (1850), Coq. Viv. 2: p. 246, no. 214).

Holotype was in collection Dupont, present whereabouts unknown, (24 \times ? mm), (fig. 27 \times 11 mm).

Type locality not mentioned, designated (C, M & W) Mactan Island, (Cebu), Philippines.

Nomenclatural status, an available name.

Taxonomic status, a valid species.



Figure in Coq. Viv. 2.

coralloides Perry, 1811.

Published in Conchology: pl. 25, fig. 6.

Holotype representative (C, M & W) figure in Perry, pl. 25, fig. 6, (fig. $49 \times 27 \text{ mm}$).

Type locality Sumatra and Ceylon (Sri Lanka).

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost holotype, a limited description and an unrecognizable figure, possibly a synonym (form) of *C. argillaceus* Perry, 1811, or possibly a synonym of *C. monile* Hwass, 1792, if the type locality is correct then the latter is most likely.



corbieri (subspecies pennaceus Born) Blöcher, 1994.

Published in World Shells (10): p. 50, figs 5 - 7.

Holotype in SMNS, $(54.6 \times 29.4 \text{ mm})$.

Type locality Salary, southwestern Madagascar, (in lagoon in shallow water).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. pennaceus* Born, 1778.





corbula Sowerby II, 1858.

Published in Thes. Conch. 3 (*Conus*), pt. 18: p. 42, no. 365, pl. 23 (209), fig. 573.

Holotype in NHMUK, $(69.5 \times 34.5 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. textile Linnaeus, 1758.





cordigera Sowerby II, 1866.

Published in Thes. Conch. 3 (*Conus*), pt. 24 - 25: App. p. 329, no. 437, pl. 21 (207), fig. 498, (as C. *nobilis*).

Holotype in NHMUK, $(61.9 \times 31.6 \text{ mm})$.

Type locality Cuyo Island, Philippines.

Nomenclatural status, an available name.

Taxonomic status, a valid species or possibly a subspecies of *C. nobilis* Linnaeus, 1758.





<u>coriolisi</u> (subspecies *orbignyi* Audouin) Moolenbeek & Richard in Röckel, Richard & Moolenbeek, 1995.

Published in Mem. Mus. natn. Hist. nat. Paris, no. 167, 14 (9): p. 578, figs 12 & 13.

Holotype in MNHN, $(44.6 \times 16 \text{ mm})$.

Type locality Banc Capel, Coral Sea, (off New Caledonia), (240 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. orbignyi* Audouin, 1831.





coromandelicus E.A. Smith, 1894.

Published in Ann. Mag. nat. Hist. 6 (14): p. 159, pl. 4, figs 1 & 2.

Holotype in ZSIC, $(37 \times 14 \text{ mm})$.

Type locality Coromandel coast, Bay of Bengal, (80 - 110 fths). Nomenclatural status, an available name.

Taxonomic status, a valid species, probably not in genus *Conus*, but in genus *Conorbis*.



coronacivica (Röding), 1798.

Published in Mus. Bolten 2: p. 38, no. 466/3, not figured. Lectotype representative (Kohn) figure in Martini, 1773, pl. 61, fig. 678. (fig. 47 x 27 mm).

Type locality not mentioned, (Martini indicated West Indies). Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. regius* Gmelin, 1791.



coronaducalis (Röding), 1798.

Published in Mus. Bolten 2: p. 38, no. 464/2*, not figured. Lectotype representative (Kohn) figure in Martini, 1773, pl. 62, fig. 693. (fig. 51 x 27 mm).

Type locality not mentioned, (Martini indicated Mauritius). Nomenclatural status, an available name.

Taxonomic status, a synonym of C. fuscatus Born, 1778, (a synonym (form) of C. imperialis Linnaeus, 1758).



coronalis (Röding), 1798.

Published in Mus. Bolten 2: p. 38, no. 478/9, not figured. Holotype whereabouts unknown, figure cited in Gmelin, 1791, p. 3389, no. 39, (fig. 38 x 19 mm), neotype (Kohn by inference) in NHMUK, (27.5 x 16.4 mm).

Type locality not mentioned, neotype from Australia.

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of *C. coronatus* Gmelin, 1791, p. 3389, same type.





Neotype of C. coronatus in NHMUK.

coronatus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3389, no. 39, not figured.

Neotype (Kohn) in NHMUK, $(27.5 \times 16.4 \text{ mm})$.

Type locality not known, neotype from Australia.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





<u>coronatus</u> (variety <u>ammiralis</u> Linnaeus) Gmelin, 1791. Published in Syst. Nat. 13th ed. 1 (6): p. 3379, no number, not figured.

Lectotype representative (Walls) figure in Martini, 1773, vignette 26, fig. 1. (fig. $38 \times 20 \text{ mm}$). Type locality not known.

Nomenclatural status, an available name, an invalid name, a homonym of *C. coronatus* Gmelin, 1791, (p. 3389, no. 39). Taxonomic status, an objective junior synonym of *C. architalassus* Lightfoot, 1786, same type, (a synonym of *C. ammiralis* Linnaeus, 1758).



coronatus Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 7, sp. 263 (313). Lectotype (G. Raybaudi) in NHMUK, (9 x 4.6 mm). Type locality Ticao Island, Philippines, (on the reef). Nomenclatural status, an available name, an invalid name, a homonym of C. coronatus Gmelin, 1791, (p. 3389 & p. 3379), renamed C. papalis Weinkauff, 1875. Taxonomic status, a synonym of C. papalis Weinkauff, 1875.



"coronensis" "Röckel & Korn" Anonymous, 1992.

Published in World Shells (3): p. 89, no. 136, not figured.

No specimen designated, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name and no author, (Röckel & Korn did not introduce this name).

Taxonomic status, none, (stated to be a specimen of *C. telatus* [presumably Reeve, 1848]).

corrugatus Sowerby II, 1870.

Published in Proc. zool. Soc. Lond. unnumbered (38), pt. 2: p. 257, pl. 22, fig. 7.

Holotype in NHMUK, (19 \times 8.8 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. jaspideus Gmelin, 1791.





coruscus "S" in Lightfoot, $1\overline{786}$.

Published in Cat. Portland Mus: p. 153, no. 3352, not figured. Specimen(s) named sold at auction, present whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

costatus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3388, no. 35, not figured.

Holotype was in collection N. Gualtieri, in MSNP but no longer identifiable, (fig. $49 \times 31 \text{ mm}$).

Type locality not known.

Nomenclatural status, an available name.


Taxonomic status, an invalid name, not in Family Conidae, in Family Strombidae.

costatus Holten, 1802.

Published in Enum. Syst. Conch: p. 38, no. 483, not figured. Lectotype representative (Kohn) figure in Chemnitz, 1795, pl. 181, fig. 1746, (fig. 65 x 34 mm).

Type locality not mentioned, (Chemnitz indicated China). Nomenclatural status, an available name, an invalid name, a homonym of *C. costatus* Gmelin, 1791.

Taxonomic status, a synonym of C. sulcatus Hwass, 1792.




costatus "Chemnitz" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 398, no. 80, not figured.

Lectotype representative (Kohn) figure in Chemnitz, 1795, pl. 181, fig. 1746, (fig. 65 x 34 mm, orig. desc. app. 62 x 33 mm). Type locality China.

Nomenclatural status, an available name, an invalid name, a homonym of C. costatus Gmelin, 1791 and of Holten, 1802. Taxonomic status, an objective junior synonym of C. costatus Holten, 1802 and of C. asper Lamarck, 1810, same type, (a synonym of C. sulcatus Hwass, 1792).



couderti Bernardi, 1860.

Published in J. Conch. Paris, 8: p. 212, pl. 4, figs 3 & 4, (1861, Monographie de Genre Conus: p. 11, pl. 2, fig. 17).

Holotype was in collection Coudert (of Bordeaux, France), present whereabouts unknown, $(25 \times 14 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. erythraeensis* Reeve, 1843.



Figures 3 & 4 in Journ. Conch.



Figure in Monographie.

coxeni Brazier, 1875.

Published in Proc. zool. Soc. Lond. unnumbered (43), pt. 1: p. 34, pl. 4, fig. 10.

Holotype was in collection C. Coxen of Brisbane, Australia, present whereabouts unknown, (fig. 32 x 15 mm. orig. desc. 31.7 x 14.8 mm).

Type locality Moreton Bay, (Queensland, Australia).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. cyanostoma* Adams, 1853.



Figure in Brazier.

coxianus Sowerby III, 1895.

Published in Proc. malac. Soc. Lond. 1 (5): p. 215, pl. 13, fig. 7.

Holotype in NHMUK, $(39.3 \times 21 \text{ mm})$.

Type locality Zeyla Bay, Somaliland, (Somalia).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. locumtenens* Blumenbach, 1791.





<u>crassus</u> Sowerby II, 1858.

Published in Thes. Conch. 3 (*Conus*), pt. 18: p. 25, no. 203, pl. 12 (198), figs 254 & 255.

Lectotype (Walls) in NHMUK, (44.2 x 28.8 mm).

Type locality Fiji Islands, (McGillivray).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. eburneus* Hwass, 1792 or possibly a valid species.





<u>crebremaculata</u> (variety ammiralis Linnaeus) Dautzenberg, 1937. Published in Mem. Mus. r. Hist. nat. Belg. 2 (18): p. 19, not figured.

Lectotype representative (Walls) figure in Chemnitz, 1788, pl. 141, fig. 1309, (fig. 65 \times 37 mm).

Type locality not mentioned, (Chemnitz indicated East Indian Ocean).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. ammiralis Linnaeus, 1758.



<u>crebrisulcus</u> Sowerby II, 1857, (spelt *crebrisulcatus* on pl. 14). Published in Thes. Conch. 3 (*Conus*), pt. 17: p. 21, no. 173, pl. 14 (200), fig. 321.

Holotype in NHMUK, $(12.2 \times 6 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (juvenile) of *C. jaspideus* Gmelin, 1791.





crenulatus (Röding), 1798.

Published in Mus. Bolten 2: p. 46, no. 581/80, not figured.

Two specimens named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

crenulatus Kiener, 1845 (1850).

Published in Coq. Viv. 2: pl. 109, fig. 1, (1850, Coq. Viv. 2: Sup. p. 355, no. 318).

Holotype was in collection Lorois, present whereabouts unknown, $(32 \times ? \text{ mm})$, holotype representative (Kohn) figure in C. V. pl. 109, fig. 1, (fig. $32 \times 14 \text{ mm}$).

Type locality not mentioned, designated (C, M & W) coast of Louisiana, Gulf of Mexico.

Nomenclatural status, an available name, an invalid name, a homonym of *C. crenulatus* Deshayes, 1835 (fossil), renamed *C. armiger* Crosse, 1858.

Taxonomic status, a synonym of C. armiger Crosse, 1858.



Figure in Coq, Viv. 2.

crepusculum Reeve, 1844.

Published in Conch. Icon. 1 (*Conus*): pl. 45, sp. 251, (published Feb), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 178, not figured, published Jun).

Holotype was in collection J. Adamson, auctioned in 1848, present whereabouts unknown, (fig. $30 \times 17 \text{ mm}$).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. furvus Reeve, 1843.



Figure in Conch. Icon.

cretaceus Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 99, fig. 1, (1849, Coq. Viv. 2: p. 264, no. 231).

Holotype was in collection A.C. Bernardi, acquired by M. Gubba then in MHNH, destroyed in World War 2, $(32 \times ? mm)$, holotype representative (Kohn) figure in C. V. pl. 99, fig. 1, (fig. 32 x 16 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. mindanus Hwass, 1792.



<u>cretheus</u> "Ch" (variety mediterraneus Hwass) Nardo, 1847. Published in Sinon. Moderna, pt. 2: p. 41/42, sp. 5, figs 509 - 510.

Type series was in collection S. Chiereghin, present whereabouts unknown, (fig ? mm).

Type locality Gulf of Venice, (base of sponges).

Nomenclatural status, an available name.

Taxonomical status, a synonym (form) of *C. ventricosus* Gmelin, 1791.



Figure in Chiereghin MSS.

crioulus Tenorio & Afonso, 2004.

Published in Visaya vol. 1, no. 2: p. 30, pl. 5.

Holotype in MNCM, $(19.1 \times 10.7 \text{ mm})$.

Type locality Praia Real, North coast of Maio Island, Cape Verde Islands, (half buried in sand in crevices and rock platforms, in very shallow water, $0.25-0.5~\rm mtrs$).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





crocatus Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 424, no. 136, not figured.

Holotype in MHNG, $(44 \times 20 \text{ mm, orig. desc. } 50 \times ? \text{ mm})$. Type locality "Mers des Grandes Indes" (Indian Ocean).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





croceus Sowerby I & II, 1833.

Published in Conch. Ill. (Conus): pl. 29, fig. 27.

Lectotype representative (Kohn) figure in C. I. pl. 29, fig. 27, $(fig. 44 \times 25 mm).$

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. daucus* Hwass, 1792, recognised by Sowerby I & II in Index, 1841.



croceus E.A. Smith, 1877.

Published in Ann. Mag. nat. Hist. 4 (19): p. 223, not figured. Holotype in NHMUK, (25.3 \times 8.5 mm, orig. desc. 27 \times 9 mm).

Type locality not known.

Nomenclatural status, an available name, an invalid name, a homonym of *C. croceus* Sowerby I & II, 1833, renamed *C. hypochlorus* Tomlin, 1937.

Taxonomic status, a synonym of C. hypochlorus Tomlin, 1937





crosseanus Bernardi, 1861.

Published in J. Conch. Paris, 9: p. 168, pl. 6, figs 3 & 4, (Monographie de Genre Conus: p. 13, pl. 1, figs 2 - 6, spelt *C. crosscanus*).

Lectotype (C, M & W) in NHMUK, (66.8 x 39.4 mm, orig. desc. 66 x 38 mm).

Type locality New Caledonia.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. marmoreus* Linnaeus, 1758.





crotchii Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 6, sp. 254 (304). Holotype was in collection H. Cuming, present whereabouts unknown, (fig. 29 x 17 mm), (three specimens in NHMUK type collection are not part of type material).

Type locality Saldanha Bay, South Africa, (erroneous), corrected (C, M & W) to Santa Monica, Boavista, Cape Verde Islands.

Nomenclatural status, an available name.

Taxonomic status, a valid species.



Figure in Conch. Icon.

crucifer (Röding), 1798.

Published in Mus. Bolten 2: p. 48, no. 617/110, not figured. Lectotype representative (C, M & W) figure in Martini, 1773, pl. 55, fig. 605, (fig. 61 x 38 mm).

Type locality not mentioned, (Martini indicated Nicobar Islands), [erroneous].

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of *C. achatinus* (Röding), 1798, same type, (a synonym of *C. ermineus* Born, 1778).



<u>cruzensis</u> (subspecies <u>centurio</u> Born) Nowell-Usticke, 1968. Published in Caribbean Cones from St. Croix and the Lesser Antilles: p. 12, pl. 2, fig. 997.

Lectotype (Nowell-Usticke $\overline{1971}$), in AMNH, (38 x 24 mm).

Type locality Ham Bay, St. Croix, Virgin Islands, (Leeward Islands).

Nomenclatural status, an available name.

Taxonomic status, a synonym (subadult) of C. centurio Born, 1778.





cumingii Reeve, 1848.

Published in Conch. Icon. 1 (Conus): Sup. pl. 3, sp. 282.

Holotype in NHMUK, $(34.3 \times 17.2 \text{ mm})$.

Type locality Island of Mindanao, Philippines.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cumingii Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 8, sp. 277 (327). Lectotype (C, M & W) in NHMUK, (53.8 x 26.9 mm).

Type locality Salango, Ecuador, (South America), (about 7 fths). Nomenclatural status, an available name, an invalid name, a homonym of C. cumingii Reeve, 1848.

Taxonomic status, a synonym of C. virgatus Reeve, 1849.





(Pupuriconus) <u>cuna</u> Petuch, 1998.

Published in Conchiglia 30 (287): p. 30 & figs.

Holotype in FMNH, $(20 \times 10 \text{ mm})$.

Type locality Moro Tupo Isl., San Blas Islands, Panama (East coast), (in sand under coral rubble on coral reef, 3 mtrs). Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. mus Hwass, 1792.





cuneatus Sowerby II, 1873.

Published in Proc. zool. Soc. Lond. unnumbered (41), pt. 1: p. 146, pl. 15, fig. 5.

Holotype in NBC-ZMA, $(33 \times 20.2 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomical status, a synonym (form) of C. malacanus Hwass, 1792.





cuneiformis E.A. Smith, 1877.

Published in Q. Jl. Conch. 1: p. 202, & fig.

Lectotype (C, M & W) in NHMUK, $(25.2 \times 14.1 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. inscriptus* Reeve, 1843.





cuneolus Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 37, sp. 205a & b, (published Dec), (1844, Proc. zool. Soc. Lond. pt. 11, no. 130: p. 173, not figured, published Jun).

Lectotype (C, M & W) in NHMUK, $(33.1 \times 20.6 \text{ mm})$.

Type locality not known, designated (C, M & W) Sal Island, Cape Verde Islands.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





<u>curassaviensis</u> (variety *cedonulli* Linnaeus) Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 602, no. 1C, (1798, Tab. Enc. pl. 316, fig. 4).

Lectotype representative (C, M & W) figure in d'Argenville, 1757, Sup. pl. 1, fig. X, (fig. 50 x 24 mm).

Type locality Island of Curacao, (erroneous), corrected (Vink & von Cosel) Aruba, (Netherlands Antilles).

Nomenclatural status, an available name.

Taxonomic status, a valid species.



curralensis Rolán, 1986.

Published in Publcoes. Occ. Soc. port. malac. (6): p. 10, fig. 1b.

Holotype in MNCM, $(20.7 \times 12.3 \text{ mm})$.

Type locality Curral, Santa Luzia Island, Cape Verde Islands, (stones & rubble, 1 - 3 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cutisanguina (Röding), 1798.

Published in Mus. Bolten 2: p. 48, no. 619/111, not figured. Lectotype representative (C, M & W) figure in Martini, 1773, pl. 55, fig. 605, (fig. 61 x 38 mm).

Type locality not mentioned, (Martini indicated Nicobar Islands), [erroneous],

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of C. achatinus (Röding), 1798 and of C. crucifer (Röding), 1798, same type, (a synonym of C. ermineus Born, 1778).



cuvieri Crosse, 1858.

Published in Revue Mag. Zool. 2 (10): p. 123.

Lectotype of C. cervus Sowerby I & II, 1838 and C. deshayesii Reeve, 1843 in IRSN, (51.2 x 27.4 mm).

Type locality of C. cervus and C. deshayesii Swan River, Australia (erroneous corrected to Djibouti, (Somalia).

Nomenclatural status, an available name, a new replacement name (nomen novum) for *C. deshayesii* Reeve, 1843.

Taxonomic status, a valid species.





Lectotype of ${\it C. cervus}$ in IRSN.

vidua cuyoensis Lorenz & Barbier, 2012.
Published in Acta Conch. 11: p. 3, pls 1 & 2.
Holotype in MNHN, no.24815, (29.30 x ? mm).
Type Locality Cuyo Islands, Philippines, (in shallow water).
Nomenclatural status, an available name.
Taxonomic status, a synonym (colour form) of C. vidua Reeve,
1843, a synonym of C. bandanus Hwass, 1792.





(Chelyconus) cyanostoma A. Adams in H. & A. Adams, 1853.

Published in The Genera of Recent Mollusca 1: p. 253, not figured, (1854, Proc. zool. Soc. Lond. pt. 21, no. 255: p. 116, no. 3, (as Conus), not figured).

Holotype in NHMUK, (27 x 15.2 mm).

Type locality West Africa, (Cuming), (erroneous), corrected (C, M

Type locality West Africa, (Cuming), (erroneous), corrected (C, M & W) Keppel Bay, Queensland, Australia.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





cyanosus
(variety textile euetrios Sowerby) Lauer, 1987.
Published in Rossiniana (35): p. 17, no. 13, not figured,
(Rossiniana (36): p. 12, fig. 13c).
Specimens described in collection Antoine, (34.5 x 16 & 36 x 18 mm).

Locality La Reunion.

Nomenclatural status, an unavailable name (nomen nudum), described as a variety after 1960.

Taxonomic status, none, (a specimen of *C. textile* Linnaeus, 1758).





Figures in Lauer.

<u>cylindraceus</u> Broderip & Sowerby I, 1830.
Published in Zool. J. Lond. 5: Sup. p. 51, pl. 40, fig. 5.
Holotype representative (Kohn) figure in Broderip & Sowerby I, pl. 40, fig. 5, (fig. 36 x 12 mm).
Type locality not known, designated (C, M & W) New Caledonia.
Nomenclatural status, an available name.
Taxonomic status, a valid species.

