

Section F fabula to futunaensis

Copyright © 2011.R Filmer Conditions of use.

The content of this website is provided for personal and scientific use and may be downloaded for this purpose. It may not be used in whole or part for any commercial activity and publication of any of the content on the internet is limited to the Cone Collector website (www.TheConeCollector.com). Authors wishing to publish any of the pictures may do so on a limited basis but should inform M Filmer so that the original owner of the rights to the picture can be acknowledged. (mike@mfilmer.fsnet.co.uk)

Version 1.0 October 2011.

fabula Sowerby I & II, 1833.

Published in Conch. Ill. (Conus): pl. 24, figs 5 & 5*.

Lectotype representative (Kohn) figure in C. I. pl. 24, fig. 5, $(\text{fig. }25 \times 13 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. scabriusculus* Dillwyn, 1817, recognised by Sowerby II in 1858, (a synonym of *C. coffeae* Gmelin, 1791).





fantasmalis Rolán, 1990.

Published in Iberus, Sup. 2: p. 37, pl. 1, fig. 8, pl. 2, fig. 8 & pl. 5, fig. N.

Holotype in MNCM, $(29 \times 16.5 \text{ mm})$.

Type locality Navio Quebrado Bay, Maio Island, Cape Verde Islands, (rocks).

Nomenclatural status, an available name.






fasciata (variety virgo Linnaeus) Menke, 1828.

Published in Synopsis Methodica Molluscorum Generum Omnium et Specierum Earum etc, Pyrmont, 1st ed: p. 43, not figured. Type series whereabouts unknown, cited figure in Martini, 1773, pl. 53, fig. 589, (fig. 38 x 20 mm).

Type locality not known.

Nomenclatural status, an available name, an invalid name, a homonym of *C. fasciatus* Schröter, 1803 and of Perry, 1811. Taxonomic status, a synonym of *C. lividus* Hwass, 1792 as suspected by Menke, (2nd ed), figure cited is the same as that for *C. detritus* Menke, 1830.



Cited figure in Martini.

fasciata (variety mediterraneus Hwass) Requien, 1848.

Published in Catalogue des Coquilles de l'Ile Corse (Corsica): p. 86, no. 614, not figured.

Type series whereabouts unknown, (? mm).

Type locality not known, [presumably Corsica].

Nomenclatural status, an available name, an invalid name, a homonym of *C. fasciatus* Schröter, 1803, of Perry, 1811, of Kiener, 1845 (1850) and of *C. fasciata* Menke, 1828.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.

fasciatus Schröter, 1803.

Published in Arch. Zool. Zoot. 3 (2): p. 69, not figured.

Holotype whereabouts unknown, (app. 33 x 26 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost holotype, no locality or figure and an inadequate description.

fasciatus Perry, 1811.

Published in Conchology: pl. 24, fig. 3.

Holotype in NHMUK, $(50 \times 29 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name, an invalid name, a homonym of *C. fasciatus* Schröter, 1803.

Taxonomic status, a synonym of *C. genuanus* Linnaeus, 1758.





fasciatus "Meuschen" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 390, no. 66, not figured.

Specimen named whereabouts unknown, cited species in Meuschen, 1766, p. 354, no. 994, not figured.

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. coffeae* Gmelin, 1791.

Taxonomic status, none, (a specimen of *C. coffeae* Gmelin, 1791).

fasciatus Kiener, 1845 (1850).

Published in Coq. Viv. 2: pl. 109, fig. 2, (1850, Coq. Viv. 2: p. 311, no. 277).

Holotype was in collection Lorois, present whereabouts unknown, $(42 \times ? \text{ mm})$, $(\text{fig. } 42 \times 21 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. fasciatus* Schröter, 1803, of Perry, 1811 and of *C. fasciata* Menke, 1828, renamed *C. anadema* Tomlin, 1937. Taxonomic status, a synonym of *C. splendidulus* Sowerby I & II, 1833.



fasciatus A. Adams, 1854 ("1853").

Published in Proc. zool. Soc. Lond. pt. 21, no. 255: p. 119, no. 20, not figured.

Lectotype (C, M & W) in NHMUK, $(27 \times 14 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name, an invalid name, a homonym of *C. fasciatus* Schröter, 1803, of Perry, 1811, of Kiener, 1845 (1850), of *C. fasciata* Menke, 1828 and of Requien, 1848, renamed *C. bifasciatus* Sowerby, 1857.

Taxonomic status, a synonym of C. attenuatus Reeve, 1844.





fasciatus "Martini" Sowerby II, 1858.

Published in Thes. Conch. 3 (*Conus*), pt. 18: p. 35, no. 306, pl. 20 (206), figs 487 - 489.

Type series whereabouts unknown, cited figure in Martini, 1773, pl. 52, fig. 573, (fig. 58 x 23 mm), (figs in T. C. 69 x 31, 69 x 29 & 66 x 30 mm).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. fasciatus* Schröter, 1803, of Perry, 1811, of Kiener, 1845 (1850), of Adams, 1854, of *C. fasciata* Menke, 1828, and of Requien, 1848.

Taxonomic status, a synonym of *C. ochroleucus* Gmelin, 1791, figure cited is lectotype of *C. ochroleucus* Gmelin, 1791 and of *C. eburneus* (Röding), 1798.



Figures in Thes. Conch. 3.



Cited figure in Martini.

<u>fascifer</u> (forma cholmondeleyi Melvill) Lauer, 1987.

Published in Rossiniana (36): p. 13, no. 18, fig. 14b.

Specimens described whereabouts unknown, cited figures in

Basteria 47 (5 - 6): p. 140, figs 411a & b, (figs 48 x 25 mm).


Locality Mozambique, Tanzania & South East Africa.

Nomenclatural status, an unavailable name (nomen nudum),

described as a form after 1960.

Taxonomic status, none, (a specimen of *C. textile* Linnaeus,

1758).



Cited figure in Basteria.

felinus Link, 1807.

Published in Beschr. Nat.-Samml. Univ. 3: p. 104, not figured. Type series whereabouts unknown, (? mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. spectrum Linnaeus, 1758.

felitae Rolán, 1990.

Published in Iberus, Sup. 2: p. 45, pl. 1, fig. 18, pl. 2, fig. 18, pl. 3, fig. B, pl. 6, fig. D & pl. 8, fig. C

Holotype in MNCM, $(12.7 \times 6.6 \text{ mm})$.

Type locality Rabo de Junco, northern Mordeira Bay, Sal Island, Cape Verde Islands, (rocks, 0.5 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





felix Fenzan, 2012.

Published in Acta Conch. 11: p. 11, pls 1 & 2. Holotype in MNHN, no.24818, $(28.20 \times 15.20 \text{ mm})$.

Type locality Umkomaas, Mkomazi River mouth, 50 km south of Durban, Kwa-zulu Natal, South Africa, (dredged in 90-100 meters on rubble overgrown by sponges and soft coral). Nomenclatural status, an available name.

Management at the second of a second





fenellus "Chemnitz" E.A. Smith, 1876.

Published in J. Linn. Soc. 12 (64): p. 535, not figured.

Specimen described whereabouts unknown, cited figure in Chemnitz, 1795, pl. 183, figs 1782 & 1783, (fig. 1782, 36 x 16 & fig. 1783, 34 x 16 mm).

Type locality not mentioned.

Nomenclatural status, an unavailable name, (nomen nudum), cited as a synonym of *C. magus* Born, 1778.

Taxonomic status, none, (a specimen of C. magus Born, 1778).



Cited figure in Chemnitz.

Dauciconus fenzani Petuch & Sargent, 2011.

Published in Visaya vol. 3, no. 3: p. 40, pl. 2, fig. 1.

Holotype in LACM, $(47.1 \times 21.8 \text{ mm})$.

Type locality NW of Isla Coiba, Golfo de Chiriqui, Panama, (dredged from 150 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. poormani Berry, 1968.





fergusoni Sowerby II, 1873.

Published in Proc. zool. Soc. Lond. unnumbered (41), pt. 1: p. 145, pl. 15, fig. 1.

Type series whereabouts unknown, (fig. 145 x 85 mm).

Type locality Panama, (West Coast).

Nomenclatural status, an available name.

Taxonomic status, a valid species, although a junior synonym of *C. fulvocinctus* Crosse, 1872, *C. fergusoni* takes precedence (nomen protectum), ICZN Opinion 1502, (1988), see Section 1 B (5).




Figure in Pro. Zoo. Soc.

fernambucinus (Röding), 1798.

Published in Mus. Bolten 2: p. 39, no. 492/18, not figured. Lectotype representative (Kohn) figure in Martini, 1773, pl. 57, fig. 632, (fig. 38 x 25 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. taeniatus Hwass, 1792.



fernandesi Tenorio, Afonso & Rolàn, 2008.

Published in Vita Malacologica 6: p. 5, pl. 2, figs 1 - 11.

Holotype in MNCM, $(21.0 \times 13.0 \text{ mm})$.

Type locality Porto Novo, South coast of Santo Antão Island, Cape Verde Islands, (on the rocky walls of the bay of the harbour, on algae, in crevices and under stones, in 2 to 4 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





ferrugatus Sowerby I & II, 1834.

Published in Proc. zool. Soc. Lond. pt. 2, no. 15: p. 19, not figured.

Type series whereabouts unknown, $(37.4 \times 18.4 \text{ mm})$.

Type locality Gulf of California near Guayamas Island.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. monilifer* Sowerby, 1833, (a synonym (form) of *C. regularis* Sowerby, 1833).

ferrugineus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 649, no. 49, (1798, Tab. Enc. pl. 326, fig. 4).

Lectotype (Kohn) in MHNG, $(52 \times 28 \text{ mm})$.

Type locality Guinea, (West Africa), (erroneous), corrected (Delsaerdt) New Guinea.

Nomenclatural status, an available name.

Taxonomic status, a valid species





ferrugineus Bosc, 1801.

Published in Hist. Nat. Coq. 5: p. 128, not figured.

Type series whereabouts unknown, figures cited, in Martini, 1773, pl. 59, figs 660 & 662, (fig. 660, 58 x 36 & fig. 662, 39 x 23 mm).

Type locality coast of Guinea, [presumably New Guinea]. Nomenclatural status, an available name, an invalid name, a homonym of *C. ferrugineus* Hwass, 1792 and of *C. ferrugineus* (Röding), 1798.

Taxonomical status, a synonym of C. capitaneus Linnaeus, 1758.





Cited figures in Martini.

ferugineus (Röding), 1798.

Published in Mus. Bolten 2: p. 41, no. 515/38, not figured. Lectotype representative (Kohn) figure in Martini, 1773, pl. 56,

fig. 626, (fig. $45 \times 26 \text{ mm}$).

Type locality not mentioned. Nomenclatural status, an available name, an invalid name, a homonym of *C. ferrugineus* Hwass, 1792.

Taxonomic status, a synonym of C. spurius Gmelin, 1791.



feruginosus Mawe, 1823.

Published in The Linnean System of Conchology: p. 90, not figured.

Specimen(s) named whereabouts unknown, (? mm).

Locality South seas.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

festivus Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 413, no. 116, not figured.

Lectotype (Kohn) in ZMUC, (23 x 13 mm, orig. desc. app. 25 x 14 mm).

Type locality Moluccas Islands, (Indonesia), (Valentyn).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. pertusus Hwass, 1792.



figulinus Linnaeus, 1758.


Published in Syst. Nat. 10th ed. (1): p. 715, no. 267, not figured.

Lectotype (Kohn) in LSL, $(65 \times 42 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.





fijiensis Moolenbeek, Röckel & Bouchet, 2008.

Published in Vita Malacologica 6: p. 38, pl. 2, figs 13 - 14 & pl. 4, fig. 40.

Holotype in MNHN, (17.8 x 6.8 mm).

Type locality Fiji, South-East of Viti Levu, (80 - 120 mtrs). Nomenclatural status, an available name.

Taxonomic status, a valid species.





fijisulcatus Moolenbeek, Röckel & Bouchet, 2008.

Published in Vita Malacologica 6: p. 38 & 43, pl. 3, figs 27 - 28 & pl. 4, fig. 42.

Holotype in MNHN, $(54.9 \times 25.6 \text{ mm})$.

Type locality Fiji, Vanua Levu, Natewa Bay, (220 - 224 mtrs).

Nomenclatural status, an available name.





filamentosus Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 6, sp. 260 (310). Holotype in NHMUK, (28.5 x 13.5 mm).

Type locality not known, designated (Filmer) the Philippines. Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. conspersus* Reeve, 1844, (a synonym (form) of *C. spectrum* Linnaeus, 1758).





(Ammirales) filicinctus Schepman, 1913.

Published in The Prosobranchia of the Siboga Expedition, pt. 5, Toxoglossa: p. 384, pl. 25, fig. 1.

Holotype in NBC-ZMA, $(27.5 \times 15 \text{ mm})$.

Type locality Madura Bay, (East Java), Indonesia, (sand, 69 - 91 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (subadult) of *C. voluminalis* Reeve, 1943





filmeri Rolán & Röckel, 2000.

Published in Argonauta 8 (2): p.35, figs 77 - 80.

Holotype in SMNS, $(33.1 \times 19.5 \text{ mm})$.

Type locality Saco Mar, southern Angola, (in calm water at 2 mtrs depth, under rocks, buried in fine sand).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





filosus (Röding), 1798.


Published in Mus. Bolten 2: p. 44, no. 559/69, not figured. Lectotype representative (Kohn) figure in Favanne & Favanne,

1780, pl. 15, fig. C, (fig. 49 \times 30 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. maldivus Hwass, 1792.



finkli Petuch, 1987.

Published in New Caribbean Molluscan Faunas: p. 111, pl. 25, figs 10 & 11.

Holotype in USNM, $(29 \times 14 \text{ mm})$.

Type locality off Punto Fijo, Falcon, Gulf of Venezuela, (dredged, 35 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. cancellatus* Hwass, 1792.





fischoederi Röckel & da Motta, 1983.

Published in Bull. Inst. malac. Tokyo, 1 (8): p. 117, pl. 40, figs 3 & 4.

Holotype in NMSF, $(33.7 \times 16.4 \text{ mm})$.

Type locality Coron, Palawan, Philippines, (sand & coral rubble, 100 ft).

Nomenclatural status, an available name.





flamingo Petuch, 1980.

Published in Proc. biol. Soc. Wash. 93 (2): p. 299, fig. 1. Holotype in USNM, $(19.4 \times 9.5 \text{ mm})$.

Type locality off Dania, Broward County, Florida, (47.5 mtrs). Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. flavescens* Sowerby, 1834.





flammeacolor Petuch, 1992.

Published in Conchiglia 23 (264): p. 38, figs 5 & 6 Holotype in CMNH, $(10.5 \times 6 \text{ mm})$.

Type locality Cabo Falso and Cayos Vivorillo, Honduras, (East coast), (coral rubble, 20 mtrs).

Nomenclatural status, an available name.



flammeus (Röding), 1798.

Published in Mus. Bolten 2: p. 44, no. 550/61, not figured. Holotype whereabouts unknown, cited figures in Martini, 1773, pl. 55, figs 606 & 607, (fig. 606, 42 x 25 & fig. 607, 61 x 33 mm). Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), an objective junior synonym of *C. achatinus* (Röding), 1798, (fig. 606, same type) and of *C. barathrum* (Röding), 1798, (fig. 607, same type), (fig. 606 is a synonym of *C. eburneus* Hwass, 1792 and fig. 607 is a synonym of *C. ermineus* Born, 1778).





Cited figures in Martini.

flammeus Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 279, no. 100, not figured.

Lectotype representative (Walls) figure in Tab. Enc. 1798, pl. 336, fig. 1, (fig. 49 \times 26 mm).

Type locality Africa, [erroneous].

Nomenclatural status, an available name, an invalid name, a homonym of C. flammeus (Röding), 1798, renamed C. phlogopus Tomlin, 1937.

Taxonomic status, a subspecies of C. spurius Gmelin, 1791.



<u>flammulata</u> (variety mediterraneus Hwass) Bucquoy, Dautzenberg & Dollfus, 1882.

Published in Mollusques Marins Roussillon, (Paris), 1 (2): p. 83, pl. 13, fig. 16.

Type series whereabouts unknown, (fig. $17 \times 10 \text{ mm}$).

Type locality Barbary, (North Africa).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.



Textilia solangeae f. "flammulata" Bozzetti, 2010.


Published in Malacologia no.68, p.13 & figs.

Specimens illustrated whereabouts unknown (no dimensions given). Type locality Lavanono, Madagascar.

Nomenclatural status, an unavailable name, described as a form after 1961.

Taxonomic status, none, (specimens are *C. solangeae* Bozzetti, 2004).





Specimens illustrated in Malacologia.

flavescens "Gray" Sowerby I & II, 1834.

Published in Conch. Ill. (Conus): pl. 55, fig. 68.

Lectotype (Kohn) in NHMUK, $(23.5 \times 12 \text{ mm})$.

Type locality S. Australia, (erroneous), corrected (Vink) Bahamas.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





flavescens "Gray" Kiener, 1845 (1846).

Published in Coq. Viv. 2: pl. 68, fig. 1, (1846, Coq. Viv. 2: p. 58, no. 48).

Type series was in collection Dupont, present whereabouts unknown, (20 x ? mm), (fig. 20 x 11 mm).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. flavescens* Sowerby, 1833.

Taxonomic status, a synonym of C. cardinalis Hwass, 1792.



flavescens (variety mediterraneus Hwass) Coen, 1933.

Published in Memorie R. Comit. talassogr. ital. (192): p. 71, not figured.

Specimen(s) described in collection Danilo Sandri in MCG, destroyed in World War 2, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. pallida* Bucquoy, Dautzenberg & Dollfus, 1882 and not described, indicated or figured.

Taxonomic status, none, (a specimen of *C. ventricosus* Gmelin, 1791).


<u>flavescens</u> (subspecies *imperialis* Linnaeus) Barros E Cunha, 1933. Published in Mems Estud. Mus. zool. Univ. Coimbra, 1 (71): p. 18, no. 10, not figured.

Two syntypes in MZUC, $(42 \times 25 \& 39 \times 20.5 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of C. flavescens Sowerby, 1834 and of Kiener, 1845 (1846).

Taxonomic status, a synonym of C. imperialis Linnaeus, 1758.



Syntype 1 42 x 25 mm.



Syntype 2 39 x 20.5 mm.

flavidus Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 265, no. 57, not figured.

Holotype in MHNG, $(63 \times 36 \text{ mm})$.

Type locality Guinee, (West Africa), (erroneous), corrected (Wils) New Guinea.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





flavocinctus Link, 1807.

Published in Beschr. Nat.-Samml. Univ. 3: p. 103, not figured Type series whereabouts unknown, cited figure in Martini, 1773, pl. 83, (error for 53) fig. 588, (fig. 40 x 23 mm). Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost type, no locality an unclear figure and an inadequate description.



Cited figure in Martini.

flavus Röckel, 1985.

Published in Spixiana 8 (2): p. 166, figs 1a, 2a, 3b & 5a. Holotype in ZSM, $(55.1 \times 24.7 \text{ mm})$.

Type locality Batangas Bay, Luzon, Philippines, (10 mtrs). Nomenclatural status, an available name.

Taxonomic status, a valid species.



flavusalbus Rolán & Röckel, 2000.

Published in Argonauta 8 (2): p. 38, figs 92 - 96.

Holotype in MNCM, $(23.7 \times 14.3 \text{ mm})$.

Type locality Das Pipas Bay, Angola, (under stones & rocks in 2 - 3 mtrs).

Nomenclatural status, an available name.





fletcheri Petuch & Mendenhall, 1972.
Published in Veliger 15 (2): p. 96, figs 1 & 2.

Holotype in CAS, $(105 \times 44.5 \text{ mm})$.

Type locality Southwest of the Penghu Island Group, Taiwan, (trawled, 90 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. pergrandis* Iredale, 1937.





flindersi Brazier, 1898 ("1897").

Published in Proc. Linn. Soc. N.S.W. 22 (4) no. 88: p. 780, no. 3, not figured.

Holotype in SAMA, (28 x 15 mm, orig. desc. 29 x 19 mm). Type locality Flinders Island, (Victoria, Australia), (under stones).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. anemone Lamarck, 1810.



floccatus Sowerby I & II, 1841.

Published in Conch. Ill. (Conus): Index, p. 4, no. 112, (1839, pl. 153, no. 112 was wrongly identified as C. dux Lamarck, 1810). Lectotype representative (Delsaerdt) figure in C. I. 1839, pl. 153, fig. 112, (fig. 63 x 29 mm).

Type locality not mentioned, (Luzon, Philippines in Thes. Conch. 3).

Nomenclatural status, an available name, Sowerby implied it was a new replacement name (nomen novum) for $C.\ dux$ Lamarck, 1810, however Lamarck did not introduce this name.



floridanus Gabb, 1869.

Published in Am. J. Conch. 4 (4): p. 195, pl. 15, fig. 4.

Holotype in ANSP, $(48.5 \times 23 \text{ mm})$.

Type locality Tampa Bay, Florida.

Nomenclatural status, an available name.

Taxonomic status, a junior synonym of *C. anabathrum* Crosse, 1865, an application to establish *C. floridanus* Gabb as having precedence over *C. anabathrum* Crosse was rejected, ICZN Opinion 1539, (1989), see Section 1 B (6).





floridensis Sowerby II, 1870.

Published in Proc. zool. Soc. Lond. unnumbered (38), pt. 2: p. 256, pl. 22, fig. 11.

Holotype in NHMUK, $(42.5 \times 20 \text{ mm})$.

Type locality Florida, (Mr. Waters).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. anabathrum Crosse, 1865.





floridulus A. Adams & Reeve, 1848.

Published in The Zoology of the Voyage of "HMS Samarang" Mollusca (London), no. 3, pt. 1: p. 18, pl. 5, figs 9a & b.

Holotype in NHMUK, $(37 \times 20 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





floridus Sowerby II, 1858.

Published in Thes. Conch. 3 (Conus), pt. 18: p. 47, no. 345, frontispiece fig. 558.

Two syntypes in NHMUK, (69 x 35 & 77 x 39 mm, latter figured). Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. striatus* Linnaeus, 1758.









Syntype no.2.

Syntype no.1 figured.

fluctifer Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 382, no. 49, not figured.

Lectotype representative (Walls) figure in Tab. Enc. 1798, pl. 331, fig. 5, (fig. 138 x 64 mm, orig. desc. app. 127 x 64 mm). Type locality Java, Mozambique & Zanzibar (Favanne & Favanne, 1780), (erroneous).

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of *C. prometheus* Hwass, 1792, same type, (a synonym of *C. pulcher* Lightfoot, 1786).



"Jaspidiconus fluviamaris" Petuch & Sargent, 2011.

Published in Rare and Unusual Shells from the Florida Keys: pl.3-14, fig.D. (published July 2011)

Specimen named whereabouts unknown (16.2 mm)

Locality off Hillsboro Inlet: also found off Key Largo,

Plantation Key and the dry Tortugas.

Nomenclatural status, an unavailable name (nomen nudum) not intended to describe a new species.

Taxonomic status, none (a specimen of *C. vanhyningi* Rehder, 1994, (a synonym (form) of *C.mindanus* Hwass, 1792)



Plate 3-14, figure D.

Jaspidiconus **fluviamaris** Petuch & Sargent, 2011.

Published in Visaya Vol. 3, no. 3: p. 45, pl. 5, figs 1 - 5.

Holotype in LACM, $(16.2 \times 8.5 \text{ mm})$.

Type locality past the Second Reef, off Hillsboro Inlet, Pompano Beach, Broward County, Florida, (on sand bottom, 10 mtrs). Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. mindanus Hwass, 1792.





folliculus (Röding), 1798.

Published in Mus. Bolten 2: p. 40, no. 503/28, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

fontonae Rolán & Trovão in Rolán, 1990.

Published in Iberus, Sup. 2: p. 28, pl. 1, fig. 10, pl. 2, fig. 10, pl. 3, fig. C, pl. 6, fig. C & pl. 8, fig. B.

Holotype in MNCM, $(18.7 \times 11.3 \text{ mm})$.

Type locality Fontona Bay, Sal Island, Cape Verde Islands, (stones & rubble).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





"(Genuanoconus) genuanus f. foscaclaudiae" Assi, 2010.

Published in Malacologia no. $\overline{67}$, p.14 & figs.

Specimen described was to be deposited in MNHN but not received and not illustrated, $(49.22 \times 29.28 \text{ mm})$.

Type locality Senegal.

Nomenclatural status an unavailable name, described as a form after 1961.

Taxonomic status, none (a specimen of *C. genuanus* Linnaeus, 1758).

fosteri Clench & Aguayo in Clench, 1942.

Published in Johnsonia 1 (6): p. 34, pl. 12, fig. 5.

Holotype in MCZ, $(61.5 \times 27 \text{ mm})$.

Type locality off Sagua la Grande, Santa Clara Province, Cuba, (about 260 fths).

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. villepinii* Fischer & Bernardi, 1857.





fragilissimus Petuch, 1979.

Published in Nemouria (23): p. 14, figs 22 - 25.

Holotype in DMNH, (30 x 12 mm), (not present in May 1988, a paratype is present, $34 \times 15 \text{ mm}$).

Type locality off South coast of Harmil Isl., Dahlak Archipelago, Eritrea Province, Ethiopia, (coral rubble, 3 mtrs).

Nomenclatural status, an available name.





Paratype in DMNH

franciscanus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 698, no. 87, (1798, Tab. Enc. pl. 337, fig. 5).

Lectotype (Kohn) in MHNG, $(55 \times 30.5 \text{ mm})$.

Type locality Africa.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. guinaicus Hwass, 1792.





<u>franciscanus</u> (variey mediterraneus Hwass) Philippi, 1836. Published in Enumeratio Molluscorum Siciliae 1, (Berlin): p. 238, pl. 12, fig. 18.

Holotype in ZMB, $(36 \times 15 \text{ mm})$.

Type locality not mentioned, (presumably Sicily, Italy). Nomenclatural status, an available name, an invalid name, a homonym of *C. franciscanus* Hwass, 1792, (although not proposed as a new variety (p. 238) it is so indicated on pl. 12, fig. 18). Taxonomic status, a synonym of *C. ventricosus* Gmelin, 1791.

franciscoi Rolán & Röckel, 2000.

Published in Argonauta 8 (2): p. 36, figs 82 - 86.

Holotype in MNCM, $(28.4 \times 15.6 \text{ mm})$.

Type locality Chapeu Armado, Angola, (buried in sand between stones).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





<u>frauenfeldi</u> Crosse, 1865.

Published in J. Conch. Paris, 13: p. 307, no. 7, pl. 10, figs 1 & 1a.

Lectotype (Tomlin) in NHMUK, $(43.2 \times 23.6 \text{ mm. orig. desc. } 42 \times 23 \text{ mm})$.

Type locality Madagascar, (erroneous), corrected (G. Raybaudi) Padang, South West Sumatra, East Indian Ocean.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. magus Linnaeus, 1758.





frausseni Tenorio & Poppe, 2004.

Published in Visaya vol. no. 1: p. 20, pl. 1, figs 1 - 3.

Holotype in NMPM, $(47.0 \times 21.0 \text{ mm})$.

Type locality offshore Aliguay Island, Philippines, (80 - 200 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





fraxineus (Röding), 1798.

Published in Mus. Bolten 2: p. 40, no. 506/31, not figured. Lectotype representative (Kohn) figures in Chemnitz, 1788, pl.

138, figs 1277 & 1278, (figs 49 x 31 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. glaucus Linnaeus, 1758.



Eugeniconus friedae da Motta, 1991.

Published in Conchiglia 22 (258): p. 12, & figs.

Holotype in MHNG, $(50.7 \times 25 \text{ mm})$.

Type locality Dutch Bay Point, Northwest of Colombo, Sri Lanka, (10 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. nobilis* Linnaeus, 1758.





frigidus Reeve, 1848.

Published in Conch. Icon. 1 (Conus): Sup. pl. 3, sp. 284.

Holotype in NHMUK, $(26 \times 15 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





frisbeyae Clench & Pulley, 1952.

Published in Tex. J. Sci. 4 (1): p. 59, pl. A, fig. 1.

Holotype in MCZ, $(32.5 \times 14.4 \text{ mm})$.

Type locality Campeche Banks, Yucatan, Mexico, (East coast), (trawled, 23 fths).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. armiger Crosse, 1853.





fritillaria Lichtenstein, 1794.

Published in Catalogus rerum naturalium rarissamarum 2: p. 31, no. 171, not figured.

Type material whereabouts unknown, dimensions not mentioned. Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost type, not figured, no locality and an inadequate description.

frostiana Brazier, 1898 ("1897").

Published in Proc. Linn. Soc. N.S.W. 22 (4), no. 88: p. 781, no. 5, not figured,

Holotype in SAMA, $(18 \times 10 \text{ mm})$.

Type locality Solomon Islands, (Kenyon).

Nomenclatural status, an available name.

Taxonomic status, a synonym (subadult) of *C. monachus* Linnaeus, 1758.





fucatus Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 7, sp. 271 (321). Two syntypes in NHMUK, (18.7 x 9.2 mm. figured & 15.9 x 8.1 mm). Type locality Philippines.

Nomenclatural status, an available name.

Taxonomic status, a synonym (juvenile) of $C.\ magus$ Linnaeus, 1758.









Syntype no. 1 Figured.

Syntype no.2.

fulgens "S" Lightfoot, 1786.

Published in Cat. Portland Mus: p. 8, no. 142, p. 115, no. 2533, p. 148, no. 3255, & p. 161, no. 3510, not figured.

Specimen(s) named sold at auction, present whereabouts unknown, (
? mm).

Locality China, (nos 0142 & 3255).

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

"fulgens" "Röckel et al" Anonymous, 1993.

Published in World Shells (4): p. 92, not figured.

Specimens named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name, named but not described, indicated or figured and no author, Röckel did not introduce this name.

Taxonomic status, none, (stated to be a specimen of C. stramineus Lamarck, 1810).

fulgetrum Sowerby I & II, 1834.

Published in Conch. Ill. (Conus): pl. 56, fig. 82.

Lectotype (Wils) in NMWC, $(22 \times 14 \text{ mm})$.

Type locality not mentioned, (Red Sea, Thes. Conch. 3).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. miliaris Hwass, 1792.





fulgurans Hwass in Bruquière, 1792.

Published in Enc. Méth. 1: p. 687, no. 76, (1798, Tab. Enc. not figured).

Lectotype representative (Kohn) figure in Martini, 1773, pl. 58, fig. 644. (fig. $39 \times 19 \text{ mm}$).

Type locality Africa.

Nomenclatural status, an available name.

Taxonomic status, an objective junior synonym of *C. fulmineus* Gmelin, 1791, same type, (a nomen dubium).



fulmen Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 39, sp. 215, (Reeve states, in error, that this species is described in Proc. zool. Soc. Lond. pt. 11, no. 130),

A syntype in NMWC, $(71.5 \times 35 \text{ mm. figured})$.

Type locality Capul Island, Philippines, (on reefs).

Nomenclatural status, an available name.

Taxonomic status, a valid species, although a junior synonym of *C. modestus* Sowerby, 1833, *C. fulmen* takes precedence (nomen protectum), ICZN Opinion 1699, (1992), see Section 1 B (7).





fulmineus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3388, no. 33, not figured.

Holotype was in collection B. Feldmann, present whereabouts unknown, holotype representative (Kohn) figure in Martini, 1773, pl. 58, fig. 644, (fig. 39 x 19 mm). Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost holotype, no locality, an unclear figure and a limited description.



fultoni Sowerby III, 1887.

Published in Thes. Conch. 5 (*Conus*), pt. 44: 2nd Sup. p. 273, no. 527, pl. 36 (512*), fig. 758.

Holotype in NHMUK, $(23.4 \times 14.1 \text{ mm})$.

Type locality Singapore, uncertain.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. ruppellii* Reeve, 1848 (a synonym (form) of *C. boeticus* Reeve, 1844.





fulvobullatus da Motta, 1982.

Published in Publcoes. Occ. Soc. port. malac. (1): p. 13, fig. 12.

Holotype in MHNG, $(52.5 \times 25 \text{ mm})$.

Type locality off East coast of Malaysia, (trawled).

Nomenclatural status, available name.

Taxonomic status, a synonym (form) of C. magus Linnaeus, 1758.





fulvocinctus Crosse, 1872.

Published in J. Conch. Paris, 20: p. 214, not figured, (1873, J. Conch. Paris, 21: p. 248, pl. 11 fig. 5)

Conch. Paris, 21: p. 248, pl. 11, fig. 5).

Holotype in NHMUK, $(72.7 \times 42 \text{ mm})$.

Type locality West Africa, (erroneous).

Nomenclatural status, an available name.

Taxonomic status, a forgotten name (nomen oblitum), although a senior synonym of C. fergusoni Sowerby, 1873, C. fergusoni takes precedence (nomen protectum), ICZN Opinion 1502, (1988), see Section 1 B (5).





fulvostriatus Fenaux, 1942.

Published in Bull. Inst. oceanogr. Monaco, (814): p. 2, fig. 4.

Type series whereabouts unknown, (fig. 71 x 39 mm).

Type locality Ile de Bourbon, (Reunion).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. quercinus Lightfoot, 1786.



Figure in Bull. Inst.

fulvus Schröter, 1803.

Published in Arch. Zool. Zoot. 3 (2): p. 73, not figured.

Holotype whereabouts unknown, (app. 39 x 20 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost holotype, no figure or locality and an inadequate description.

fulvus Sowerby III, 1889.

Published in J. Conch. Lond. 6 (1): p. 10, pl. 1, fig. 1.

Holotype was in collection S.D. Bairstow possibly in OMNH but no longer identifiable, (fig. $21 \times 11 \text{ mm}$).

Type locality South Africa.

Nomenclatural status, an available name, an invalid name, a homonym of *C. fulvus* Schröter, 1803.

Taxonomic status, a synonym of C. tinianus Hwass, 1792.



Figure in Journ. Conch.

fulvus Fenaux, 1943.

Published in Bull. Inst. oceanogr. Monaco, (835): p. 4, fig. 8. Type series whereabouts unknown, (fig. 28 x 13 mm). Type locality New Guinea.

Nomenclatural status, an available name, an invalid name, a homonym of *C. fulvus* Schröter, 1803 and of Sowerby, 1889. Taxonomic status, doubtful (nomen dubium), lost type, an inadequate description and an unclear figure, possibly a synonym (form) of *C. magus* Linnaeus, 1758.



Figure in Bull. Inst.

fumigatus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 704, no. 94, (1798, Tab. Enc. pl. 336, fig. 7).

Lectotype (Kohn) in MHNG, (35 x 21 mm). Type locality American Seas, (erroneous). Nomenclatural status, an available name. Taxonomic status, a valid species.





<u>furnae</u> (subspecies *verdensis* Trovão) Rolán, 1990.

Published in Iberus, Sup. 2: p. 42, pl. 1, fig. 9 & pl. 2, fig. 9.

Holotype in MNCM, $(21.1 \times 13 \text{ mm})$.

Type locality Furna Bay, Brava Island, Cape Verde Islands, (under rocks).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. verdensis* Trovão, 1979.





furvus Reeve, 1843.

Published in Conch. Icon. 1 (Conus): pl. 13, sp. 69 (published May), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 79, not figured, published Dec).

A probable syntype in NMWC, (45.4 x 19 mm), 3 syntypes recently rediscovered in NHMUK, (43.40 x 19.85, 41.10 x 20.10 & 41.20 x 18.90 mm).

Type locality Ticao & Masbate Isles, Philippines, (in sandy mud, low tide).

Nomenclatural status, an available name.

Taxonomic status, a valid species.













Syntype 3 in NHMUK. ©NHMUK.

Syntype 2 in NHMUK. ©NHMUK





Syntype 4 in NHMUK. ©NHMUK.

<u>fusca</u> (variety mediterraneus Hwass) Bucquoy, Dautzenberg & Dollfus, 1882.

Published in Mollusques Marins Roussillon, (Paris), 1 (2): p. 83, pl. 13, fig. 14.

Type series whereabouts unknown, (fig. 20 x 9 mm).

Type locality Barbary, (North Africa), (in sponges).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.





fuscatus Born, 1778.

Published in Ind. Mus. Caes. Vindob: p. 126, (1880, Test. Mus. Caes. p. 148, not figured).

Lectotype (Kohn) in NHMW, $(53 \times 31 \text{ mm})$.

Type locality Mauritius, (Martini).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. imperialis Linnaeus, 1758.





fuscocingulatopsis de Gregorio, 1885.

Published in Boll. Soc. malac. ital. 11: p. 94, not figured.

Type series in MPUP but no longer identifiable, (? mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.

<u>fuscoflavus</u> Röckel, Rolán & Monteiro, 1980.

Published in Cone Shells from Cape Verde Islands: p. 71, figs 38 - 41.

Holotype in NMSF, $(20.3 \times 12.1 \text{ mm})$.

Type locality Sal Rei, Boavista Isl., Cape Verde Islands, (on rocks with algae).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





fuscolineatus Sowerby III, 1905.

Published in Proc. malac. Soc. Lond. 6 (5): p. 282, fig. 6. Holotype in NHMUK, (22 x 12.5 mm).

Type locality Sierra Leone, (probably erroneous), designated (Rolán & Röckel) Ponta de Noronha, near Moçamedes Bay, Angola. Nomenclatural status, an available name. Taxonomic status, a valid species.





fuscomaculatus E.A. Smith, 1877.

Published in Ann. Mag. nat. Hist. 4 (19): p. 224, not figured. Holotype in NHMUK, (37.5 x 16 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. stramineus Lamarck, 1810.





fuscoolivaceus (variety catus Hwass) Dautzenberg, 1937.

Published in Mem. Mus. r. Hist. nat. Belg. 2 (18): p. 62, pl. 1, fig. 8.

Holotype in IRSN, $(30.4 \times 17 \text{ mm})$.

Type locality Tjilaeot, Eureun, (West Java, Indonesia), (de Priester).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. catus Hwass, 1792.





fuscus (subspecies lugubris Reeve) Barros E Cunha, 1933.

Published in Mems Estud. Mus. zool. Univ. Coimbra, 1 (71): p. 194, no. 167, not figured.

Eight syntypes in MZUC, (from $15.5 \times 9 \text{ mm}$ to $11 \times 7 \text{ mm}$). Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.





Syntype 15 x 8.4 mm in MZUC.

<u>fusiformis</u> G. Fischer von Waldheim, 1807, (originally spelt fusifornis in error, corrected (Kohn).

Published in Mus. Démidoff 3: p. 144, nos 179 - 180, not figured. Lectotype (Kohn), (no. 179) in ZMMU, (44.5 \times 19 mm, orig. desc. no. 180, 65 \times 27 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. glans Hwass, 1792.



fusiformis Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 423, no. 129, not figured.

Holotype in MHNG, $(48 \times 21 \text{ mm})$.

Type locality Pacific Ocean, uncertain.

Nomenclatural status, an available name, an invalid name, a homonym of C. fusiformis Fischer, 1807, renamed C. attractus Tomlin, 1937.

Taxonomic status, a synonym of C. compressus Sowerby, 1866.





fusiformis Pease, 1861 ("1860").

Published in Proc. zool. Soc. Lond. unnumbered (28), pt. 3: p. 398, not figured.

Lectotype (Kay) in NHMUK, $(6.5 \times 3 \text{ mm})$.

Type locality Sandar Island, (Sandwich Island, Hawaii).

Nomenclatural status, an available name, an invalid name, a homonym of *C. fusiformis* Fischer, 1807 and of Lamarck, 1810, renamed *C. parvus* Pease, 1868.

Taxonomic status, not in Family Conidae in Family Turridae.





fustigatus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 623, no. 18, (1798, Tab. Enc. pl. 320, fig. 1).

Lectotype (Kohn) in MHNG, $(41 \times 27 \text{ mm})$.

Type locality Asiatic Ocean.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. pulicarius* Hwass, 1792, (p. 622, no. 17).





fusus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3390, no. 45, not figured.

Lectotype representative (Kohn) figure in Martini, 1773, pl. 52, fig. 576, (fig. 20 \times 11 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym (juvenile) of C. terebra Born, 1778.



<u>futunaensis</u> (subspecies neptunus Reeve) Moolenbeek & Röckel, 1996.

Published in Bull. Mus. natn. Hist. nat. Paris 4 (18), sect. A (3 - 4): p. 392, figs 1 - 3.

Holotype in MNHN, $(45.5 \times 19.5 \text{ mm})$.

Type locality Futuna Island, (South Pacific), (355 - 369 mtrs). Nomenclatural status, an available name.

Taxonomic status, a subspecies of C. neptunus Reeve, 1843.

