

Section HI habui to istriensis

Copyright © 2011.R Filmer Conditions of use.

The content of this website is provided for personal and scientific use and may be downloaded for this purpose. It may not be used in whole or part for any commercial activity and publication of any of the content on the internet is limited to the Cone Collector website (www.TheConeCollector.com). Authors wishing to publish any of the pictures may do so on a limited basis but should inform M Filmer so that the original owner of the rights to the picture can be acknowledged. (mike@mfilmer.fsnet.co.uk)

Version 1.0 October 2011.

habui Lan, 2003 ("2002").

Published in Bull. Malac. R of China 26: p. 1, figs 1 & 2. Holotype in NMNS, (30 x 15.2 mm).

Type locality exact locality not made available, area around Okino-daito-jima east of Okinawa Japan to the Philippines. (trawled by coral boats in fairly deep water).

Nomenclatural status, an available name.

Taxonomic status, a synonym (subadult) of *C. samiae* da Motta, 1982 (a synonym (form) of *C. sulcatus* Hwass, 1792).





halitropus
Published in Proc. biol. Soc. Wash. 56: p. 88, not figured.
Holotype in USNM, (32.5 x 13.8 mm).
Type locality Honolulu, Hawaii, (dredger dumpings).
Nomenclatural status, an available name.
Taxonomic status, a synonym of C. obscurus Sowerby, 1833.





halli da Motta, 1983.

Published in Publcoes. Occ. Soc. port. malac. (2): p. 3, figs 1 - 8.

Holotype in MHNG, $(52.8 \times 27.8 \text{ mm})$.

Type locality Pasir Putih, about 180 kms East of Surabaya, Java, Indonesia, (coral rubble).

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. hyaena* Hwass, 1792, recognised by da Motta in 1985.





hamamotoi Yoshiba & Koyama, 1984.

Published in Venus, Kyoto, 43 (2): p. 115, pl. 1, figs 1 - 5. Holotype in WPMN, $(23 \times 13.7 \text{ mm})$.

Type locality 1 to 1.5 kms West of Ogokuda Beach, Kushimoto-machi, Nishimoro-gun, Wakayama Prefecture, Japan, (dredged, 60 - 70 mtrs).

Nomenclatural status, an available name.





hamanni Fainzilber & Mienis, 1986.

Published in Bull. malac. R. of China 12 (1 - 4): p. 1, figs 1 - 3.

Holotype in HUJ, (29.4 x 13 mm).

Type locality The Creek, Jeddah, Saudi Arabia, (coarse sand and coral rubble, 35 - 40 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





hamilli Crosse, 1858.

Published in Revue Mag. Zool. 2 (10): p.122.

Lectotype of *C. piperatus* Reeve in NHMUK, (27.7 x 14.9 mm).

Type locality of *C. piperatus* not known.

Nomenclatural status, an available name, a new replacement name (nomen novum), for *C. piperatus* Reeve, 1844, unjustified as Reeve had already renamed this species *C. dillwynii* Reeve, 1849. Taxonomic status, a synonym of *C. erythraeensis* Reeve, 1843.





Lectotype of C. piperatus.

hammatus Bartsch & Rehder, 1943.

Published in Proc. Biol. Soc. Wash. 56: p.86, not figured. Holotype in USNM, $(22.5 \times 12 \text{ mm})$.

Type locality Kauai, Hawaii, (coral & sand, 179 - 253 fths). Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. circumactus* Iredale, 1929.





hanleyi Sowerby II, 1857.

Published in Thes. Conch. 3, (*Conus*) pt. 17: p. 20, no. 160, pl. 17 (203), figs 399 & 400.

Two syntypes in NHMUK, (fig. 399, 29 x 16.5 mm. & fig. 400, 16.9 \times 9.1 mm).

Type locality Mediterranean.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.









Syntype no.1 in NHMUK (fig.399)

Syntype no.2 in NHMUK (fig.400)





harasewychi Petuch, 1987.

Published in New Caribbean Molluscan Faunas: p. 31, pl. 5, figs 7 & 8.

Holotype in USNM, $(26 \times 14 \text{ mm})$.

Type locality North of Palm Beach Inlet, Palm Beach, Florida, corrected (Petuch) to Little Sal Cay Bahamas (coral rubble, 30 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. cardinalis Hwass, 1792.





harlandi Petuch, 1987.

Published in New Caribbean Molluscan Faunas: p. 74, pl. 12, figs 9 - 12.

Holotype in USNM, $(33 \times 17 \text{ mm})$.

Type locality Utila Island, Bay Islands, Honduras, (East coast), (sand & weed, 60 ft).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





havanensis Aguayo & Farfante, 1947.

Published in Revta. Soc. malac. Carlos de la Torre 5 (1): p. 11, & fig.

Holotype in MHNP, $(20.2 \times 11.3 \text{ mm})$.

Type locality Arenas de la Chorrera, off Havana, Cuba.

Nomenclatural status, an available name.



hawaiiensis "Bartsch & Rehder" Kaicher, 1956.

Published in Indo-Pacific Sea Shells: sect. 5, pl. 5, no. 6, & fig.

Specimen named whereabouts unknown, (fig. $32 \times 17 \text{ mm}$). Locality Hawaii.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or indicated.

Taxonomic status, none, (a specimen of *C. suturatus* Reeve, 1844).



Figure in Kaicher.

hayesi Korn, 2001 ("2000").

Published in Conchiglia 32 (297): p.15, pls 1 & 2.

Holotype in NMSA, $(18.7 \times 11 \text{ mm})$.

Type locality off Port Grosvenor, Transkei, South Africa, (on sandy reef in 70 - 90 mtrs).

Nomenclatural status, an available name, an invalid name, a homonym of *C. hayesi* Arnold, 1909 (fossil), renamed *C. brianhayesi* Korn, 2001.

Taxonomic status a synonym of C. brianhayesi Korn, 2001).





hedgesi Sowerby III, 1913.

Published in Ann. Mag. nat. Hist. 8 (11): p. 558, pl. 9, fig. 4. Holotype in NHMUK, $(32.8 \times 16 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. iodostoma Reeve, 1843.





<u>helenae</u> (subspecies *recurvus* Broderip) Schwengel, 1955. Published in Nautilus, Philad. 69 (1): p. 15, pl. 2, figs 14 - 15.

Holotype (Hanna), [erroneous ? lectotype], in USNM, (45.2 x 19.2 mm), lectotype representative (Lauer) figure 14, (fig. 44 x 18 mm).

Type locality Curu, Gulf of Nicoya, Costa Rica, (West Coast). Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. scalaris* Valenciennes, 1832.





helgae Blöcher, 1992.

Published in Acta Conch. (3): p. 35, pl. 3, figs 23 - 27.

Holotype in SMNS, $(36.6 \times 19 \text{ mm})$.

Type locality off Grand Récif, Tulear, S.W. Madagascar, (sand & coral rubble, 75 - 90 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species, probably not in genus *Conus* but in genus *Conorbis*, see introduction sect. F (4).





helvolus "S" Lightfoot, 1786.

Published in Cat. Portland Mus: p. 4, no. 18, not figured. Specimen named sold at auction, present whereabouts unknown, (?mm).

Locality not mentioned

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

henckesi Coltro, 2004.

Published in Strombus no. 011: p. 3, pl. 2, fig. A, pl. 9, figs H1 & P1 - 11.

Holotype in MZUSP, $(15.6 \times 8.5 \text{ mm})$.

Type locality Mar Grande, Itaparica Island, Bahia State, Brazil, (coral sand bottom at 1 - 2 mtrs on outside reefs on open sea area).

Nomenclatural status, a valid name. Taxonomic status, a valid species.





hennequini Petuch, 1993 ? ("1992").

Published in Conchiglia 24 (265): p. 10, figs 9, 10 & 11. Holotype in MNHN, $(17.5 \times 9.5 \text{ mm})$.

Type locality La Vauclin, Martinique, French West Indies, (Windward Islands), (2 mtrs).

Nomenclatural status, an available name.





henoquei Bernardi, 1860.

Published in J. Conch. Paris, 8: p. 380, pl. 13, fig. 4, (1861, Monographie de Genre Conus: p. 9, pl. 2, fig. 13).

Holotype in MNHN, $(37.5 \times 21.5 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. fumigatus* Hwass, 1792.





hepaticus Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 97, fig. 3, (1849, Coq. Viv. 2: p. 227, no. 195).

Holotype was in collection Dupont, present whereabouts unknown, $(40 \times ? \text{ mm})$, $(\text{fig. } 40 \times 23 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym (subadult) of *C. quercinus* Lightfoot, 1786.




Figure in Coq. Viv.

hereditarius da Motta, 1987.

Published in Conchiglia 19 (218 - 219): p. 27, & figs.

Holotype in MHNG, $(43.4 \times 22 \text{ mm})$.

Type locality Raya Island, Phuket, Thailand, (divers, sand & rock, $40-80~{\rm ft}$).

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. blainvillii* Vignard, 1829, (a subspecies of *C. ammiralis* Linnaeus, 1758).





herillus "Ch" (variety mediterraneus Hwass) Nardo, 1847.
Published in Sinon. Moderna, pt. 2: p. 41/42, sp. 11, fig. 520.
Type series was in collection S. Chereghin, present whereabouts unknown, (fig. ? mm).

Type locality Isle of Cherso, (Mediterranean).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.




Figure in Chiereghin MSS.

hevassii A. Adams, 1854 (" $185\overline{3}$ ").

Published in Proc. zool. Soc. Lond. pt. 21, no. 255: p. 118, no. 16.

Lectotype representative of *C. pulchellus* Sowerby I & II in Conch. Ill. Pl. 54, fig 61 (fig. 36 x 19 mm).

Type locality of C. pulchellus Freemantle, (West Australia), [Dubious].

Nomenclatural status, an available name, a new replacement name (nomen novum) for *C. pulchellus* Sowerby, 1834.

Taxonomic status, a synonym of C. varius Linnaeus, 1758.



Rep.Lect. C. pulchellus.

hieroglyphus Duclos, 1833.

Published in Magasin Zool. Paris, 3: no page number, pl. 23. Holotype in MNHN, (23 x 13 mm). (this specimen belonged to G.P. Deshayes and was loaned to Duclos who deliberately misrepresented it as measuring 36 mm. and then mentioned the Deshayes specimen as a different species, fig. $34 \times 18 \text{ mm}$).

Type locality not known, designated (Vink) Aruba, (Netherlands Antilles).

Nomenclatural status, an available name.





hilli Petuch, 1990.

Published in Nautilus, Philad. 104 (2): p. 68, figs 36 & 37. Holotype in USNM, (21 x 12 mm).

Type locality Portobelo, Panama, (East Coast), (trawled, 26 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. magellanicus* Hwass, 1792.





hirasei Kuroda, 1935.

Published in Venus, Kyoto, 5 (1): p. 54, not figured.

Specimen named destroyed in World War 2, $(62.5 \times 31.5 \text{ mm})$. Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none, (a specimen of C. hirasei Kuroda, 1956).

(Rhizoconus ?) hirasei Kuroda, 1956.

Published in Venus, Kyoto, 19 (1): p. 3, in English, p. 14, in Japanese, fig. 10.

Holotype destroyed in World War 2, (62.5 x 31.5 mm).

Type locality Kashiwajima, Tosa, Japan.

Nomenclatural status, an available name.

Taxonomic status, a valid species.




Figure in Venus

hivanus Moolenbeek, Zandbergen & Bouchet, 2008.

Published in Vita Malacologica 6: p. 30, pl. 2, figs 13 - 14 & pl.4, fig. 18.

Holotype in MNHN, $(15.7 \times 8.6 \text{ mm})$.

Type locality Marquesas Archipelago, Nuku Hiva, (90 - 120 mtrs).

Nomenclatural status, an available name.





holemani Nowell-Usticke, 1968.

Published in Caribbean Cones from St. Croix and the Lesser Antilles: p. 21, pl. 3, fig. 1011.

Lectotype (Nowell-Usticke 1971) in AMNH, (64 x 34.5 mm).

Type locality St. Thomas, (Windward Islands), & St. Vincent, (Leeward Islands).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. cedonulli Linnaeus, 1767.





honkeri Petuch, 1988.

Published in Neogene History of Tropical American Mollusks: p. 158, pl. 36, figs 1 & 2.

Holotype in USNM, $(37 \times 16 \text{ mm})$.

Type locality off Los Monges Island, Gulf of Venezuela, (trawled, 35 mtrs).

Nomenclatural status, an available name, Taxonomic status, a valid species.





hopwoodi Tomlin, 1936.

Published in J. Conch. Lond. 20 (8): p. 254.

Holotype of C. gracilis Sowerby III, 1875 in NMWC, (30.5 x 10 mm).

Type locality of *C. gracilis* not mentioned.

Nomenclatural status, an available name, a new replacement name (nomen novum) for *C. gracilis* Sowerby, 1875.

Taxonomic status, a valid species.





Holotype of *C. gracilis* Sowerby.

howelli Iredale, 1929.

Published in Rec. Aust. Mus. 17 (4): p. 182, pl. 40, figs 1 & 8. Holotype in AMS, $(27 \times 13.3 \text{ mm})$.

Type locality Montague Isl., N.S.W., Australia, (trawled, 50 - 60 fths).

Nomenclatural status, an available name.





huberorum da Motta, 1989.

Published in Conchiglia 21 (242 - 245): p. 9, & figs.

Holotype in MHNG, $(48.2 \times 24.7 \text{ mm})$.

Type locality Coromandel coast, Bay of Bengal, North of Madras, (India), (20 - 40 fths).

Nomenclatural status, an available name.

Taxonomic status, a synonym, (teratological form) of *C. loroisii* Kiener, 1845 (1846) (, recognised later by da Motta.





humilis von Salis Marschlins, 1793.

Published in Reisen Provinzen Königsreichs, Neapel: App. p. 364, not figured.

Type material whereabouts unknown.

Type locality Taranto and Naples, (Italy).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.

humilis "Smith" Galindo, 1977.

Published in Index & Register of Seashells: p. 115, not figured. Specimen named whereabouts unknown, (? mm).

Locality Japan.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none.

hunti Wils & Moolenbeek, 1979.


Published in Bijdr. Dierk. 49 (2): p. 257, figs 7 - 9 & 11 - 14. Holotype in NBC-ZMA, (18.6 x 9.9 mm).

Type locality off St. James, West of Barbados, Windward Islands, (155 - 180 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. sanderi* Wils & Moolenbeek, 1979, p.255, recognised later by both authors.





<u>hyaena</u> Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 656, no. 55, (1798, Tab. Enc. pl. 327, figs 5 & 7).

Lectotype (Kohn) in MHNG, (61 \times 33.5 mm). Type locality West Africa, [erroneous]. Nomenclatural status, an available name.





hybridus Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 83, fig. 1, (1850, Coq. Viv. 2: p. 256, no. 223).

Holotype was in collection J.P.B. Delessert, present whereabouts unknown, $(42 \times ? mm)$, $(fig. 42 \times 22 mm)$.

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a valid species.



Figure in Coq. Viv.

hypochlorus Tomlin, 1937.

Published in Proc. malac. Soc. Lond. 22 (4): p. 206.

Holotype of C. croceus Smith in NHMUK, (25.3 x 8.5 mm).

Type locality of C. croceus not known.

Nomenclatural status, an available name, a new replacement name (nomen novum) for *C. croceus* Smith, 1877.





Holotype of C. croceus

iansa Petuch, 1979.

Published in Proc. biol. Soc. Wash. 92 (3): p. 524, figs 4g & h. Holotype in USNM, $(12 \times 7 \text{ mm})$.

Type locality Abrolhos Arch., 2 kms East of Sta. Barbara Isl., Bahia State, Brazil, (mud, 25 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. mindanus Hwass, 1792.





iberogermanicus Röckel, Rolán & Monteiro, 1980.

Published in Cone Shells from Cape Verde Islands: p. 75, figs 48 - 50 & pl. 3, row 5.

Holotype in NMSF, $(32 \times 17.5 \text{ mm})$.

Type locality Sal-rei, Boavista Isl., Cape Verde Islands, (under rocks, $0-3\ \mathrm{mtrs}$).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. irregularis* Sowerby, 1858.





ichinoseana Kuroda, 1955.

Published in Venus, Kyoto, 18 (4): p. 293, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name, (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none, (a specimen of *C. ichinoseana* Kuroda, 1956).

Asprella (Conasprella ?) <u>ichinoseana</u> Kuroda, 1956.

Published in Venus, Kyoto, 19 (1): p. 10, in English, p. 15, in Japanese, pl. 1, fig. 5.

Holotype in KIMN, $(58 \times 21 \text{ mm})$.

Type locality off Tosa, Japan.

Nomenclatural status, an available name.





ignobilis Olivi, 1792.


Published in Zoologia Adriatica, Ossia Catalogo Ragionata degli Animali del Golfo e Lagune di Venezia & Bassano, Venice: p. 133, not figured.

Lectotype representative (Kohn) figure in Gualtieri, 1742, pl. 20, fig. P. (fig. $26 \times 13 \text{ mm}$).

Type locality Ischia, Adriatic Sea.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.



ignotus Cargile, 1998.

Published in Siratus 2 (14): p. 9, figs 1 - 4.

Holotype in SBMNH, $(20.6 \times 10.5 \text{ mm})$.

Type locality West side of Quito Sueno Bank, (off Nicaragua, East coast), (25 - 35 mtrs)

Nomenclatural status, an available name.





(Lithoconus) iheringi Frenquelli, 1946.

Published in Notas Mus. La Plata 11 (88): p. 234, pl. 1, figs 1 - 8 & pl. 2, figs 4 - 7.

A syntype (non fossil) in MBA, (42 x 22 mm), a syntype (fossil) in MACN, (32 x 18 mm) and a syntype (fossil) in MUP, (47 x 26 mm).

Type locality Cape San Antonio, Buenos Aires, Argentina. Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. carcellesi* Martins, 1945, (a




Figure in Frenguelli, 1946

(Leptoconus) <u>ikatt</u> (forma thalassiarchus Sowerby) Shikama, 1979. Published in Sci. Rep. Yokosuka Cy. Mus. (26): p. 4, pl. 1, figs 11 & 12.

Specimen described in NSMT, $(58.7 \times 31.3 \text{ mm})$, [missing in May 1989 & April 1995, a second specimen is present $(65 \times 32.8 \text{ mm})$]. Locality Zamboanga, Mindanao Island, Philippines.

Nomenclatural status, an unavailable name (nomen nudum), described as a form after 1960.

Taxonomic status, none, (a specimen of C. thalassiarchus Sowerby, 1834).





Second specimen in NSMT.

(Profundiconus) ikedai Ninomiya, 1987.

Published in Venus, Kyoto, 46 (1): p. 7, pl. 1, figs 1 - 4 & pl. 2, figs 5 & 6.

Holotype in NSMT, $(25.5 \times 11.2 \text{ mm})$.

Type locality S.W. off Jogashima Islet, Miura Peninsula, Japan, (dredged, 20 - 280 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





(Leptoconus) illawarra Garrard, 1961.

Published in \overline{J} . malac. Soc. Aust. 1 (5): p. 31, pl. 1, fig. 2. Holotype in AMS, (29 x 15 mm).

Type locality East of Stanwell Park, N.S.W., Australia, (trawled, 75 fths).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. sydneyensis Sowerby, 1887.





<u>imelus</u> (variety mediterraneus Hwass) de Gregorio, 1885. Published in Boll. Soc. malac. ital. 11: p. 95, not figured. Type series in MPUP but no longer identifiable, (? mm). Type locality Mediterranean, (in Sponges).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.

immaculata (variety betulinus Linnaeus) Dautzenberg, 1906.

Published in J. Conch. Paris, 54: p. 27, not figured.

Holotype in IRSN, $(130 \times 81 \text{ mm})$.

Type locality Ambodifoutra, coast East of Ste Marie, (Island), Madagascar.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. betulinus* Linnaeus, 1758.





immelmani Korn, 1998.

Published in Conchiglia 30 (288): p. 11 & figs.

Holotype in NMSA, (89.7 \times 46 mm) not, as originally reported, in NMC.

Type locality Southern Natal (holotype) & Northern Transkei (paratypes), (South Africa), (alongside coral reef at the base of narrow ledges, 35 - 50 mtrs).

Nomenclatural status, an available name.





(Virroconus) imperator Woolacott, 1956 ("1954 - 55").
Published in Proc. R. zool. Soc. N.S.W. (no number): p. 72, fig.

Holotype in AMS, $(42.1 \times 23.6 \text{ mm. orig. desc. } 40 \times 25 \text{ mm})$. Type locality Trinity Bay, Queensland, Australia.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. parvulus* Link, 1807, (a synonym of *C. biliosus* (Röding), 1798).



imperialis Linnaeus, 1758.

Published in Syst. Nat. 10th ed. (1): p. 712, no. 251, not figured.

Lectotype (Kohn) in LSL, $(65 \times 37 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.





imperialis (Röding), 1798.

Published in Mus. Bolten 2: p. 45, no.575/-, not figured.

Specimen named whereabouts unknown, (? mm).

Type locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

inaequalis Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 7, sp. 270 (320). Holotype in NHMUK, (17 x 9 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.





incarnatus Reeve, 1844.

Published in Conch. Icon. 1 (*Conus*): pl. 41, sp. 221, (published Jan), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 174, not figured, published Jun).

Three syntypes in NHMUK, (figured syntype 47.00×26.00 mm, syntype $2.65.20 \times 34.30$ mm & syntype $3.47.60 \times 26.30$ mm. Type locality Malacca, (Malaysia), (mud banks).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. hyaena Hwass, 1792.









Figured syntype in NHMUK. ©NHMUK.

Syntype no. 2 in NHMUK. ©NMHUK.

incinctus Fenaux, 1942.

Published in Bull. Inst. oceanogr. Monaco, (814): p. 3, fig. 6. Type series whereabouts unknown, (fig. 27 x 12.5 mm).

Type locality Australia.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. anemone Lamarck, 1810.



Figure in Bull. Inst.

inconstans E.A. Smith, 1877.

Published in Ann. Mag. nat. Hist. (4) 19: p. 224, not figured.

Lectotype (Vink) in NHMUK, (22.4 x 12.3 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





incurvus "Broderip" Sowerby I & II, 1833.

Published in Conch. Ill. (Conus): pl. 33, fig. 36.

Lectotype representative (Kohn) figure in C. I. pl. 33, fig. 36, (fig. $42 \times 18 \text{ mm}$).

Type locality Monte Christe, (Manta, Ecuador).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. regularis Sowerby, 1833).



<u>indiae</u> (subspecies archithalassus Hwass) (Röding), 1798. Published in Mus. Bolten 2: p. 43, no. 545/57, not figured. Lectotype representative (Kohn) figures in Chemnitz, 1788, pl. 138, figs 1282 & 1283. (figs 77 x 47 mm).

Type locality not mentioned, (Chemnitz East Indies), [erroneous]. Nomenclatural status, an available name.

Taxonomic status, a synonym of C. pulcher Lightfoot, 1786.



indicus "Chemnitz" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 422, no. 136, not figured.

Specimen named whereabouts unknown, cited figure in Chemnitz, 1788, pl. 140, fig. 1295, (fig. $56 \times 27 \text{ mm}$).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of $C.\ magus$ Linnaeus, 1758.

Taxonomic status, none, (a specimen of C. magus Linnaeus, 1758).



<u>indicus</u> (variety *magus* Linnaeus) "Chemnitz" Küster, 1838.

Published in Mollusca Gastropoda Conea (Kegelschnecken): pl. 6, fig. 1.

Type series whereabouts unknown, (fig. $55 \times 27 \text{ mm}$), cited figure in Chemnitz, 1788, pl. 140, fig. 1295, ($57 \times 28 \text{ mm}$).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. magus Linnaeus, 1758.



Figure in Küster.

indicus "Chemnitz" Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Emend. p. 2, not figured. Specimen named whereabouts unknown, cited figure in Chemnitz, 1788, pl. 140, fig. 1295, (fig. 57 x 28 mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. magus* Linnaeus, 1758.

Taxonomic status, none, (a specimen of C. magus Linnaeus, 1758).



Cited figure in Chemnitz.

"indicus" (variety inscriptus Reeve) Röckel, 1979.

Published in Hawaii. Shell News 27 (3): p. 11, fig. 2.

Two specimens named in collection D. Röckel now in SMNS but no longer identifiable, (figs $41 \times 21 \& 25 \times 16 \text{ mm}$).

Locality South India & Sri Lanka.

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name, and publication invalid by disclaimer.

Taxonomic status, none, (specimens are a form of C. inscriptus Reeve, 1843).









Specimens in SMNS.

induratus Reeve, 1849.

Published in Conch. Icon. 1 (*Conus*): Sup. pl. 7, sp. 268 (318). Lectotype representative (Röckel & Moolenbeek) figure in C. I. pl. 7, sp. 268 (318), (fig. 21 x 10.5 mm).

Type locality Red Sea.

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. erythraeensis* Reeve, 1843.



infinitus Rolán, 1990.

Published in Iberus, Sup. 2: p. 39, pl. 1, fig. 12, pl. 2, fig. 12 & pl. 5, fig. P.

Holotype in MNCM, $(20.8 \times 12 \text{ mm})$.

Type locality Pau Seco Bay, Maio Island, Cape Verde Islands, (under rocks, 1 - 2 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





<u>inflatus</u> Sowerby I & II, 1833.

Published in Conch. Ill. (Conus): pl. 33, fig. 41.

Type series whereabouts unknown, (fig. 38 x 21 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium).



Figure in Conch. Ill.

informis Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 699, no. 88, (1798, Tab. Enc. pl. 337, fig. 8).

Lectotype (Kohn) in MHNG, $(55.5 \times 25.5 \text{ mm})$.

Type locality America, [erroneous].

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. mozambicus* Hwass, 1792 (p.696, no.85).





informis (Hwass) Reeve, 1843.

Published in Conch. Icon. 1 (Conus): pl. 5, sp. 24.

Four syntypes in NHMUK, figured syntype 49.75×23.14 mm, syntype $2~46.55 \times 23.00$ mm, syntype $3~38.60 \times 19.60$ mm & syntype $4~19.30 \times 10.70$ mm.

Type locality not mentioned.

Nomenclatural status, an available name, a homonym of *C. informis* Hwass, 1792. (Reeve explains that his species is not the same as that of Hwass although he lists it under Hwass's name).

Taxomonic status, a senior objective synonym of $C.\ simplex$ Sowerby, 1858, same type.



Figure in Conch Icon.



The second secon

Figured syntype in NHMUK. ©NHMUK.

infrenatus Reeve, 1848.

Published in Conch. Icon. 1 (Conus): Sup. pl. 3, sp. 285.

Holotype in NHMUK, $(26 \times 13.8 \text{ mm})$.

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





infumata (variety achatinus Hwass) Dautzenberg, 1937.

Published in Mem. Mus. r. Hist. nat. Belg. 2 (18): p. 12, not figured.

Lectotype representative (Walls) figure in Chemnitz, 1788, pl. 142, fig. 1320, (fig. 60 \times 31 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. achatinus Hwass, 1792.



innexus A. Adams, 1854 ("1853").


Published in Proc. zool. Soc. Lond. pt. 21, no. 255: p. 118, no. 17, not figured.

Three syntypes in NHMUK, (no. 1, 58.6 x 26.5, no. 2,50.6 x 23.8 & no.; 3, 48.1 x 21.4 mm).

Type locality Natal, (South Africa), (Cuming), [dubious]. Nomenclatural status, an available name.

Taxonomic status, a synonym of C. consors Sowerby, 1833.







Syntype no.2 in NHMUK. Syntype no.3 in NHMUK.



innotabilis E.A. Smith, 1892.

Published in Proc. zool. Soc. Lond. unnumbered (60), pt. 3: p. 487, pl. 40, fig. 1.

Holotype in NHMUK, $(17.4 \times 9.6 \text{ mm. orig. desc. } 18 \times 10 \text{ mm})$. Type locality Rocky Point, 4 miles North of Richmond River also Port Stephens, New South Wales, (Australia), (Brazier). Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of C. cyanostoma A. Adams, 1854.





inornatus Mawe, 1823.

Published in The Linnean System of Conchology: p. 89, not figured.

Specimen named whereabouts unknown, (? mm).

Locality Sumatra, (Indonesia).

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

inquinatus Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 5, sp. 251 (301). Type series was in collection H. Cuming, present whereabouts unknown, (fig. 50 x 26 mm).

Type locality West of Africa.

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. ermineus Born, 1778.



Figure in Conch Icon

inscriptus Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 29, sp. 164, (published Nov), (1844, Proc. zool. Soc. Lond. pt. 11, no. 130: p. 171, not figured, published Jun).

Three syntypes were in collection H. Cuming, in NHMUK type collection but cannot be traced at present, $(34.5 \times 18 \text{ mm.})$ figured, $32.5 \times 17.0 \text{ mm.}$ & $29.0 \times 14.0 \text{ mm}$).

Type locality not known.

Nomenclatural status, an available name.

Taxonomical status, a valid species.



Figure in Reeve.

insculptus Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 99, fig. 2, (1850, Coq. Viv. 2: p. 309, no. 275).

Type series was in collection Largilliert, present whereabouts unknown, (25 x ? mm), (fig. 25 x 10 mm), a probable syntype is in MNHN, (21.5 x 8.5 mm).

Type locality China Sea.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





Probable syntype in MNHN.

insignis Sowerby I & II, 1833.

Published in Conch. Ill. (Conus): pl. 28, fig. 17.

Lectotype representative (Kohn) figure in C. I. pl. 28, fig. 17, $(fig. 38 \times 21 \text{ mm})$.

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. acuminatus* Hwass, 1792, recognised by Sowerby in Index, 1841, p. 1, no. 17, (a synonym of *C. locumtenens* Blumenbach, 1791).



<u>insignis</u> (variety *figulinus* Linnaeus) Dautzenberg, 1937. Published in Mem. Mus. r. Hist. nat. Belg. 2 (18): p. 108, pl. 1, fig. 6.

Holotype in IRSN, $(75 \times 49 \text{ mm})$.

Type locality Amboine, (Ambon, Indonesia).

Nomenclatural status, an available name, an invalid name, a homonym of *C. insignis* Sowerby, 1833.

Taxonomic status, a synonym (form) of *C. loroisii* Kiener, 1845 (1846).





<u>insolitus</u> (form mediterraneus Hwass) "Monterosato" Gaglini in Settepassi, 1985 ("1972").

Published in Atl. Malac. Moll. Marin. 3: p. 8, not figured. Specimen described in collection T. di I. Monterosato in MCR but no longer identifiable, (37 \times 20 mm).

Locality unknown.

Nomenclatural status, an unavailable name (nomen nudum), described as a form after 1960.

Taxonomic status, none, (a specimen of C. ventricosus Gmelin, 1791).

insularis Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3389, no. 38, not figured.

Holotype was in collection J.F. Bolten acquired by F.C. Schmidt, probably in MNG, destroyed in World War 2, holotype

representative (Kohn) figure in Martini, 1773, pl. 62, fig. 683. (fig. 43 \times 24 mm).

Type locality not known, designated (Vink & von Cosel) Gros Islet Bay, St. Lucia, Lesser Antilles.

Nomenclatural status, an available name.

Taxonomic status, a subspecies of C. cedonulli Linnaeus, 1767.



inter "Jousseaume" O. & J.B. Lozet, 1978.

Published in Le Peigne de Venus, Coquillages du Monde Entier: p. 103, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none, (stated to be a specimen of *C. crocea* Broderip, 1833 (p. 99), as Broderip did not introduce this name, presumably an error for *C. croceus* Sowerby I & II, 1833, (a synonym of *C. daucus* Hwass, 1792).

intermedius Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 23, sp. 129, (published Aug), (1844, Proc. zool. Soc. Lond. pt. 11, no. 130: p. 169, not figured, published Jun).

Holotype of C. rosea Sowerby I & II in NHMUK, (53.5 x 22 mm). Type locality of C. rosea Annaa (Anaa Isl., Tuamotu Archipelago), (on reefs).

Nomenclatural status, an available name, a new replacement name (nomen novum) for *C. rosea* Sowerby I & II, 1833 and Broderip, 1833, a homonym of *C. intermedius* Lamarck, 1810 (fossil), renamed *C. mappa* Crosse, 1858.

Taxonomic status, a synonym of C. eldredi Morrison, 1955.





Holotype of C. rosea Sowerby in NHMUK.

<u>interrupta</u> "Stalio MSS" (variety mediterraneus Hwass) Coen, 1933. Published in Memorie R. Comit. talassogr. ital. (192): p. 70 & p. 175, no. 145, not figured.

Type series was in collection Stallio, probably in MCV, (? mm). Type locality Tremiti, Adriatic.

Nomenclatural status, an available name, an invalid name, a homonym of C. interruptus Wood, 1828 and of Broderip & Sowerby, 1829.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.

interruptus Wood, 1828.

Published in Index Testaceologicus, 2nd. ed: Sup. p. 8, pl. 3, fig. 2.

Holotype was in collection F.J. Stainforth, present whereabouts unknown, (app. 51 mm), holotype representative (Kohn) figure in Wood, pl. 3, fig. 2. (fig. 15 \times 9 mm).

Type locality not known (Galapagos, Reeve).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. brunneus Wood, 1828.



interruptus Broderip & Sowerby I, 1829.

identifiable, $(46 \times 20 \text{ mm})$.

Published in Zool. J. Lond. 4: p. 379, not figured. Holotype was in ZSML, transferred to NHMUK but no longer

Type locality near Mazatlán, (Mexico, West coast).

Nomenclatural status, an available name, an invalid name, a homonym of *C. interruptus* Wood, 1828.

Taxonomic status, a synonym of C. ximenes Gray, 1829.

interstinctus Guppy, 1866.

Published in Q. Jl. geol. Soc. Lond, (22): p.288, pl. 16, fig. 3. Type series whereabouts unknown, $(fig. 57 \times 27.5 \text{ mm})$.

Type locality Jamaica.

Nomenclatural status, an available name.

Taxonomic status, a synonym (possibly a fossil form) of C. granarius Kiener, 1848 (1847), (a subspecies of C. mappa Lightfoot, 1786).



Figure in Guppy.

iodostoma Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 28, sp. 159, (published Oct), (1844, Proc. zool. Soc. Lond. pt. 11, no. 130: p. 170, not figured, published Jun).

Four syntypes in NHMUK, (42 x 19.5 mm. figured, 47 x 22, 42 x 19 & 42 x 19 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





Figured syntype in NHMUK.

ione Kuroda, 1935.

Published in Venus, Kyoto, 5 (1): p. 54, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured.

Taxonomic status, none, (a specimen of *C. ione* Fulton, 1938).

ione Fulton, 1938.

Published in Proc. malac. Soc. Lond. 23 (1): p. 55, pl.3, fig. 2. Holotype in NHMUK, $(58 \times 28.4 \text{ mm})$.

Type locality Kii, Japan.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





<u>irregularis</u> Sowerby II, 1858.

Published in Thes. Conch. 3 (Conus), pt. 18: p. 29, no. 242, pl. 18 (204), figs 418 - 419.

Three syntypes in NHMUK, (fig. 418, $28.3 \times 18.1 \text{ mm}$, no. 2, $28 \times 18.1 \text{ mm}$ 17.7 & no. 3, 31.8 \times 19.3 mm).


Type locality West Africa.


Nomenclatural status, an available name.

Taxonomic status, a valid species.



Figured syntype in NHMUK. Syntype no. 2 in NHMUK.





Syntype no. 3 in NHMUK.

isabelarum Tenorio & Afonso, 2004.

Published in Visaya, vol. 1, no. 2: p. 29, pl. 4.

Holotype in MNCM, $(22.8 \times 13.5 \text{ mm})$.

Type locality Baia de Pau Seco, West coast of Maio Island, Cape Verde Islands, (under rocks and dead coral slabs on sandy bottom, 2-4 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





isabella (Röding), 1798.

Published in Mus. Bolten 2: p. 44, no. 552/63, not figured.

Specimen named whereabouts unknown, (? mm).

Type locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none.

<u>istriensis</u> "Ch" (variety mediterraneus Hwass) Nardo, 1847. Published in Sinon. Moderna, pt. 2: p. 41/42, sp. 9, figs 517 - 518.

Type series was in collection S. Chereghin, present whereabouts unknown, (fig. ? mm).

Type locality Istria, Mediterranean.

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. ventricosus* Gmelin, 1791.




Figure in Chiereghin MSS.