

# Section V vanhyningi to vulpinus

Copyright © 2011.R Filmer Conditions of use.

The content of this website is provided for personal and scientific use and may be downloaded for this purpose. It may not be used in whole or part for any commercial activity and publication of any of the content on the internet is limited to the Cone Collector website (<a href="www.TheConeCollector.com">www.TheConeCollector.com</a>). Authors wishing to publish any of the pictures may do so on a limited basis but should inform M Filmer so that the original owner of the rights to the picture can be acknowledged. (mike@mfilmer.fsnet.co.uk)

Version 1.0 October 2011.

vanhyningi (subspecies verrucosus Hwass) Rehder, 1944.

Published in Nautilus, Philad. 47 (3): p. 105, not figured. Holotype in USNM, (17 x 10.4 mm).

Type locality off Pompano Beach, Broward County, Florida, (beached).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of *C. mindanus* Hwass, 1792.





vanikenensis Galindo, 1977.

Published in Index & Register of Seashells: p. 123, not figured. Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described, indicated or figured. Taxonomic status, none.

moluccensis vappereaui Monteiro, 2009.

Published in Visaya vol.2, no.5: p. 80 & figs.

Holotype in MNCM,  $(47.7 \times 26.7 \text{ mm})$ .

Type locality Arue, Tahiti Island, Society Islands, French Polynesia, (60 - 70 mtrs, on the outer slope of the reef barrier, on sand bottom in caves).

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. moluccensis* Küster, 1838.





variegatus Kiener, 1845 (1848).

Published in Coq. Viv. 2: pl. 106, figs 1 & 1a, (1849 (1850), Coq. Viv. 2: p. 261, no. 228).

Type series was in collection Largilliert, present whereabouts unknown, (25 x ? mm), lectotype representative (Röckel & Fernandes) figure in C. V. pl. 106, fig. 1a, (fig. 18 x 11 mm). Type locality not mentioned, designated (Rolán & Röckel) Santa Maria Bay, Angola.

Nomenclatural status, an available name. Taxonomic status, a valid species.



varius Linnaeus, 1758.

Published in Syst. Nat. 10th ed. (1): p. 715, no. 270, not figured.

Lectotype (Kohn), in LSL,  $(33.5 \times 16 \text{ mm})$ .

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





<u>vaubani</u> Röckel & Moolenbeek in Röckel, Richard & Moolenbeek, 1995.

Published in Mem. Mus. natn. Hist. nat. Paris, no. 167, 14 (9): p. 587, figs 3, 14, 15 & 56.

Holotype in MNHN,  $(25.8 \times 11.7 \text{ mm})$ .

Type locality South New Caledonia, (435 mtrs).

Nomenclatural status, an available name.





vautieri Kiener, 1845 (1847).

Published in Coq. Viv. 2: pl. 100, fig. 3, (1850, Coq. Viv. 2: Sup. p. 350, no. 313).

Holotype was in collection A.C. Bernardi acquired by M. Gubba then in MHNH, destroyed in World War 2,  $(32 \times ? mm)$ ,  $(fig. 32 \times 18 mm)$ .

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. pulicarius* Hwass, 1792.



Figures in Coq. Viv.

### vayssetianus Crosse, 1872.


Published in J. Conch. Paris, 20: p. 154 & p. 349, pl. 16, fig. 1.

Type series was in collection Rossiter, present whereabouts unknown, (14  $\times$  7.5 mm).

Type locality New Caledonia, (Rossiter).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. exiguus Lamarck, 1810.



Figures in Crosse

vayssierei (subspecies mediterraneus Hwass) Pallary, 1906.

Published in Ann. Soc. Linn. de Lyon and in J. Conch. Paris, 54 (2): p. 78, pl. 4, fig. 1.

Two syntypes in MNHN,  $(28.5 \times 16.5 \& 28 \times 13 \text{ mm})$ .

Type locality Houmt Souk Adjim, El Kantara, Gulf of Gabes, (Tunisia).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.





Syntype in MNHN.

veillardi da Motta, 1990.

Published in Conchiglia 22 (253 - 255): p. 44 & figs.

Holotype in MHNG,  $(52.4 \times 25.3 \text{ mm})$ .

Type locality Glorieuses Islands, western Indian Ocean, (lagoons, near rocks, 5 - 7 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of C. gubernator Hwass, 1792.





**velaensis** Petuch, 1993 ? ("1992").

Published in Conchiglia 24 (265): p. 14, figs 18 & 19.

Holotype in CMNH,  $(31 \times 15 \text{ mm})$ .

Type locality North of Cape La Vela, Goajira Peninsula, Colombia, (East coast), (35 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





venezuelanus Petuch, 1987.

Published in New Caribbean Molluscan Faunas: p. 113, pl. 25, figs 18 & 19.

Holotype in USNM,  $(25 \times 12 \text{ mm}, \text{ orig. desc. } 27 \text{ mm})$ .

Type locality off Puerto Cabello, Gulf of Triste, Venezuela, (trawled, 25 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species or possibly a synonym (form) of *C. cancellatus* Hwass, 1792.





ventricosus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3397, no. 71, not figured.

Lectotype representative (Kohn) figure in Kammerer, 1786, pl. 6, fig. 3, (fig.  $59 \times 32 \text{ mm}$ ).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a valid species.



ventricosus (Röding), 1798.

Published in Mus. Bolten 2: p. 49, no. 623/114, not figured. Lectotype representative (Kohn) figure in Chemnitz, 1788, pl. 142, fig. 1317, (fig. 80 x 45 mm).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. ventricosus* Gmelin, 1791.

Taxonomic status, an objective junior synonym of *C. achatinus* Gmelin, 1791, same type.



ventricosus
(form mediterraneus Hwass) "Auctorum" Gaglini in
Settepassi, 1985 ("1972").

Published in Atl. Malac. Moll. Marin. 3: p. 10, not figured. Specimen described in collection T. di I. Monterosato in MCR but no longer identifiable, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), described as a form after 1960.

Taxonomic status, none, (a specimen of C. ventricosus Gmelin, 1791).

venulatus Hwass in Bruquière, 1792.

Published in Enc. Méth. 1: p. 695, no. 84, (1798, Tab. Enc. pl. 337, fig. 9).

Lectotype (Kohn) in MHNG, (43 x 28.5 mm).

Type locality America, [erroneous].

Nomenclatural status, an available name.

Taxonomic status, a valid species.





venustus (Röding), 1798.

Published in Mus. Bolten 2: p. 47, no. 599/94, not figured.

Holotype whereabouts unknown, (? mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of  $C.\ amadis$  Gmelin, 1791, as indicated by Röding.

verdensis Trovão, 1979.

Published in Amphitrite 1 (1): p. 5, pl. 1, fig. 3. Holotype in NHMUK, (17.1 x 11.1 mm).

Type locality Santa Luzia Isl., Cape Verde Islands, (12 mtrs). Nomenclatural status, an available name. Taxonomic status, a valid species.





"verhoefi" "Liltved" Röckel & Moolenbeek, 1996.

Published in Vita Marina 44 (1 - 2): p. 47, pl. 1, figs 5 & 6.

Specimen described in MNHN, (27.6 x 10.9 mm).

Locality Pitcairn Islands, (South Pacific), (Liltved).

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name, Liltved never published this name although he presumably had intended to do so.

Taxonomic status, none, (the specimen is labelled "paratype no. 2" of C. tirardi Röckel & Moolenbeek, 1996).

vermiculatus Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 34, no. 17, not figured.

Holotype was in MNHN, present whereabouts unknown, (33 x 22 mm), cited figures in Tab. Enc. 1798, pl. 321, figs 1 & 8, (fig. 1, 33 x 22 & fig. 8, 36 x 23 mm).

Type locality Asia, Africa and America, (warm seas).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. chaldaeus (Röding), 1798.





Cited figures in Tab. Enc.

verreauxii Kiener, 1845 (1846).

Published in Coq. Viv. 2: pl. 60, fig. 5, (1849 (1850), Coq. Viv. 2: p. 249, no. 217).

Holotype was in collection Verreaux, present whereabouts unknown,  $(40 \times ? \text{ mm})$ ,  $(fig. 40 \times 21 \text{ mm})$ .

Type locality coasts of the Cape of Good Hope, (South Africa), corrected (Filmer) to Nusa Tengara Indonesia.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. spectrum* Linnaeus, 1758.



Fig. in Coq. Viv.

verriculum Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 38, sp. 208a & b, (published Dec), (1844, Proc. zool. Soc. Lond. pt. 11, no. 130: p. 173, not figured, (published Jun).

Two syntypes were in collection F.J. Stainforth, present whereabouts unknown, (54 x 29 & 51 x 29 mm), provisional lectotype representative (Lauer) figure in C. I. pl. 38, sp. 208b, (fig.  $54 \times 29 \text{ mm}$ ).

Type locality Ceylon etc., (Sri Lanka).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. textile Linnaeus, 1758.



verrucosus Hwass in Bruquière, 1792.

Published in Enc. Méth. 1: p. 708, no. 100, (1798, Tab. Enc. pl. 333, fig. 4).

Type figure designated (Clench) figure in Martini, 1773, pl. 55, fig. 612 C, (fig. 21 x 11 mm), lectotype representative (Kohn) figure in Tab. Enc. pl. 333, fig. 4,  $(23 \times 11 \text{ mm})$ .

Type locality Senegal & Mozambique, (both erroneous), corrected (Clench) Puerto Plata, Santo Domingo, (Dominican Republic). Nomenclatural status, an available name.

Taxonomic status, a synonym (granulose form) of *C. jaspideus* Gmelin, 1791.



Figure in Martini.



Figure in Tab. Enc.

versicolor (variety ignobilis Olivi) Scacchi, 1836.

Published in Catalogus Conchyliorum Regni, Neapolitani: p. 9, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not known.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none, (a specimen of *C. ventricosus* Gmelin, 1791).

verulosus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 719, no. 115, (1798, Tab. Enc. pl. 341, fig. 7).

Lectotype representative (Kohn) figure in Tab. Enc. pl. 341, fig. 7, (fig. 45  $\times$  23 mm, orig. desc. 45  $\times$  20 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. granulatus Linnaeus, 1758.



vespertinus Sowerby I, 1825.

Published in Cat. Shells Tankerville: p. 91, no. 2446, pl. 8, figs 3 & 4.

Holotype in NMWC,  $(48 \times 21 \text{ mm})$ .

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. timorensis Hwass, 1792.






vexillum Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3397, no. 68, not figured.

Lectotype representative (Kohn) figure in Rumphius, 1705, pl. 31, fig. S (error for fig. V, (fig.  $68 \times 39 \text{ mm}$ ).

Type locality not known, (Rumphius indicates Amboina, (Ambon, Indonesia).

Nomenclatural status, an available name.



<u>vezoi</u> (subspecies *pennaceus* Born) Korn, Niederhőfer & Blőcher, 2000.

Published in Stuttg. Beitr. Naturk. ser. A, no. 610, 25 S: p.20. pl. 2, figs 1 - 9.

Holotype in SMNS,  $(50.1 \times 26.7 \text{ mm})$ .

Type locality Maromitiliky 25 kms airline South of Pointe Beheloka, South of Toliara [Tulear], South West Madagascar, (subtidal inside shallow lagoon and near outside reef edge). Nomenclatural status, an available name.

Taxonomical status, a synonym (form) of C. pennaceus Born, 1778.





### viaderi Fenaux, 1942.

Published in Bull. Inst. oceanogr. Monaco, (814): p. 3, fig. 9. Type series whereabouts unknown, (fig. 30 x 13 mm).

Type locality Mauritius, [erroneous].

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. locumtenens Blumenbach, 1791.



Figure in Bull. Inst.

vicarius Linnaeus, 1767.

Published in Syst. Nat. 12th ed. (1): p. 1167, no. 299, not figured.

Lectotype representative (Kohn) figure in d'Argenville, 1742, pl. 15, fig. H, (fig.  $55 \times 25 \text{ mm}$ ).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ammiralis Linnaeus, 1758.



<u>vicarius</u> (variety ammiralis Linnaeus) Hwass in Bruguière, 1792. Published in Enc. Méth. 1: p. 659, no. 57E, (1798, Tab. Enc. pl. 328, fig. 2).

Type series whereabouts unknown, (fig.  $71 \times 37 \text{ mm}$ ).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. vicarius* Linnaeus, 1767.

Taxonomic status, a synonym of C. ammiralis Linnaeus, 1758.



Figure in Tab. Enc.

vicarius (Röding), 1798.

Published in Mus. Bolten 2: p. 45, no. 567/76 & 568/-, not figured.

Lectotype representative (Kohn) figure in Knorr, 1771, pl. 24, fig. 4, (fig.  $33 \times 16 \text{ mm}$ ).

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. vicarius* Linnaeus, 1767 and of Hwass, 1792. Taxonomic status, a synonym of *C. locumtenens* Blumenbach, 1791.



vicarius Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 274, no. 85, not figured.

Holotype in MNHN,  $(45 \times 24 \text{ mm})$ .

Type locality Asian Seas, uncertain.

Nomenclatural status, an available name, an invalid name, a homonym of *C. vicarius* Linnaeus, 1767, of Hwass, 1792 and of (Röding), 1798, renamed *C. abbreviata* Dautzenberg, 1937. Taxonomic status, a synonym of *C. ammiralis* Linnaeus, 1758.





vicarius (variety ammiralis australis Lamarck) Küster, 1837.

Published in Syst. Conch.-Cab. 4 (1): p. 26, no. 14, Var. E, not figured.

Specimen described whereabouts unknown, (? mm), cited figure in Tab. Enc. 1798, pl. 328, fig. 6, (fig. 66 x 31 mm). Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), described as a variety of a variety.

Taxonomic status, none, (a specimen of C. ammiralis Linnaeus, 1758).



Cited figure in Tab. Enc.

<u>vicarius</u> (variety ammiralis archithalassus Hwass) Küster, 1837. Published in Syst. Conch.-Cab. 4 (1), p. 26, no. 14, Var. E, not figured.

Specimen described whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), described as a variety of a variety.

Taxonomic status, none, (a specimen of *C. ammiralis* Linnaeus, 1758).

(Rhizoconus) <u>vicdani</u> (subspecies sugimotonis Kuroda) Lan, 1978. Published in Bull. malac. R. of China (5): p. 66, pl. 1, figs 1 - 6.

Holotype in MHNG,  $(78.7 \times 43.8 \text{ mm})$ .

Type locality Panlao [Panglao], Bohol, Philippines, (120 fths). Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. sugimotonis* Kuroda, 1928.





vicdani Kosuge, 1980.

Published in Bull. Inst. malac. Tokyo, 1 (4): p. 62, pl. 18, figs 1 - 5.

Holotype in IMT,  $(47.9 \times 23.8 \text{ mm})$ .

Type locality Balut Island, Davao, Philippines.

Nomenclatural status, an available name, an invalid name, a homonym of *C. vicdani* Lan, 1978, renamed *C. sulcocastaneus* Kosuge, 1981.





victor Broderip, 1842.

Published in Proc. zool. Soc. Lond. pt. 10, no. 111: p. 54, not figured.

Lectotype (Finet & Caillez, inadvertently designated as holotype), in NHMUK, (32  $\times$  16 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a subspecies of *C. nobilis* Linnaeus, 1758, although a junior synonym of *C. vincoomnes* Lichtenstein, 1794, C. *vincoomnes* Lichtenstein, 1794 should be forgotten (nomen oblitum), (designated Geiger, 2003) and *C. victor* Broderip, 1842 should take precedence (nomen protectum), see Section 1 C (8).





#### victoriae Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 37, sp. 202, (published Dec), (1844, Proc. zool. Soc. Lond. pt. 11, no. 130: p. 172, not figured, published Jun).

Two syntypes were in collection G.P. Deshayes, possibly in UCBL, (figs  $40.5 \times 19 \& 37 \times 19 \text{ mm}$ ).

Type locality Mouth of Victoria River, New Holland, (Northern Territory, Australia), [dubious].

Nomenclatural status, an available name.





Figures in Conch. Icon.

(Textilia) vicweei Old, 1973.

Published in Veliger 16 (1): p. 58, figs 1 - 3 & 5.

Holotype in AMNH,  $(66 \times 31 \text{ mm})$ .

Type locality North West of Sumatra, Indonesia, (dubious), (trawled, sand & sponge, 50 - 60 fths).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





vidua Reeve, 1843.

Published in Conch. Icon. 1 (*Conus*): pl. 8, sp. 45, (published Apr), (Proc. zool. Soc. Lond. pt. 11, no. 125: p. 79 & no. 130: p. 169, not figured, published Dec).

Two syntypes in NHMUK, (fig. 45a, 72 x 41 mm. & fig. 45b, 47.5 x 26 mm).

Type locality Capul Is., Philippines, (on reefs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. bandanus Hwass, 1792.









Figured syntype 72 x 41 mm.

Figured syntype  $47.5 \times 26 \text{ mm}$ .

vikingorum Petuch, 1993 ? ("1992").

Published in Conchiglia 24 (265): p. 15, figs 20 & 21.

Holotype in CMNH,  $(37 \times 20 \text{ mm})$ .

Type locality off Puerto Colombia, Mouth of Magdalena River, Atlantico State, Colombia, (East coast), (35 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. daucus Hwass, 1792.





villepinii P. Fischer & Bernardi, 1857 ("1856").

Published in J. Conch. Paris, 5: p. 292, pl. 9, fig. 12, (1861, Monographie de Genre Conus: pl. 1, fig. 7).

Holotype was in collection Beau & Schramm, present whereabouts unknown,  $(32 \times 13 \text{ mm})$ .

Type locality Marie Galante, (Leeward Islands), (Beau & Schramm). Nomenclatural status, an available name.



Figure in Monographie.

vimineus Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Sup. pl. 7, sp. 269 (319). Holotype in NHMUK, (29 x 10.5 mm).

Type locality Cagayan, Mindanao Island, Philippines, (sandy mud, 25 fths).

Nomenclatural status, an available name.

Taxonomic status, a valid species





<u>vincomnes</u> variety nobilis Linnaeus) Lichtenstein, 1794. Published in Catalogus rerum naturalium rarissamarum 2: p. 23, no. 74, not figured.

Holotype wherabouts unknown, [presumably sold at auction], (dimensions not mentioned).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a senior synonym of *C. victor* Broderip, 1842, *C. nobilis var. vincoomnes* Lichtenstein, 1794 should be forgotten (nomen oblitum), (designated Geiger 2003) and *C. nobilis* Linnaeus subspecies *victor* Broderip, 1842 should take precedence (nomen protectum), see Section 1 C (8)

vinctus A. Adams, 1854 ("1853").

Published in Proc. zool. Soc. Lond. pt. 21, no. 255: p. 118, no. 12, not figured.

Holotype in NHMUK,  $(40 \times 21 \text{ mm})$ .

Type locality Australia, (Cuming).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. monachus* 

Linnaeus, 1758.





viola Cernohorsky, 1977.

Published in Nautilus, Philad. 91 (2): p. 72, figs 1 - 3. Lectotype of *C. violaceus* Reeve in NHMUK, (41 x 15 mm).

Type locality of *C. violaceus* Reeve Matnog, Luzon, Philippines, (on reefs).

Nomenclatural status, an available name, a new replacement name (nomen novum) for  $C.\ violaceus$  Reeve, 1844.





Lectotype of *C. violaceus* Reeve.

<u>violacea</u> (variety achatinus Hwass) Barros E Cunha, 1933.
Published in Mems Estud. Mus. zool. Univ. Coimbra 1 (71): p. 162, no. 138, not figured.

Two syntypes in MZUC, (70 x 38 & 67 x 35 mm), missing in 2011. Type locality India & China, (two different localities). Nomenclatural status, an available name, an invalid name, a homonym of C. violaceus Gmelin, 1791, of Link, 1807 and of Reeve, 1843.

Taxonomic status, a synonym of C. achatinus Hwass, 1792.

### violaceus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3391, no. 51, not figured.

Holotype representative (Kohn) figures in Martini, 1777, pl. 2, figs 18 & 19, (fig. 50 x 17 mm).

Type locality not known.

Nomenclatural status, an available name.





violaceus Link, 1807.

Published in Beschr. Nat.-Samml. Univ. 3: p. 106, not figured. Holotype (Kohn) in ZMUC,  $(45 \times 20 \text{ mm})$ .

Type locality not mentioned.

Nomenclatural status, an available name, an invalid name, a homonym of *C. violaceus* Gmelin, 1791.

Taxonomic status, a synonym of *C. glans* Hwass, 1792.





### violaceus Reeve, 1844.

Published in Conch. Icon. 1 (*Conus*): pl. 44, sp. 241, (published Feb), (Proc. zool. Soc. Lond. pt. 11, no. 130: p. 176, not figured, published Jun).

Lectotype (Cernohorsky) in NHMUK, (41 x 15 mm), (paralectotype  $36.5 \times 14$  mm, figured, in NHMUK).

Type locality Matnog, Luzon, Philippines, (on reefs).

Nomenclatural status, an available name, an invalid name, a homonym of *C. violaceus* Gmelin, 1791 and of Link, 1807, renamed *C. viola* Cernohorsky, 1977.

Taxonomic status, a synonym of C. viola Cernohorsky, 1977.





<u>violascens</u> (subspecies *figulinus* Linnaeus) Barros E Cunha, 1933. Published in Mems Estud. Mus. zool. Univ. Coimbra, 1 (7): p. 37, no. 29, not figured.

Two syntypes in MZUC,  $(48 \times 30 \& 42 \times 27 \text{ mm})$ .

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. loroisii Kiener, 1845 (1846).





Syntype 48 x 30 mm MZUC.

Syntype 42 x 27 mm MZUC.

## viperinus Lauer, 1986.

Published in Conchiglia 18 (212 - 213): p. 28, figs 1 - 3. Holotype in MNHN,  $(48.4 \times 25 \text{ mm})$ .

Type locality Island of Bohol, South Philippines, (on dead coral, 5 - 15 mtrs), corrected (Lauer 1990) to Sulu Archipelago, Philippines.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. omaria* Hwass, 1792.





virgatus Reeve, 1849.

Published in Conch. Icon. 1 (Conus): Emend. p. 1. (pl. 16, sp. 87).

Three syntypes in NHMUK, (56 x 29, 48 x 25 & 46.5 x 23 mm last probably figured).

Type locality Salango, Ecuador, (dredged, sandy mud, 12 - 16 fths).

Nomenclatural status, an available name.

Taxonomic status, a valid species.









Probable figured syntype in NHMUK

Syntype in NHMUK.


## virgineus Link, 1807.

Published in Beschr. Nat.-Samml. Univ. 3: p. 106, not figured. Type series whereabouts unknown, cited figures in Martini, 1773, pl. 63, figs 701 & 702, (fig. 701, 32 x 16 mm. & fig. 702, 29 x 17 mm).

Type locality not mentioned.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost type, no locality or description and unclear figures.



Cited figures in Martini.

virgineus (variety capitaneus Linnaeus) Wils, 1972.

Published in Family Conidae 8: p. 91, no. 156, not figured.

Specimen described in IRSN, (68.8 x 42.4 mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), described as a variety after 1960.

Taxonomic status, none, (a specimen of *C. capitaneus* Linnaeus, 1758).





Specimen described in IRSN.


## virgo Linnaeus, 1758.

Published in Syst. Nat. 10th ed. (1): p. 713, no. 253, not figured.

Lectotype representative (Kohn) figure in Rumphius, 1705, pl. 31, fig. E, (fig. 63  $\times$  30 mm).

Type locality not known, (1767 "Africano Oceano"), designated (Wils) Ambon, (Indonesia).

Nomenclatural status, an available name.



viridis Sowerby II, 1857.

Published in Thes. Conch. 3 (*Conus*), pt. 17: p. 20, no. 162, pl. 5 (191), fig. 102.

Lectotype representative (Wils) figure in T. C. pl. 5, fig. 102, (fig. 22 x 12.5 mm), (a specimen in NHMUK Type collection, (35.5 x 20.5 mm) is not part of type material and is a specimen of C. balteatus Sowerby, 1833).

Type locality not known, designated Hurghada, Red Sea, (Verbinnen & Wils).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. rattus Hwass, 1792.



viridulus Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 31, no. 9, not figured.

Holotype was in MNHN, present whereabouts unknown,  $(53 \times 27 \text{ mm})$ , cited figure in Chemnitz, 1788, pl. 139, fig. 1289,  $(48 \times 27 \text{ mm})$ . Type locality Australian Ocean, uncertain.

Nomenclatural status, an available name.

Taxonomic Status, a synonym (form) of *C. imperialis* Linnaeus, 1758.



Cited figure in Chemnitz.

visagenus Kilburn, 1974.

Published in Durban Mus. Novit. 10 (6): p. 81, pl. 1, figs 1 - 4. Holotype in NMSA, (31.2 x 23.3 mm).

Type locality East of Durban, Natal, South Africa, (180 fths). Nomenclatural status, an available name.

Taxonomic status, a valid species.





visseri Delsaerdt, 1990.

Published in Gloria Maris 29 (1): p. 1 & figs 1 - 4.

Holotype in NBC-ZMA,  $(8.7 \times 5.3 \text{ mm})$ .

Type locality Ka Lhim Beach, Patong Bay, Phuket, Thailand, (beached).

Nomenclatural status, an available name.

Taxonomic status, a juvenile of uncertain status (incertae cedis), probably a synonym(juvenile) of *C. coffeae* Gmelin, 1791.





vitellus "Solander's MSS" Dillwyn, 1817.

Published in Desc. Cat. Recent Shells 1: p. 390, no. 67, not figured.

Specimen named whereabouts unknown, (? mm).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), cited as a synonym of *C. vittatus* Hwass, 1792.

Taxonomic status, none, (a specimen of C. vittatus Hwass, 1792).

# vitifera Perry, 1811.

Published in Conchology: pl. 25, fig. 2.

Holotype whereabouts unknown, (fig. 81 x 41 mm).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, doubtful (nomen dubium), lost type, no locality, an inadequate description and an unclear figure.



vittatus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 704, no. 95, (1798, Tab. Enc. pl. 335, fig. 3).

Lectotype (Kohn) in MHNG,  $(39 \times 23 \text{ mm})$ .

Type locality Indian Ocean, [erroneous].

Nomenclatural status, an available name.

Taxonomic status, a valid species.





vitulinus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 648, no. 47, (1798, Tab. Enc. pl. 326, fig. 3).

Lectotype (Kohn) in MHNG,  $(50 \times 29 \text{ mm})$ .

Type locality Asiatic Ocean.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. planorbis* Born, 1778.





vividus Sowerby III, 1914.

Published in Ann. Mag. nat. Hist. 8 (14): p. 476, pl. 19, fig. 8. Holotype in NHMUK,  $(4.5 \times 2.5 \text{ mm})$ .

Type locality New Caledonia.

Nomenclatural status, an available name.

Taxonomical status, a juvenile of uncertain status (incertae cedis).





voluminalis "Hinds" Reeve, 1843.

Published in Conch. Icon. 1 (Conus): pl. 37, sp. 206.

Holotype was in collection Sir E. Belcher, present whereabouts unknown, (fig.  $37 \times 20 \text{ mm}$ ).

Type locality Straits of Malacca, (between Malaya and Sumatra), (Hinds).

Nomenclatural status, an available name.



Figure in Conch. Icon.

### voluminalis Hinds, 1844.

Described in The Zoology of the Voyage of "H.M.S. Sulphur", 2, Mollusca: p. 7, no. 4, pl. 1, figs 8 & 9.

Holotype was in collection Sir E. Belcher, present whereabouts unknown, (fig.  $48 \times 19 \text{ mm}$ ).

Type locality Straits of Malacca, (between Malaya & Sumatra), (in mud flats).

Nomenclatural status, an available name, an invalid name, a homonym of *C. voluminalis* Reeve, 1843.

Taxonomic status, an objective junior synonym of *C. voluminalis* Reeve, 1843, same type.





Figures in Hinds.

vulcanus Tenorio & Afonso, 2004.

Published in Visaya vol. 1, no. 2: p. 25, pl. 1.

Holotype in MNCM,  $(25.2 \times 14.5 \text{ mm})$ .

Type locality Porto Ferriera, East coast of Boavista, Cape Verde Islands, (rocky reef, large rock platforms with algae and sand, 1 - 3 mtrs).

Nomenclatural status, an available name.





vulpinus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 648, no. 48, (1798, Tab. Enc. pl. 326, figs 6 & 8).

Lectotype (Kohn) in MHNG,  $(51 \times 26 \text{ mm})$ .

Type locality Guinea, (? New Guinea).

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. planorbis* Born, 1778, (recognised later by Hwass).





vulpinus Schubert & Wagner, 1829.

Published in Neues Systematisches Conchylien-Cabinet 12: p. 56, pl. 222, fig. 3073.

Lectotype representative (Kohn) figure in Schubert & Wagner pl. 222, fig. 3073, (fig. 50 x 27 mm).

Type locality var. A, African Ocean, [erroneous], var. B, Guinea. (? New Guinea).

Nomenclatural status, an available name, an invalid name, a homonym of *C. vulpinus* Hwass, 1792.

Taxonomic status, a synonym of *C. vulpinus* Hwass, 1792, (a synonym of *C. planorbis* Born, 1778).

