

Section XYZ xanthicus to zylmanae

Copyright © 2011.R Filmer Conditions of use.

The content of this website is provided for personal and scientific use and may be downloaded for this purpose. It may not be used in whole or part for any commercial activity and publication of any of the content on the internet is limited to the Cone Collector website (www.TheConeCollector.com). Authors wishing to publish any of the pictures may do so on a limited basis but should inform M Filmer so that the original owner of the rights to the picture can be acknowledged. (mike@mfilmer.fsnet.co.uk)

Version 1.0 October 2011.

xanthicus Dall, 1910.

Published in Proc. U.S. natn. Mus. 38 (1741): p. 225, not figured.

Holotype in USNM, $(42 \times 22.5 \text{ mm})$.

Type locality off Guayamas, Mexico, (West coast), (sand, 71 $\,$ fths).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





xanthocinctus Petuch, 1980.

Published in Proc. biol. Soc. 99 (1): p. 13, figs 11 & 12.

Holotype in MORG, (47 x 22 mm).

Type locality Cape Frio, Rio de Janeiro State, Brazil, (trawled 100 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym of *C. clenchi* Martins, 1943, (a synonym of *C. lemniscatus* Reeve, 1849).





xicoi Röckel, 1987.


Published in Publcoes. Occ. Soc. port. malac. (9): p. 45, figs 11 & 12.

Holotype in NMSF, $(25.5 \times 14.7 \text{ mm})$.

Type locality Santiago Beach, 30 kms North of Luanda, Angola, (intertidal, rocks & sand).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





ximenes Gray, 1839.


Published in The Zoology of Beechey's Voyage, (London): p. 119, pl. 33, fig.2.

Lectotype (Kohn) in NHMUK, (40 x 19 mm).

Type locality Panama, (West coast).

Nomenclatural status, an available name.





yemenensis Bondarev, 1997.

Published in World Shells (23): p. 66 & figs.

Holotype in NHMUK, $(30.2 \times 12.5 \text{ mm})$.

Type locality Gulf of Aden, offshore Yemen, Sikha I area, (sandy silt, 50 - 60 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.





yemanjae van Mol, Tursch & Kempf, 1967.

Published in Ann. Inst. oceanogr. Univ. Recife, 45 (2): p. 251, fig. 15, & pl. 8, fig. 1.

Holotype in MNHN, (12.1 x 8.9 mm). Type locality off mouth of San Francisco River, Brazil, (dredged, 63 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. selenae van Mol, Tursch & Kempf, 1967, (p. 250), recognized by the three authors in 1971.





(Rhizoconus) yoshioi Azuma, 1973.

Published in Venus, Kyoto, 32 (1): p. 10, pl. 1, fig. 2.

Holotype in BRIT, $(42.5 \times 22.5 \text{ mm})$.

Type locality off Kirimezaki, Kii Peninsula, Japan, (20 - 30 fths).

Nomenclatural status, an available name.

Taxonomic status, a synonym (form) of C. sazanka Shikama, 1970.





zandbergeni Filmer & Moolenbeek, 2010.

Published in Miscellanea Malacologica 4(5): p.75 & figs.

Holotype in NBC-ZMA, $(38.10 \times 19.30 \text{ mm})$.

Type locality Sulu Sea, The Philippines (in sand, 10-25 meters).

Nomenclatural status, an available name.





zapatosensis Röckel, 1987.

Published in Spixiana 10 (3): p. 288, figs 4, 5, 8 & 12.

Holotype in ZSM, $(40.1 \times 19.1 \text{ mm})$.

Type locality Zapatos Island, Negros, Philippines, (100 mtrs). Nomenclatural status, an available name.

Taxonomic status, a valid species.





zealandicus Hutton, 1873.

Published in Catalogue of Marine Molluscs of New Zealand, (collection Mus. & Geol. Survey): p. 23, not figured. Holotype in NMNZ, (14.7 x 8 mm).

Type locality Bay of Islands, New Zealand, [erroneous, specimen must have reached New Zealand with a ship].

Nomenclatural status, an available name.

Taxonomic status, a synonym of C. ventricosus Gmelin, 1791.





zebra Lamarck, 1810.

Published in Annls. Mus. Hist. nat. Paris, 15: p. 273, no. 82, not figured.

Holotype (Kohn) in MNHN, $(26.6 \times 14.5 \text{ mm})$.

Type locality Asiatic Ocean.

Nomenclatural status, an available name.

Taxonomic status, a valid species.





zebra Mawe, 1823.

Published in The Linnean System of Conchology: p. 87, not figured.

Specimen named whereabouts unknown, (? mm).

Locality New Guinea.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none, (presumably a specimen of $C.\ zebra$ Lamarck, 1810).

zebroides Kiener, 1845 (1848).

Published in Coq. Viv. 2: pl. 105, fig. 5, (1849 (1850), Coq. Viv. 2: p. 257, no. 224).

Holotype was in collection Largilliert, present whereabouts unknown, (45 x ? mm), holotype representative (Rolán & Röckel) figure in C. V. pl. 104, fig. 2, (fig. 45 x 26 mm)

Type locality not mentioned, designated (Rolán & Röckel) Lucira Bay, Angola.

Nomenclatural status, an available name. Taxonomic status, a valid species.



"<u>zerokini</u>" (form marmoreus crosseanus Bernardi) Prigent, 1984. Published in Rossiniana (22): p. 19, not figured. Specimen named whereabouts unknown, (? mm). Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), no intention to introduce a new name, named but not described, indicated or figured and named as a form after 1960. Taxonomic status, none, (a specimen of *C. crosseanus* Bernardi, 1861, a synonym of *C. marmoreus* Linnaeus, 1758).

zeylanicus Gmelin, 1791.

Published in Syst. Nat. 13th ed. 1 (6): p. 3389, no. 41, not figured.

Lectotype representative (Kohn) figure in Martini, 1777, pl. 2, fig. 20, (fig. $44 \times 23 \text{ mm}$).

Type locality not known.

Nomenclatural status, an available name.

Taxonomic status, a valid species.



ziczac Megerle von Mühlfeld, 1816.

Published in Mag. Ges. naturf. Fr. Berl. 8 (1): p. 4, pl. 1, fig. 3.

Holotype in NHMW, $(8.2 \times 4.6 \text{ mm})$.

Type locality Mediterranean Sea, [erroneous].

Nomenclatural status, an available name.



zonatus "S" Lightfoot, 1786.

Published in Cat. Portland Mus: p. 18, no. 401 & p. 146, no. 3216, not figured.

Two specimens named sold at auction, present whereabouts unknown, ($? \ mm$).

Locality not mentioned.

Nomenclatural status, an unavailable name (nomen nudum), named but not described or figured.

Taxonomic status, none, (probably a specimen of C. zonatus Hwass, 1792).


zonatus Hwass in Bruguière, 1792.

Published in Enc. Méth. 1: p. 613, no. 9, (1798, Tab. Enc. pl. 318, fig. 4).

Lectotype representative (Kohn) figure Tab. Enc. pl. 318, fig. 4, $(fig. 57 \times 32 \text{ mm})$.

Type locality Asiatic Ocean.

Nomenclatural status, an available name.



(Leptoconus) zukiae Shikama, 1979.

Published in $\overline{\text{Sci. Rep.}}$ Yokosuka Cy. Mus. (26): p. 3, pl. 1, figs 9 & 10.

Holotype in NSMT, $(70.5 \times 37.3 \text{ mm})$.

Type locality off the Philippines.

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of *C. mustelinus* Hwass, 1792.





zulu Petuch, 1979.

Published in Nemouria (23): p. 19, figs 28 - 31.

Holotype in DMNH, ((62 x 35 mm).

Type locality off mouth of Umfolozi River, Zululand coast, Natal, South Africa, (app. 40 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a synonym (colour form) of $C.\ betulinus$ Linnaeus, 1758.





(Magelliconus) zylmanae Petuch, 1998.

Published in Conchiglia 30 (287): p.28, & figs 6 & 7.

Holotype in FMNH, (22 x 13 mm).

Type locality northern Great Bahama Bank, off Paradise Isl.,

North of New Providence Isl., Bahamas, (hardpan bottom, dense algae & strong currents, 6 - 7 mtrs).

Nomenclatural status, an available name.

Taxonomic status, a valid species.



